



CHALMERS
UNIVERSITY OF TECHNOLOGY

Fission of highly excited nuclei investigated in complete kinematic measurements

Downloaded from: <https://research.chalmers.se>, 2026-04-05 15:38 UTC

Citation for the original published paper (version of record):

Rodriguez-Sanchez, J., Benlliure, J., Taieb, J. et al (2013). Fission of highly excited nuclei investigated in complete kinematic measurements. EPJ Web of Conferences, 62.
<http://dx.doi.org/10.1051/epjconf/20136207009>

N.B. When citing this work, cite the original published paper.

Fission of highly excited nuclei investigated in complete kinematic measurements

J.L. Rodríguez-Sánchez¹, J. Benlliure¹, J. Taieb³, H. Álvarez-Pol¹, L. Audouin², Y. Ayyad^{1,a}, G. Bélier³, G. Boutoux³, E. Casarejos⁴, A. Chatillon³, D. Cortina-Gil¹, T. Gorbinet³, A. Heinz⁵, A. Kelić-Heil⁶, N. Kurz⁶, B. Laurent³, J.-F. Martin³, C. Paradela¹, E. Pellereau³, B. Pietras¹, A. Prochazka⁶, D. Ramos¹, C. Rodríguez-Tajes⁷, D. Rossi⁶, H. Simon⁶, L. Tassan-Got², J. Vargas¹, and B. Voss⁶

¹Universidad de Santiago de Compostela, 15782 Santiago de Compostela, Spain

²Institut de Physique Nucléaire d'Orsay, 91406 Orsay, France

³CEA DAM, 91297 Arpajon, France

⁴Universidad de Vigo, 36200 Vigo, Spain

⁵Chalmers University of Technology, 412 96 Gothenburg, Sweden

⁶GSI Helmholtzzentrum für Schwerionenforschung GmbH, 64291 Darmstadt, Germany

⁷Grand Accélérateur National d'Ions Lourds, 14076 Caen, France

Abstract. Fission is an extremely complex mechanism that requires a dynamical approach to describe the evolution of the process in terms of intrinsic and collective excitations of the nuclear constituents. In order to determine these effects a complex experimental setup was mounted at GSI, which allowed us for the first time the full identification in charge and mass of all fission fragments thanks to a magnetic separation and the use of the inverse kinematic technique. Moreover, we also measured the neutron multiplicities and the light-charged particles emitted in coincidence with fission. These complete kinematic measurements will be used to define sensitive observables to dissipative and transient effects in fission. In this manuscript we present the first results for the total fission cross sections.

1. Introduction

In 1940 the first description of the nuclear fission was proposed by Bohr and Wheeler [1] as a purely statistical model. At the same time and with the same purpose, Kramers [2] proposed a dynamical description of the fission process based on the coupling between internal and collective degrees of freedom through a dissipation parameter (β) which reduces the fission decay width respect to the statistical model. Both descriptions were under discussion during four decades, however, in the 80's different measurements of pre-scission neutron multiplicities [3] provided values much larger than expected from the predictions using the Bohr-Wheeler's statistical model, which indicated the existence

^aPresent address: Research Center for Nuclear Physics, Osaka University, Ibaraki, Osaka 567-0047, Japan




Figure 1. Schematic drawing of the experimental set-up used to investigate the reaction $^{208}\text{Pb} + p$ in inverse kinematics.

of a dissipative effect in the fission. The introduction of the dissipation implies that the system needs time to populate the phase space and reach the equilibrium. Under this idea, Grangé [4] investigated the dependence of the fission decay width with the time solving the Fokker-Planck equation (FPE). The solution showed that the fission is highly suppressed at the onset of the process, requiring a certain transient time to reach the stationary fission rate given by Kramers. These dynamical effects appear on fissioning systems with high excitation energy, low angular momentum and near the ground-state deformation [4]. For this reason, we investigated the reaction ^{208}Pb impinging onto a liquid hydrogen target at energies between 370 and 650 A MeV. Moreover the use of the inverse kinematic technique allowed us to measure all particles emitted in coincidence with the fission fragments providing a complete access.

2. Experiment

The fission setup situated at the Cave C at GSI is shown in Figure 1 where a small plastic-scintillator ($50 \times 30 \text{ mm}^2$) was used as trigger of the data acquisition. The characterization of the beam is made with an ionization chamber (MUSIC) [5], that provides the energy loss with high-resolution, and a Time Projection Chamber (TPC) [6], which provides the horizontal X and vertical Y position of the ions before the target with a resolution of $300 \mu\text{m}$ (FWHM). A cylindrical target (11.24 mm long and 30 mm diameter) was filled with liquid-hydrogen which was isolated by two windows consisting of aluminized-mylar foils of $35 \mu\text{m}$. After the target, the identification of the fission fragments was made with a double ionization chamber (Twin MUSIC) (active volume of 50 cm long, 20 cm high and 20 cm wide) [7], which measures the nuclear charge and the X coordinate of both fission fragments with a resolution of $50 \mu\text{m}$ (FWHM) and two multi-wire proportional chambers (MWPC), that provides the X and Y coordinates of the fission fragments with a resolution (FWHM) of $200 \mu\text{m}$ and 1 mm [8], respectively. These position measurements allowed us to reconstruct the path-length and the $B\rho$ of both fission fragments. Moreover, a ToF Wall [9] composed of 28 plastic-scintillator was placed behind the second MWPC to measure the time-of-flight with a resolution around 40 ps (FWHM). In addition, two pipes filled with helium gas were mounted between the target and the double ionization chamber and between the magnet and second MWPC to reduce the energy and angular straggling of the fission fragments. To detect the light-charged particles in coincidence with the fission a segmented plastic-scintillator wall was placed around the first He pipe in front of the double ionization chamber. Finally, the neutron detector LAND [10] (1 m long, 2 m high and 2 m wide) was used to determine the neutron multiplicities.




Figure 2. Total fission cross sections measured in the present work in comparison to previously measured data of the reactions ${}^{208}\text{Pb} + p$ and $p + {}^{nat}\text{Pb}$.

3. Results

The total fission cross sections for proton-induced fission of ${}^{208}\text{Pb}$ were measured at 370, 500 and 650 A MeV. Measurements were performed with a hydrogen target of (86 ± 1) mg/cm². The number of projectiles was determined from the reaction trigger provided by the small plastic-scintillator and the MUSIC while the number of fission events was determined using the double ionization chamber. Their statistical uncertainties given by Poisson statistics according to the observed number of counts are below 1% and are negligible in the fission cross-section uncertainties. The main contributions to the uncertainties are attributable to uncertainties of the target thickness, attenuation of the beam intensity in the target, fission events lost on the pipe and the detection efficiency of the TWIN chamber. The largest corrections are attributable to the attenuation of the beam intensity in the target considering the attenuation to half the target (values obtained for the energies 370, 500 and 650 MeV were 3.6%, 3.8% and 4.0%, respectively.), the correction of the fission events lost on the pipe and the detection efficiency of the TWIN chamber (around 88%). The secondary reactions were also calculated with the nuclear reaction rate in half the target obtaining values smaller than 2.2% for all energies. Moreover, the fission fragments could undergo nuclear reactions between the target and the TWIN chamber and not be counted as fission events. This corresponding probability was calculated with the nuclear reaction rate of the different layers of matter, obtaining values below 0.5% for three energies.

Our results are shown in Figure 2 (open squares). In this figure we also show similar data found in the literature for the reactions $p + {}^{nat}\text{Pb}$ and ${}^{208}\text{Pb} + p$ at different energies. As can be seen in the figure, our data have in general better accuracy than any of the previous ones. They seem consistent with the measurements between 50 and 200 MeV, which were performed by V.S. Bychenkov [11] (full squares), if one looks at the downward trend and also with measurement performed by T. Enqvist [12] (full triangle up) at the FRS if one looks at the upward trend. One can also see a discrepancy at 1 GeV between the measurements performed by T. Enqvist [12] (full triangle up) and M. Gloris [13] (open diamond) but our results are only consistent with the Enqvist's measurement. Moreover, our results are in agreement with the values reported by E. Hagebo [14] (open cross) and R. Brandt [15] (star) who measured in direct kinematics the reaction $p + {}^{nat}\text{Pb}$ at 600 and 590 MeV, respectively. However,

the cross section obtained by B. Fernández [16] (full circle) at the FRS has a non-negligible deviation with respect to our results, which can be due to the limited acceptance of the spectrometer requiring large corrections. Also, the measurements performed by A.A. Kotov [17] (full triangles down) have a deviation, but these were performed in direct kinematics with a proton beam impinging on a ^{nat}Pb target, which could prevent fission fragments to escape from the target reducing the total fission cross section if one does not apply the accurate corrections. However, these measurements confirm the upward trend of our results. On the other hand our value of fission cross section at 500 MeV is in agreement with the measurement performed by K.-H. Schmidt [18] (full star) in a similar setup at GSI. Moreover the fission cross section at 370 MeV is in agreement with the results performed by V.A. Konshin [19] (open circles).

4. Summary and conclusions

In this work we have presented the first results for proton induced fission of ^{208}Pb in inverse kinematics at 370, 500 and 650 A MeV. The combination of the inverse kinematic technique with a highly efficient detection set-up made it possible to obtain the total fission cross sections with high accuracy. The new data have made possible to clarify the previous controversial results obtained in different experiments.

References

- [1] N. Bohr and J.A. Wheeler, *Phys. Rev.* **56**, 426 (1939)
- [2] H.A. Kramers, *Physika VII* **4**, 284 (1940)
- [3] A. Gavron *et al.*, *Phys. Rev. Lett.* **48**, 835 (1982)
- [4] P. Grangé, L. Jun-Qing and H. A. Weidenmüller, *Phys. Rev. C* **27**, 2063 (1983)
- [5] M. Pfüzner *et al.*, *Nucl. Instr. Methods B* **86**, 213 (1994)
- [6] R. Janik, A. Prochazka *et al.*, *Nucl. Instrum. Methods A* **640**, 54-57 (2011)
- [7] B. Voss, “A low-mass Multi-Sampling Ionization chamber for fission studies”, Nuclear Science Symposium and Medical Imaging Conference (NSS/MIC), 1767-1770 (2011)
- [8] L. Audouin, Private communication
- [9] A. Bail, “Time of flight measurement on the SOFIA experiment”, *Advancements in Nuclear Instrumentation Measurement Methods and their Applications (ANIMMA)*, 1-4 (2011)
- [10] Th. Blaich *et al.*, *NIM A* **314**, 136-154 (1992)
- [11] V.S. Bychenkov *et al.*, *Soviet Journal of Nuclear Physics*, Vol. **17**, 496 (1973)
- [12] T. Enqvist *et al.* *Nucl. Phys. A.* **686**, 481 (2001)
- [13] M. Gloris, *et al.*, *Nucl. Inst. Meth. A* **463** 593 (2001)
- [14] E. Hagebo and T. Lund. *J. inorg. nucl. Chem.* **37** 1569 (1975)
- [15] R. Brandt *et al.* *CERN* **71**, 2 (1971)
- [16] B. Fernández-Dominguez *et al.* *Nucl. Phys. A* **747**, 227 (2005)
- [17] A.A. Kotov *et al.* *Phys. Rev. C* **74**, 034605 (2006)
- [18] K.-H. Schmidt *et al.* *Phys. Rev. C* **87**, 034601 (2013)
- [19] V.A. Konshin *et al.*, *Yadernaya Fizika* Vol. **2**, 682 (1965)