



CHALMERS
UNIVERSITY OF TECHNOLOGY

Strategies and challenges with the microbial conversion of methanol to high-value chemicals

Downloaded from: <https://research.chalmers.se>, 2026-04-06 05:57 UTC

Citation for the original published paper (version of record):

Zhan, C., Li, X., Yang, Y. et al (2021). Strategies and challenges with the microbial conversion of methanol to high-value chemicals. *Biotechnology and Bioengineering*, 118(10): 3655-3668.
<http://dx.doi.org/10.1002/bit.27862>

N.B. When citing this work, cite the original published paper.



The Smarter Solution

Flexible, powerful, industrial

Many engineering parameters need to be considered to establish a successful mAbs process. The BioFlo® 320 offers a wide range of options to meet your needs. It combines industrial design, flexibility between interchangeable autoclavable and single-use bioreactors, and universal gas control strategy for your applications.

- > **New Scale Up Assist** software:
Fast and easy calculation of important process parameter based on P/V or constant tip speed
- > Compatible with the BioBLU® Single-use Bioreactor portfolio
- > **Scalable:** Extensive working volume range of 250 mL – 40 L
- > **Efficient:** Multi-unit control of up to eight systems from a single interface



www.eppendorf.com/BioFlo320

REVIEW

Strategies and challenges with the microbial conversion of methanol to high-value chemicals

Chunjun Zhan^{1,2,3,4}  | Xiaowei Li^{4,5} | Yankun Yang^{1,2,3}  | Jens Nielsen^{4,5,6} |
 Zhonghu Bai^{1,2,3} | Yun Chen^{4,5} 

¹National Engineering Laboratory for Cereal Fermentation Technology, Jiangnan University, Wuxi, China

²The Key Laboratory of Industrial Biotechnology, School of Biotechnology, Ministry of Education, Jiangnan University, Wuxi, China

³The Key Laboratory of Carbohydrate Chemistry and Biotechnology, School of Biotechnology, Ministry of Education, Jiangnan University, Wuxi, China

⁴Department of Biology and Biological Engineering, Chalmers University of Technology, Gothenburg, Sweden

⁵Novo Nordisk Foundation Center for Biosustainability, Chalmers University of Technology, Gothenburg, Sweden

⁶BioInnovation Institute, Copenhagen N, Denmark

Correspondence

Zhonghu Bai, National Engineering Laboratory for Cereal Fermentation Technology, Jiangnan University, 214122 Wuxi, China.
 Email: baizhonghu@jiangnan.edu.cn

Yun Chen, Department of Biology and Biological Engineering, Chalmers University of Technology, Kemivägen 10, SE412 96 Gothenburg, Sweden.
 Email: yunc@chalmers.se

Funding information

National Natural Science Foundation of China, Grant/Award Numbers: 21908077, 31570034; Novo Nordisk Fonden, Grant/Award Number: NNF10CC1016517

Abstract

As alternatives to traditional fermentation substrates, methanol (CH₃OH), carbon dioxide (CO₂) and methane (CH₄) represent promising one-carbon (C1) sources that are readily available at low-cost and share similar metabolic pathway. Of these C1 compounds, methanol is used as a carbon and energy source by native methylotrophs, and can be obtained from CO₂ and CH₄ by chemical catalysis. Therefore, constructing and rewiring methanol utilization pathways may enable the use of one-carbon sources for microbial fermentations. Recent bioengineering efforts have shown that both native and nonnative methylotrophic organisms can be engineered to convert methanol, together with other carbon sources, into biofuels and other commodity chemicals. However, many challenges remain and must be overcome before industrial-scale bioprocessing can be established using these engineered cell refineries. Here, we provide a comprehensive summary and comparison of methanol metabolic pathways from different methylotrophs, followed by a review of recent progress in engineering methanol metabolic pathways *in vitro* and *in vivo* to produce chemicals. We discuss the major challenges associated with establishing efficient methanol metabolic pathways in microbial cells, and propose improved designs for future engineering.

KEYWORDS

energy supply, intermediate recycling, methanol, methylotrophs, redox cofactor

1. INTRODUCTION

Rapid population growth and economic developments in the world are imposing an increased demand for food, materials, and energy. As traditional supply routes are unable to meet this demand, there is much interest in evaluating alternative supply routes for these resources. Production of biofuels and high-value commodity chemicals via microbial fermentation represents a promising strategy to provide materials and energy in the world market (Luo et al., 2019; Nielsen & Keasling, 2016; Nielsen et al., 2014). However, traditional fermentation substrates mainly comprise of sugars, such as glucose, which has raised concerns about food security and biodiversity (Naik et al., 2010) (Figure 1a). Therefore, exploring the use of renewable and nonfood carbon sources is becoming extremely important for industrial biomanufacturing. CO₂ and natural gas (consisting primarily of CH₄) are two of the most important and attractive carbon sources owing to their high abundance (Haynes & Gonzalez, 2014; Xu et al., 2018). The process of converting CO₂ (e.g., metal catalysis, Fujiwara et al., 2019; and electrocatalysis, Magdesieva et al., 2002) to value-added products is promising in the sense that it will not only relieve the greenhouse effect, but also shorten the cycle of organic compound production via plant-based photosynthesis (Liu et al., 2020; Xiaoding & Moulijn, 1996) (Figure 1b). In addition, CO₂/CH₄-based autotrophic microorganisms could directly convert CO₂/CH₄ into biofuels and chemicals (Long et al., 2018), which greatly promotes CO₂/CH₄-based microbial cell factory construction (Savakis & Hellingwerf, 2015). However, a major challenge in harvesting CO₂ and CH₄ as carbon sources for microbial fermentation is that they cannot be fixed efficiently by most microbes, owing to their

gaseous form, thermodynamic instability (Naik et al., 2010), high energy-cost (Birdja et al., 2019), low catalytic activity of Rubisco (Erb & Zarzycki, 2016) and requirement for additional metabolic supplements such as ribulose-1,5-bisphosphate (RuBP) (Budzianowski, 2012; Erb & Zarzycki, 2018; B. Hu et al., 2013; G. Hu et al., 2018; Savakis & Hellingwerf, 2015). In contrast, methanol, a liquid one-carbon compound, is considered a promising C1 feedstock owing to its increasing availability (via conversion of natural gas) (Latimer et al., 2018; Linton & Niekus, 1987; Sperling, 2007), a predicted decrease in price (Bertau et al., 2014; Pfeifenschneider et al., 2017), and relatively low energy-cost compared with CO₂ and CH₄ (Schrader et al., 2009; Whitaker et al., 2015). Moreover, some methanol utilization pathways share similarities with CO₂ fixation pathways, such as the C5 regeneration pathway (Figure 1b) (Antonovsky et al., 2016; Cox & Quayle, 1975; Jakobsen et al., 2006). Therefore, a better understanding of methanol utilization pathways would not only benefit methylotrophic microorganism construction, but also provide a chassis for CO₂ and other C1 source utilization (Saeidi et al., 2014) (Figure 1b).

Native methylotrophs can grow on methanol as a single carbon and energy source, and include both methylotrophic bacteria, such as *Bacillus methanolicus* (Arfman et al., 1992; De Vries et al., 1992) and methylotrophic yeast, such as *Pichia pastoris* (Couderc & Baratti, 1980; Ellis et al., 1985). These native methylotrophs represent natural platforms for biotechnological engineering of C1 compound utilization, however the genetic tools for engineering these organisms are not as efficient as those for model organisms such as *Saccharomyces cerevisiae* and *Escherichia coli*. Moreover, drawbacks including insufficient accumulation of metabolic intermediates and




FIGURE 1 The traditional one-carbon cycle and one-carbon compound utilization for establishment of sustainable circular carbon economy. (a) The CO₂ cycle in nature. Plants convert CO₂ into different sources, that is, food and energy, to support human society development. Excessive human activity releases large amount of CO₂ exceeding the CO₂ sequestration speed by plants and leads to accumulation of CO₂ in atmosphere causing “greenhouse effect.” (b) C1 sources (i.e., CO₂ and methanol) utilization by microorganisms in biorefinery for production of chemicals and biofuels, allowing for a sustainable circular carbon economy. One-carbon feedstock (CO₂-based or methanol-based) can be used by microorganisms in a biorefinery process to replace fuels and chemicals that produced today by petroleum-refinery process. 1,3BPGA, 1,3 bisphosphoglycerate; 3PGA, 3-phosphoglycerate; DHA, dihydroxyacetone; DHAP, dihydroxyacetone phosphate; F6P, fructose 6-phosphate; G3P, glyceraldehyde-3-phosphate; H6P, hexose 6-phosphate; Pyr, pyruvate; Ru1,5BP, ribulose 1,5-bisphosphate; Ru5P, ribulose-5-phosphate; Xu5P, xylulose-5-phosphate [Color figure can be viewed at wileyonlinelibrary.com]

low carbon yield due to insufficient methanol catabolism (Jorda et al., 2014), further prohibit the use of these organisms as cell factories to produce secondary metabolites. In addition, genes related to methanol utilization pathways are tightly regulated by methanol and repressed by other carbon sources in co-substrate media, and the molecular mechanism behind this is still unclear (Zhan et al., 2017). Energy generation from methanol as a substrate is insufficient to support all biological processes, as ^{13}C labeling results show that most of the methanol assimilated is used to produce energy for methanol assimilation itself (Jordà et al., 2012). All of these factors make it necessary to take a two-step strategy when using *P. pastoris* for fermentations, whereby biomass formation is supported in the first step by a traditional carbon source (e.g., glycerol), and methanol is only used as a substrate to maintain biomass in the second step (Bahrami et al., 2008; Zhan et al., 2017). These drawbacks indicate that methanol is not an ideal carbon and energy source for fermentations using *P. pastoris*.

An alternative approach to utilize methanol is to develop synthetic methylotrophs that can efficiently metabolize methanol either as a sole carbon and energy source, or as co-substrates with other carbohydrates. To do this, metabolic engineering and synthetic biology strategies were widely used in both native and synthetic methylotrophs to optimize methanol metabolic pathways. In this review, we first provide a comprehensive summary and comparison between different methanol metabolic pathways (Table 1). Then, we discuss recent efforts toward understanding and engineering native and synthetic methylotrophs to produce biofuels and high-value commodity chemicals. Finally, we discuss remaining challenges and propose modular strategies to improve methanol utilization by synthetic methylotrophs.

2. METHANOL METABOLIC PATHWAYS IN METHYLOTROPHS

Methanol metabolic pathways can be divided into two modules (Figure 2) in almost all methylotrophs. The first is a methanol oxidation module, in which methanol is enzymatically converted to formaldehyde (HCHO). The second is a formaldehyde assimilation module, where formaldehyde could be assimilated into central carbon metabolism via one of three pathways: (1) the xylulose monophosphate (XuMP) cycle, (2) the ribulose monophosphate (RuMP) cycle, or (3) the serine cycle. Of note, besides the main fate of assimilation, formaldehyde can also be oxidized to CO_2 in dissimilation pathways (W. Zhang et al., 2017). In this process, the carbon in formaldehyde is released as CO_2 , but the proton generated is used to supply energy for biological processes.

The first module of methanol oxidation can be catalyzed by three methanol oxidoreductases, differentiated by their electron acceptor: (i) pyrroloquinoline quinone (PQQ)-dependent methanol dehydrogenase (PMDH), (ii) NAD^+ -dependent methanol dehydrogenase (NMDH) and (iii) O_2 -dependent alcohol oxidase (AOX). Among these, PMDH is the most complex as it requires the PQQ

prosthetic group to transfer electrons to cytochrome c. In *Methylobacterium extorquens*, the expression of up to 15 genes are required to realize its function (M. Zhang & Lidstrom, 2003; W. Zhang et al., 2017). Even in *betaproteobacteria* with monomeric PMDH (Kalyuzhnaya et al., 2008), at least 4–7 additional enzymes are still required to synthesize PQQ, which additionally requires molecular oxygen (Velterop et al., 1995). Use of the PMDH circuit in synthetic methylotrophs construction has therefore been limited (Davidson, 2001).

The second methanol oxidoreductase, NMDH, has a lower enzyme specificity towards methanol (K_m , 170 mM) compared to PMDH (20 μM) (Kalyuzhnaya et al., 2008; Sheehan et al., 1988; Wu et al., 2016), although incorporation of its activator protein (ACT) dramatically promotes its affinity towards methanol (9 mM) in vitro (Arfman et al., 1997; Ochsner et al., 2014). ACT acts by removing the nicotinamide mononucleotide (NMN) moiety of the NMDH-bound NAD^+ (Arfman et al., 1997; Wu et al., 2016), which increases the affinity of NMDH for methanol and NAD^+ (Arfman et al., 1991). Therefore, this simple two-component system is much easier to construct than the PMDH complex (Velterop et al., 1995). However, NMDH mainly exists in thermophilic Gram-positive methylotrophs (such as *B. stearothersophilus* and *B. methanolicus*) and has a high Gibbs free energy ($\Delta_r G' = +34.2 \text{ kJ mol}^{-1}$) (Whitaker et al., 2015). This means that the recombinant strains harboring NMDH must be grown at very high temperatures (45–55°C) to facilitate methanol oxidation, which is not feasible for most microorganisms.

The third methanol oxidoreductase, AOX, is mainly found in eukaryotic methylotrophs such as *P. pastoris*. Although this oligomeric flavoenzyme alcohol oxidase has relatively low affinity for both methanol and oxygen, its Gibbs energy is much lower ($\Delta_r G' = -99.2 \text{ kJ mol}^{-1}$) than the other two enzymes (Cereghino & Cregg, 2000; W. Zhang et al., 2017), which makes it more achievable for aerobic biological processes under standard conditions (Whitaker et al., 2015). To compensate for the low affinity towards methanol, *P. pastoris* expresses extremely high levels of AOX1, controlled by a strong methanol inducible promoter P_{AOX1} , which results in AOX accounting for more than 30% of total soluble protein (Inan & Meagher, 2001). However, as the overall standard enthalpy change for the reaction catalyzed by AOX is negative ($\Delta H_o = -112.41 \text{ kJ/mol}$), heat would be released under aerobic conditions, which may be problematic during scale-up fermentation (Jungo et al., 2007; Krainer et al., 2012). Moreover, its proton utilization efficiency is lower than both PMDH or NMDH, owing to the formation of H_2O_2 instead of NADH.

Whereas methanol assimilation using PMDH and AOX are limited to aerobic conditions, NMDH can potentially be used in both aerobic and anaerobic conditions. Some studies have shown that the thermophilic (55°C) anaerobic conversion of methanol to acetate could be realized by the addition of bicarbonate (Paulo et al., 2003). Indeed, microorganisms using NMDH to oxidize methanol has been explored under anaerobic conditions for liquid fuel production (<http://www.coskata.com>). However, the thermodynamic constraints on NMDH would require an optimal growth temperature of

TABLE 1 Comparison of different methanol pathways

Module name ¹	Substrates	Products	ATP equivalents	NADH equivalents	CO ₂ equivalents	Δ_rG' (kJ/mol) ^{a,d}	Energetic efficiency (or microbial conversion) (Cotton et al., 2020)	Location	Cofactor	KM (mM)	Biomass productivity gCDW/l/h	Temperature optima (°C)
First module												
NMDH	3 CH ₃ OH	3 HCHO	0	+3	0	+34.2 (Whitaker et al., 2015)		Cytoplasm	ACT, NAD ⁺	20	-	55
AOX	3 CH ₃ OH	3 HCHO	0	0	0	-99.2 (Whitaker et al., 2015)		Peroxisome	PYC, O ₂ , FAD	0.47	-	37
PMDH	3 CH ₃ OH	3 HCHO	0	0	0	-24.8 (Whitaker et al., 2015)		Cytoplasm	PQQ, Ca ²⁺	0.02	-	37
Second module												
RuMP cycle	3 HCHO	1 Acetyl-CoA	+1	+2	+1 ^c	-38.3 ^d	37%-49%	Cytoplasm	RuBP	-	28.4 (Bozdag, 2013)	
XuMP cycle	3 HCHO	1 Acetyl-CoA	-1	+2	+1 ^c	-78.4 ^d	31%-36%	peroxisome (Rußmayer et al., 2015)	Xu5P	-	-	
Serine cycle	3 HCHO	3 Acetyl-CoA	-6	-6	-3 ^c	-92.0 ^d	32%-37%	Cytoplasm	Serine, CH ₂ =H ₄ F	-	1.2-3.6 (Bozdag, 2013)	

Note: All calculations in the first (second) module are based on converting methanol (formaldehyde) into formaldehyde (acetyl-CoA).

^aStandard conditions are 25°C.

^bThe K_M values are obtained from BRENDA (<https://www.brenda-enzymes.org/>).

^cCO₂ equivalents did not take dissimilation pathway into consideration.

^d Δ_rG' in three different methanol utilization pathways were predicted by eQuilibrator (<http://equilibrator.weizmann.ac.il/>), pH 7, ionic strength = 0.1 M.




FIGURE 2 Methanol metabolic pathways in methylotrophs. AOX, alcohol oxidase; CAT1, catalase; DAK, dihydroxyacetone kinase; DAS, dihydroxyacetone synthase; DHA, dihydroxyacetone; DHAP, dihydroxyacetone phosphate; EMP, Embden-Meyerhof-Parnas pathway; F1,6P, fructose-1,6-bisphosphate; F6P, fructose-6-phosphate; FBA, fructose-1,6-bisphosphate aldolase; FBP, fructose-1,6-bisphosphatase; GAP, glyceraldehyde-3-phosphate; H6P, D-arabino-3-hexulo-6-phosphate; HPS, hexulose phosphate synthase; MDH, methanol dehydrogenase; OAA, oxaloacetate; PEP, phosphoenolpyruvate; PHI, phosphohexulose isomerase; PPP, pentose phosphate pathway; Ru5P, ribulose-5-phosphate; Xu5P, xylulose-5-phosphate [Color figure can be viewed at wileyonlinelibrary.com]

45–55°C, which is challenging for most microorganisms. Alternatively, HCHO must be kept at exceptionally low concentrations (<50 μM) (Woolston, King, et al., 2018). Such a low concentration of HCHO would however pose a challenge for one of the downstream enzymes in formaldehyde assimilation, which must then be engineered to have very strong substrate affinity towards HCHO.

The second module, HCHO assimilation, can occur via three possible routes: the serine pathway, the XuMP pathway, and the RuMP pathway. The serine pathway is the most carbon-efficient among the three pathways, as it can fix 3 mol of CO_2 along with 3 mol HCHO to form 3 mol acetyl-CoA. However, it is also the most energy-expensive pathway as it requires 2 mol of ATP and 2 mol of redox equivalents for each mole of acetyl-CoA produced. In contrast, the RuMP pathway is the most energy-efficient, because for each mole of acetyl-CoA produced, 2 mol of NADH plus one mole ATP are generated. Finally, the XuMP pathway generates 2 mol of NADH but consumes one mole ATP per mole of acetyl-CoA produced. These relationships are reflected by the specific maximum growth rate of bacteria expressing each of these three pathways: serine pathway (0.015–0.07 h^{-1}) < XuMP pathway (0.06–0.11 h^{-1}) < RuMP pathway (0.1–0.14 h^{-1}) (Cotton et al., 2020; Vartiainen et al., 2019).

Between all possible routes in the methanol oxidation and assimilation modules, four combinations are naturally occurring in methylotrophs. The first is the combination of NAD-dependent methanol dehydrogenase and ribulose monophosphate pathway (NMDH-RuMP pathway, NR) (Dijkhuizen et al., 1992), mainly found in thermophilic Gram-positive methylotrophic bacteria. The second is the oxygen-requiring alcohol oxidase followed by the xylulose monophosphate pathway (AOX-XuMP pathway, AX) (Schroer et al., 2010), which mainly exist in methylotrophic yeasts. The third is methanol dehydrogenase

combined with the serine pathway (NMDH-serine pathway, NS) (Vuilleumier et al., 2009), also mainly found in bacteria. The last is pyrroloquinoline quinone (PQQ) dependent MDH followed by the ribulose monophosphate (PQQ-RuMP pathway, PR) pathway, mainly found in Gram-negative methylotrophic bacteria (Keltjens et al., 2014). Among those four native methanol metabolic pathways, the NR pathway incurs the lowest energy-cost, generating 1 mol ATP and 5 mol NADH (which could be converted to energy via respiration) when 3 mol methanol is fixed. However, this pathway has a positive $\Delta_r G'$ (+34.2 kJ/mol). The PR route is the second most efficient in terms of energetics, producing 1 mol ATP and 2 mol NADH when 3 mol methanol is fixed; followed by the AX pathway, producing 2 mol NADH but consuming 1 mol ATP when 3 mol methanol fixed. The AX pathway exhibits the lowest $\Delta_r G'$ (−99.2 kJ/mol), making it the most thermodynamically feasible in synthetic methylotrophs. Finally, the pathway with the highest energy cost is the NS pathway, which consumes 6 mol ATP and 6 mol NADH to fix 3 mol methanol.

Because of these energy generation characteristics, currently NR is the most common pathway to be engineered in the construction of methylotrophic microorganisms, followed by the AX and NS pathways (Dai et al., 2017; Gonzalez et al., 2018; Meyer et al., 2018; Yu & Liao, 2018).

3. ENGINEERING THE METHANOL UTILIZATION PATHWAYS

Although C–C bond construction from C1 compounds via chemical synthesis has been proven possible, this process remains challenging due to the requirement of high temperature, pressure, and energy

(Yu & Liao, 2018). Bioconversion is a promising alternative, with a relatively low energy cost and being generally an environmentally friendly process. The current use of native methylotrophic microorganisms, however, suffers from certain drawbacks such as poor genetic availability and low metabolic yield. Therefore, engineering nonnative methylotrophic microbes is a promising route to harvest C1 sources and convert them into value-added products. Below we review recent progress in engineering nonnative methylotrophic microbes both in vitro and in vivo.

3.1 In vitro engineering of the methanol utilization pathways

Cell-free methanol utilization systems have been constructed which side-step many complex cellular processes such as gene transcription, protein folding and assembly, and competing pathways within the cellular metabolic network (Price et al., 2016). The NR pathway is preferred for in vitro constructions of methanol metabolic pathways, due to its more efficient energetics in terms of ATP and NADH production (Bogorad et al., 2014). However, a significant energy barrier of NMDH must be overcome, which requires HCHO and NADH to be maintained at low concentrations to ensure that the reaction favors methanol oxidation. To improve NMDH activity and increase its affinity to methanol, two strategies have been tested. The first is to express an NAD-dependent but ACT-independent Mdh2 enzyme. Both directed evolution and protein engineering have been used to construct such a synthetic enzyme. For example, directed molecular evolution has resulted in a mutated version of CT4-1 (A126V, A31V, A169V) which has a 6-fold higher K_{cat}/K_m for methanol ($9.3 \text{ M}^{-1} \text{ s}^{-1}$) and 10-fold lower K_{cat}/K_m for n-butanol ($48 \text{ M}^{-1} \text{ s}^{-1}$) in vitro (Wu et al., 2016). In an independent study, a phage-assisted noncontinuous evolution (PANCE) method was adapted for Mdh2 evolution, resulting in an Mdh2 variant (Q5L, A363L) with up to 3.5-fold higher V_{max} (Roth et al., 2019). The second strategy to improve NMDH activity is to compress the reaction space. For example, by using a scaffoldless self-assembly methodology to organize Mdh, Hps, and Phi into an engineered supramolecular enzyme complex via SH3-ligand interaction pairing, in vitro production of fructose-6-phosphate (F6P) from methanol improved 97-fold (Price et al., 2016). Expression of an "NADH Sink" using the *E. coli* lactate dehydrogenase as an NADH scavenger, further improved methanol consumption in vitro.

This concept described above has also been tested in vivo using *E. coli*, however the improvement in methanol consumption here was smaller compared with that achieved in vitro. Although in vitro cell-free systems have shown outstanding characteristics in biofuels production (Korman et al., 2017; Y.-H. P. Zhang, 2011), challenges remain such as stability of reaction system for multiple enzyme reactions (Bogorad et al., 2014) and toxic metabolites, for example, formaldehyde tolerance, to operate at large-scale.

Finally, to achieve the highest possible carbon conservation, a methanol condensation cycle (MCC) has been constructed by a

combination of non-oxidative glycolysis (NOG) and the RuMP pathway. This synthetic approach completely avoids the complexity of NMDH from methylotrophs. MCC was first proven to be functional in vitro by conversion of methanol to ethanol or n-butanol. Using alcohol dehydrogenase from *S. cerevisiae*, the MCC was demonstrated to produce 610 mg/L ethanol from 6200 mg/L methanol, or 170 mg/L n-butanol from 6200 mg/L methanol in vitro (Bogorad et al., 2014). However, the use of MCC in vivo has not been reported so far.

3.2 Engineering nonnative methylotrophic microbes

Although progress have been shown in vitro, it is more challenging to achieve methanol utilization in vivo due to the complexity of the cellular environment. Engineering methanol metabolic pathways in nonnative methylotrophs have been mainly carried out in *E. coli*, *S. cerevisiae* and *Corynebacterium glutamicum* (Bennett et al., 2018) to produce high-value commodities (Balk et al., 2003; Fei et al., 2014; Haynes & Gonzalez, 2014).

In *E. coli*, since the cofactor PQQ required in PR pathway cannot be synthesized (Anthony, 2004; Davidson, 2001; Matsushita et al., 1997), the NR pathway is the most frequent choice to engineer (J. E. Müller et al., 2015; Rohlhill et al., 2017; W. Brian Whitaker et al., 2017). A number of NMDH enzyme candidates from different host organisms have been evaluated in *E. coli* for methanol utilization. ^{13}C labeling experiments showed that up to 40% incorporation of carbon in glycolytic and pentose phosphate pathway (PPP) intermediates were derived from methanol, when cells were incubated in co-carbon medium (J. E. Müller et al., 2015; W. Brian Whitaker et al., 2017). By expressing an NMDH homolog from *B. stearotermophilus* (W. Brian Whitaker et al., 2017), and with supplementation of yeast extract in the growth media, cells were able to grow on methanol (from $\text{OD}_{600} = 0.04$ to 0.08) (Gonzalez et al., 2018), and convert methanol to naringenin (W. Brian Whitaker et al., 2017).

In addition to *E. coli*, the NR pathway has also been integrated into *Corynebacterium glutamicum*. ^{13}C -labeling results indicated that more than 25% methanol was incorporated into glycolytic and PPP intermediates (Tuyishime et al., 2018). Moreover, supplying methanol as a co-substrate together with glucose or ribose further improved methanol consumption (with a methanol consumption rate of 1.7 mM h^{-1}) (Leßmeier & Pfeifenschneider, Carnicer, et al., 2015; Leßmeier & Wendisch, 2015; Witthoff et al., 2015).

In addition to the NR pathway, the AX pathway has also been tested for methylotrophic microbe construction. Results indicated that the AX pathway was more efficient than the NR pathway in *S. cerevisiae* (Dai et al., 2017). Lastly, a non-naturally occurring "hybrid" pathway (NMDH-Xu5P), which was rationally designed based on the observed advantages and disadvantages of natural methanol metabolic pathways, has also been introduced into *S. cerevisiae*. In this synthetic yeast strain, ^{13}C -labeling experiments showed that more than 30% CO_2 production was derived from methanol (Espinosa et al., 2019).

4. CHALLENGES IN ENGINEERING SYNTHETIC METHYLOTROPHS

Despite significant efforts in developing synthetic methylotrophs in *E. coli*, *C. glutamicum* and *S. cerevisiae*, to date none of engineered strains can grow on methanol as the sole carbon source. Below we summarize some of the ground challenges that are limiting in methylotrophic microbe construction.

4.1 Tolerance of metabolic intermediates

While methanol is a promising carbon source, many metabolites of its catabolic pathways, including formaldehyde (Yurimoto et al., 2005), DHA (Molin & Blomberg, 2006), and H_2O_2 (Yurimoto, 2009), are toxic for cells (Table 2). Among these metabolic intermediates, formaldehyde is the most toxic to many microorganisms. Studies have shown that the growth defect of methylotrophic yeast (*Pichia methanolica*) on high methanol media is not caused directly by methanol toxicity, but rather by formaldehyde toxicity (Wakayama et al., 2016). To counter this toxicity, the formaldehyde dissimilation pathway can be activated to detoxify formaldehyde (Pfeifenschneider et al., 2017). These results suggest that the formaldehyde assimilation module may be the limiting step in methanol utilization (Papoutsakis et al., 1978). Therefore, the heterologous hosts must be engineered to either be tolerant to formaldehyde, or have efficient pathways to quickly detoxify it. Two approaches can be taken to reduce formaldehyde toxicity in methylotrophic strains. The first is to strengthen the efficiency of the formaldehyde dissimilation pathway. This strategy not only accelerates formaldehyde degradation, but also provides NADH which can be used for ATP generation. However, it leads to carbon loss in the form of CO_2 . The second approach is to convert formaldehyde into a less toxic compound. For example, in the AX pathway the affinity of dihydroxyacetone synthase (DAS, Figure 2) towards formaldehyde is much higher ($K_m = 0.43$) (Kato et al., 1982) than that of AOX for methanol ($K_m = 3$) (Nichols & Cromartie, 1980), which indicates that DAS may be well-suited for quickly relieving formaldehyde toxicity in synthetic microbes utilizing AX pathway for methanol oxidation.

4.2 Cofactor balance

To overcome the thermodynamic constraint on methanol oxidation catalyzed by NMDH ($\Delta_r G' = 34.2 \pm 6.5 \text{ kJ mol}^{-1}$), an alternative method to promote this reaction is to accelerate NAD^+ regeneration. One way to achieve this is to weaken other NAD^+ utilization pathways. For example, Meyer et al. has constructed an *E. coli* strain wherein methanol is essential for growth, by knocking out NAD^+ -dependent malate dehydrogenase and reducing the function of the NAD repressor nicotinamide mononucleotide adenylyltransferase NadR (Meyer et al., 2018). These modifications led to a rebalance toward a high $NAD^+/NADH$ ratio, which thermodynamically favored

the oxidation of methanol. Another promising method is to build an NADH-ATP generation cycle. Since NADH can be transported into the mitochondria to generate ATP, enhancing the NADH shuttle system (i.e., malate-aspartate shuttle) (Eto et al., 1999) may be a good solution that is capable of not only promoting NAD^+ regeneration, but also providing additional ATP which can be used to further increase methanol utilization.

4.3 Energy supply

Based on the reactions shown in Table 1, it is clear that even though the NR pathway is energetically self-sufficient, excessive NADH will inhibit NMDH activity. This may be the reason why an NADH sink was necessary in a previous report of microbes expressing the NR pathway to metabolize methanol (Price et al., 2016). In contrast to the RuMP cycle, all other methanol-assimilating pathways have an energy demand. So far, two methods have been used to meet this energy demand. One is to strengthen the formaldehyde dissimilation pathway, which produces ATP from the generated NADH, although it can lead to > 40% carbon loss (Jorda et al., 2014). The formaldehyde dissimilation pathway generates 2 mol NADH in two enzymatic steps, which requires a very small proteomic investment and thus more proteomically efficient than glycolysis and the TCA cycle in generating protons for energy production. Moreover, enhanced formaldehyde dissimilation would relieve formaldehyde toxicity, which is likely why the TCA cycle was inhibited while the formaldehyde dissimilation pathway was up regulated in *P. pastoris* (Rußmayer et al., 2015).

The second method to meet the energy demand of methanol utilization is by adding other carbohydrates to the culture medium (Gonzalez et al., 2018; Meyer et al., 2018; Woolston, Roth, et al., 2018). For example, when the serine cycle was introduced into *E. coli*, the engineered strains were able to co-assimilate methanol together with a pyruvate source, i.e., xylose, thereby improving the production of acetyl-CoA derived C2 compounds (Yu & Liao, 2018). Dai et al. showed that when the AX pathway was introduced into *S. cerevisiae*, the recombinant strain consumed methanol up to 2.35 g/L, leading to 11.70% increase in cell growth (OD_{600}), when supplied with 1 g/L yeast extract (Dai et al., 2017). In *C. glutamicum*, biomass was improved by up to 30% in minimal medium containing both methanol and glucose (Witthoff et al., 2015), and up to 15.7% ^{13}C -labeled methanol was used for cadaverine production in minimal medium containing both methanol and ribose (Leßmeier et al., 2015). As additional carbohydrates are required to supply the energy necessary for methanol degradation, growth with NR independent pathway on methanol as the sole carbon source would be difficult.

4.4 Intermediate recycling

The last but not least challenge is the regeneration of intermediates such as ribulose-5-phosphate (Ru5P) or xylulose-5-phosphate

TABLE 2 Tolerance for different intermediates

Compounds	Organism	Max tolerance concentration	KM (mM) ^a	Characteristics
Formaldehyde (HCHO)	<i>Yarrowia lipolytica</i>	1 mM (Vartiainen et al., 2019)	Dihydroxyacetone synthase (DAS): 0.43 (Kato et al., 1982) -1.86 (Ro et al., 1997), hexulose phosphate synthase (HPS): 0.98 (Rozova et al., 2017)-1.4 (R. Müller & Babel, 1980)	Damage proteins, nucleic acid and lipids and inhibit cell grow (Nian et al., 2013; Pilat & Prokop, 1975; Songur et al., 2010)
	<i>S. cerevisiae</i>	IC ₂₀ = 0.6 mM (North et al., 2016)		
	<i>Pichia methanolica</i>	0.3 mM (Wakayama et al., 2016)		
	<i>Candida boidini</i> 11 Bh	3.6-4.0 mM (Pilat & Prokop, 1976)		
	<i>Pichia pastoris</i>	5 mM (Sunga & Cregg, 2004)		
Dihydroxyacetone (DHA)	<i>S. cerevisiae</i>	200 mM (Molin & Blomberg, 2006; Molin et al., 2003)	Dihydroxyacetone kinase (DAK): 0.005 (Molin et al., 2003) -0.008 (Gutknecht et al., 2001)	Carbon-based toxicity (Molin et al., 2003) and the toxicity of DHA appears to be due to its intracellular conversion to an aldehyde compound, presumably methylglyoxal (Subedi et al., 2008)
Hydrogen peroxide (H ₂ O ₂)	<i>S. cerevisiae</i>	2-3 mM (Hasan et al., 2002; Poljak et al., 2003)	Catalase (CAT1): 0.65 (Hochman & Goldberg, 1991) - 15 (Santoni et al., 2004)	Result in DNA and membrane damage that cause mutagenesis and kill cells (Brunk et al., 1995; Imlay et al., 1988)
	<i>Candida albicans</i>	50 mM (González-Párraga et al., 2003)		
	<i>Yarrowia lipolytica</i>	120 mM (Biryukova et al., 2006)		

^aThe KM values are obtained from BRENDA (<https://www.brenda-enzymes.org/>).

(Xu5P). The most common solution to this is to use another carbohydrate as a co-substrate to sustain a pool of these intermediates. Several feedstocks such as gluconate and xylose have been used as precursors for these intermediates to support methanol utilization. Vorholt et al. showed that gluconate or glucose could provide Ru5P through the PPP, which significantly accelerates HCHO fixation (Meyer et al., 2018). Woolston, Roth, et al. (2018) showed that xylose is a promising co-carbon source for HCHO fixation owing to its capability of forming Xu5P or Ru5P directly. Although such co-carbon strategies could promote methanol utilization, biomass formation is not mainly supported by methanol. Engineering of alternative pathway for regeneration of C5 sugar phosphates in bacteria, as demonstrated by activating sedoheptulose biphosphatase variants, may provide some hints for the eventual resolution of this problem (Woolston, Roth, et al., 2018). Hannes et al. have shown that the Xu5P regeneration pathway has been duplicated in the evolution of *P. pastoris*, which has been compartmentalized into the peroxisome (Rußmayer et al., 2015). This indicates that engineering of enzyme compartmentalization may be an alternative strategy to improve the recycling of metabolic intermediates in methanol utilization pathways.

5. CONCLUSION REMARKS

In recent years, much attention has been focused on the development of methylotrophic organisms due to the significant advantages of methanol fermentation and its potential application in conversion of methane. However, it is clear that most of the engineered methylotrophic strains showed poor growth capability in methanol medium (Cantera et al., 2018; Duan et al., 2015; Schrader et al., 2009), which limits the engineering of methanol metabolic pathways to produce high-value products in both native and nonnative methylotrophic microorganisms. To promote the construction of methylotrophic microbial cell factories, the reiterative design-build-test-learn (DBTL) cycle of metabolic engineering (Chen et al., 2017; Nielsen & Keasling, 2016) may be employed as described below.

5.1. Design

In the process of engineering native and nonnative methylotrophy, most efforts have been focused on introducing natural pathways into model microbes. However, each natural pathway presents a different set of drawbacks, therefore different synthetic novel pathways may be better suitable for different applications. For example, computational analysis of thousands of metabolic reactions to address energy consumption and carbon loss, have led to several nonnatural formaldehyde assimilation pathways (Yang et al., 2019). Also, some synthetic or 'hybrid' pathways like variants of reductive glycine pathway (Cotton et al., 2020) have been found to outperform the native pathway. For each application, to achieve

the optimal design of a methanol-utilizing microbial cell factory, three modules should be considered: module I (methanol-degradation module)—methanol adsorption and fixation; module II (energy-producing module)—energy generation to support methanol degradation; and module III (Intermediate recycle module)—supplying sufficient intermediates for HCHO assimilation and promoting intermediate recycle (Figure 3). In addition, a balance between the characteristics of the final products, the efficiency of the designed pathway(s), and the biomass yield should be carefully considered during the design phase.

5.2. Build

Novel synthetic biology modules and tools should be employed for the construction and optimization of methanol-utilizing microbial cell factories. For instance, one key tool is for heterologous gene expression, whereby methanol utilization genes are identified and introduced into host species, for example using CRISPR/Cas9-based methodologies (Caspeta et al., 2014). To finetune the expression of these heterologous genes, regulatory elements such as gene promoters may be targeted for engineering, as many studies have shown that most genes involved in methanol utilization pathways are tightly regulated by methanol or its metabolites (Rußmayer et al., 2015; Yurimoto et al., 2000). Other enzyme characteristics can also be optimized, for example by improving substrate affinity and reducing substrate/metabolites inhibition, via directed evolution, rational protein engineering or a combination of enzyme engineering methodologies (Eriksen et al., 2014). The use of advanced biosensors, that is, the recently developed formaldehyde biosensor (Woolston, Roth, et al., 2018), should also be incorporated in the process of building methanol-utilizing microbial cell factories, to dynamically control the methanol dissimilation pathway, and thus supporting energy generation and reducing carbon loss via this route.

5.3. Test

The purpose of this step of the DBTL cycle is to characterize the constructed cell factories, which provides crucial information for the learning phase and to guide the next iterative DBTL cycle. A number of evaluation criteria can be applicable as an evaluation index, of which the cell growth rate on methanol is the most intuitive and most commonly used. In addition to the cell growth rate, adaptability to methanol can be evaluated via verification of the utilization efficiency of methanol as a carbon source, and tolerance to methanol and its downstream metabolites. Pathway efficiency and omics-based analyses can also be used in the testing phase to evaluate the microbial cell factories, for example, through transcription network analysis and metabolic flux analysis (Figure 3). As these evaluations are often costly and time-consuming, strategies to shorten the testing phase and improve evaluation efficiency, such as mechanistic/robotic high-throughput




FIGURE 3 The use of design–build–test–learn cycle for methylotrophic microorganism construction. Design: Based on available knowledge and computational-aided modeling to guide the design of efficient metabolic pathways for methanol carbon utilization. Build: using novel synthetic biology tools and modules to construct metabolic pathways in different microbial hosts for methanol utilization. Test: detailed characterization of constructed strains in terms of cell growth, product formation, and even omics-based phenotyping of strains. Learn: the generated data will be combined with computational based integrative analysis to find further engineering targets for the next cycle, to reach finally the desired methylotrophic microbial cell factory [Color figure can be viewed at wileyonlinelibrary.com]

approaches, careful selection of specific test characteristics, and/or rapid omics-based analyses would lead to significant improvements in the DBTL cycle to construct methanol-utilizing cell factories (Srivatsan et al., 2020).

5.4. Learn

A better understanding of the metabolic pathways in native methylotrophs is an important step towards constructing more efficient synthetic methylotrophic organisms. For example, based on a deep analysis of the AX pathway in *P. pastoris*, combined with lessons learned while constructing nonnative methylotrophic microbes, it is now understood that a sufficient supply of metabolic intermediates, high-efficiency energy transformation from NADH, as well as the ability to dissimilate formaldehyde, are three of the most important characteristics for efficient AX pathway construction. In addition, more and more synthetic methylotrophic constructions and characterizations through the iterative design–build–test–learn (DBTL) cycle of metabolic engineering will generate more useful data and provide new knowledge for future methylotrophic microorganism construction.

ACKNOWLEDGMENTS

We thank Rosemary Yu for help with final polishing of the manuscript. This study is financially supported by the National Natural Science Foundation of China (21908077, 31570034) and the Novo Nordisk Foundation (grant no. NNF10CC1016517).

AUTHOR CONTRIBUTIONS

Chunjun Zhan wrote the manuscript. Xiaowei Li provided related papers. Yankun Yang gave some useful suggestions. Jens Nielsen, Zhonghu Bai, and Yun Chen revised this paper and gave lots of great suggestions. All authors read and approved the final manuscript.

ORCID

Chunjun Zhan  <http://orcid.org/0000-0001-9504-6439>

Yankun Yang  <http://orcid.org/0000-0003-4815-6832>

Yun Chen  <https://orcid.org/0000-0002-2146-6008>

REFERENCES

- Anthony, C. (2004). The quinoprotein dehydrogenases for methanol and glucose. *Archives of Biochemistry and Biophysics*, 428(1), 2–9. Retrieved from <https://www.sciencedirect.com/science/article/pii/S0003986104001808?via%3Dihub>

- Antonovsky, N., Gleizer, S., Noor, E., Zohar, Y., Herz, E., Barenholz, U., Zelcbuch, L., Amram, S., Wides, A., Tepper, N., Davidi, D., Bar-On, Y., Bareia, T., Wernick, D. G., Shani, I., Malitsky, S., Jona, G., Bar-Even, A., & Milo, R. (2016). Sugar synthesis from CO₂ in *Escherichia coli*. *Cell*, 166(1), 115-125.
- Arfman, N., Van Beeumen, J., De Vries, G. E., Harder, W., & Dijkhuizen, L. (1991). Purification and characterization of an activator protein for methanol dehydrogenase from thermotolerant *Bacillus* spp. *Journal of Biological Chemistry*, 266(6), 3955-3960.
- Arfman, N., Dijkhuizen, L., Kirchof, G., Ludwig, W., Schleifer, K.-H., Bulygina, E. S., Chumakov, K. M., Govorukhina, N. I., Trotsenko, Y. A., & White, D. (1992). *Bacillus methanolicus* sp. nov., a new species of thermotolerant, methanol-utilizing, endospore-forming bacteria. *International Journal of Systematic and Evolutionary Microbiology*, 42(3), 439-445.
- Arfman, N., Hektor, H. J., Bystrykh, L. V., Govorukhina, N. I., Dijkhuizen, L., & Frank, J. (1997). Properties of an NAD (H)-containing methanol dehydrogenase and its activator protein from *Bacillus methanolicus*. *European Journal of Biochemistry*, 244(2), 426-433.
- Bahrami, A., Shojaosadati, S., Khalilzadeh, R., & Farahani, E. V. (2008). Two-stage glycerol feeding for enhancement of recombinant hG-CSF production in a fed-batch culture of *Pichia pastoris*. *Biotechnology Letters*, 30(6), 1081-1085.
- Balk, M., Weijma, J., Friedrich, M. W., & Stams, A. J. (2003). Methanol utilization by a novel thermophilic homoacetogenic bacterium, *Moorella mulderi* sp. nov., isolated from a bioreactor. *Archives of Microbiology*, 179(5), 315-320.
- Bennett, R. K., Steinberg, L. M., Chen, W., & Papoutsakis, E. T. (2018). Engineering the bioconversion of methane and methanol to fuels and chemicals in native and synthetic methylophs. *Current Opinion in Biotechnology*, 50, 81-93. <https://doi.org/10.1016/j.copbio.2017.11.010>
- Bertau, M., Offermanns, H., Plass, L., Schmidt, F., & Wernicke, H.-J. (2014). *Methanol: The basic chemical and energy feedstock of the future*. Springer.
- Birdja, Y. Y., Pérez-Gallent, E., Figueiredo, M. C., Göttele, A. J., Calle-Vallejo, F., & Koper, M. T. (2019). Advances and challenges in understanding the electrocatalytic conversion of carbon dioxide to fuels. *Nature Energy*, 4(9), 732-745.
- Biryukova, E., Medentsev, A., Arinbasarova, A. Y., & Akimenko, V. (2006). Tolerance of the yeast *Yarrowia lipolytica* to oxidative stress. *Microbiology*, 75(3), 243-247.
- Bogorad, I. W., Chen C.-T., Theisen, M. K., Wu, T.-Y., Schlenz, A. R., Lam, A. T., & Liao, J. C. (2014). Building carbon-carbon bonds using a biocatalytic methanol condensation cycle. *Proceedings of the National Academy of Sciences*, 111(45), 15928-15933.
- Bozdag, A. (2013). *Investigation of methanol and formaldehyde metabolism in Bacillus methanolicus* (Doctoral dissertation). North Carolina State University, Raleigh, NC.
- Brunk, U., Zhang, H., Roberg, K., & Öllinger, K. (1995). Lethal hydrogen peroxide toxicity involves lysosomal iron-catalyzed reactions with membrane damage. *Redox Report*, 1(4), 267-277.
- Budzianowski, W. M. (2012). Negative carbon intensity of renewable energy technologies involving biomass or carbon dioxide as inputs. *Renewable and Sustainable Energy Reviews*, 16(9), 6507-6521.
- Cantera, S., Muñoz, R., Lebrero, R., López, J. C., Rodríguez, Y., & García-Encina, P. A. (2018). Technologies for the bioconversion of methane into more valuable products. *Current Opinion in Biotechnology*, 50, 128-135.
- Caspeta, L., Chen, Y., Ghiaci, P., Feizi, A., Buskov, S., Hallström, B. M., Petranovic, D., & Nielsen, J. (2014). Altered sterol composition renders yeast thermotolerant. *Science*, 346(6205), 75-78.
- Cereghino, J. L., & Cregg, J. M. (2000). Heterologous protein expression in the methylophic yeast *Pichia pastoris*. *FEMS Microbiology Reviews*, 24(1), 45-66.
- Chen, X., Gao, C., Guo, L., Hu, G., Luo, Q., Liu, J., Nielsen, J., Chen, J., & Liu, L. (2017). DCEO biotechnology: Tools to design, construct, evaluate, and optimize the metabolic pathway for biosynthesis of chemicals. *Chemical Reviews*, 118(1), 4-72.
- Cotton, C. A., Claassens, N. J., Benito-Vaquerizo, S., & Bar-Even, A. (2020). Renewable methanol and formate as microbial feedstocks. *Current Opinion in Biotechnology*, 62, 168-180.
- Couderc, R., & Baratti, J. (1980). Oxidation of methanol by the yeast, *Pichia pastoris*. Purification and properties of the alcohol oxidase. *Agricultural and Biological Chemistry*, 44(10), 2279-2289.
- Cox, R. B., & Quayle, J. R. (1975). The autotrophic growth of *Micrococcus denitrificans* on methanol. *Biochemical Journal*, 150(3), 569-571.
- Dai, Z., Gu, H., Zhang, S., Xin, F., Zhang, W., Dong, W., Ma, J., Jia, H., & Jiang, M. (2017). Metabolic construction strategies for direct methanol utilization in *Saccharomyces cerevisiae*. *Bioresource Technology*, 245, 1407-1412. <https://doi.org/10.1016/j.biortech.2017.05.100>
- Davidson, V. L. (2001). Pyrroloquinoline quinone (PQQ) from methanol dehydrogenase and tryptophan tryptophylquinone (TTQ) from methylamine dehydrogenase. *Advances in Protein Chemistry*, 58, 95-140.
- Dijkhuizen, L., Levering, P., & De Vries, G. (1992). Methane and methanol utilizers, *The physiology and biochemistry of aerobic methanol-utilizing Gram-negative and Gram-positive bacteria* (Vol. 56, pp. 149-181). Springer.
- Duan, H., Wang, H., Ma, B., Jiang, P., Tu, P., Ni, Z., Li, X., Li, M., Ma, X., Wang, B., Wu, R., & Li, M. (2015). Codon optimization and expression of irisin in *Pichia pastoris* GS115. *International Journal of Biological Macromolecules*, 79, 21-26.
- Ellis, S. B., Brust, P. F., Koutz, P. J., Waters, A., Harpold, M. M., & Gingeras, T. R. (1985). Isolation of alcohol oxidase and two other methanol regulatable genes from the yeast *Pichia pastoris*. *Molecular and Cellular Biology*, 5(5), 1111-1121.
- Erb, T. J., & Zarzycki, J. (2016). Biochemical and synthetic biology approaches to improve photosynthetic CO₂-fixation. *Current Opinion in Chemical Biology*, 34, 72-79.
- Erb, T. J., & Zarzycki, J. (2018). A short history of rubis CO: The rise and fall (?) of Nature's predominant CO₂ fixing enzyme. *Current Opinion in Biotechnology*, 49, 100-107. <https://doi.org/10.1016/j.copbio.2017.07.017>
- Eriksen, D. T., Lian, J., & Zhao, H. (2014). Protein design for pathway engineering. *Journal of Structural Biology*, 185(2), 234-242.
- Espinosa, M. I., Valgepea, K., Gonzalez-Garcia, R. A., Scott, C., Pretorius, I. S., Marcellin, E., & Williams, T. (2019). Native and synthetic methanol assimilation in *Saccharomyces cerevisiae*. *bioRxiv*, 717942.
- Eto, K., Suga, S., Wakui, M., Tsubamoto, Y., Terauchi, Y., Taka, J., Aizawa, S., Noda, M., Kimura, S., Kasai, H., & Kadowaki, T. (1999). NADH shuttle system regulates KATP channel-dependent pathway and steps distal to cytosolic Ca²⁺ concentration elevation in glucose-induced insulin secretion. *Journal of Biological Chemistry*, 274(36), 25386-25392.
- Fei, Q., Guarnieri, M. T., Tao, L., Laurens, L. M., Dowe, N., & Pienkos, P. T. (2014). Bioconversion of natural gas to liquid fuel: Opportunities and challenges. *Biotechnology Advances*, 32(3), 596-614.
- Fujiwara, K., Tada, S., Honma, T., Sasaki, H., Nishijima, M., & Kikuchi, R. (2019). Influences of particle size and crystallinity of highly loaded CuO/ZrO₂ on CO₂ hydrogenation to methanol. *AIChE Journal*, 65(12), e16717.
- Gonzalez, J. E., Bennett, R. K., Papoutsakis, E. T., & Antoniewicz, M. R. (2018). Methanol assimilation in *Escherichia coli* is improved by co-utilization of threonine and deletion of leucine-responsive regulatory protein. *Metabolic Engineering*, 45, 67-74.
- González-Párraga, P., Hernández, J. A., & Argüelles, J. C. (2003). Role of antioxidant enzymatic defences against oxidative stress (H₂O₂) and

- the acquisition of oxidative tolerance in *Candida albicans*. *Yeast*, 20(14), 1161-1169.
- Gutknecht, R., Beutler, R., Garcia-Alles, L. F., Baumann, U., & Erni, B. (2001). The dihydroxyacetone kinase of *Escherichia coli* utilizes a phosphoprotein instead of ATP as phosphoryl donor. *The EMBO Journal*, 20(10), 2480-2486.
- Hasan, R., Leroy, C., Isnard, A. D., Labarre, J., Boy-Marcotte, E., & Toledano, M. B. (2002). The control of the yeast H₂O₂ response by the Msn2/4 transcription factors. *Molecular Microbiology*, 45(1), 233-241.
- Haynes, C. A., & Gonzalez, R. (2014). Rethinking biological activation of methane and conversion to liquid fuels. *Nature Chemical Biology*, 10(5), 331-339.
- Hochman, A., & Goldberg, I. (1991). Purification and characterization of a catalase-peroxidase and a typical catalase from the bacterium *Klebsiella pneumoniae*. *Biochimica et Biophysica Acta—Protein Structure and Molecular Enzymology*, 1077(3), 299-307.
- Hu, B., Guild, C., Suib, S., Util, J., Aresta, M., Dibenedetto, A., & Angelini, A. (2013). *Journal of CO₂ Utilization*, 1, 18-27.
- Hu, G., Li, Y., Ye, C., Liu, L., & Chen, X. (2018). Engineering microorganisms for enhanced CO₂ sequestration. *Trends in Biotechnology*, 37, 532-547.
- Imlay, J. A., Chin, S. M., & Linn, S. (1988). Toxic DNA damage by hydrogen peroxide through the Fenton reaction in vivo and in vitro. *Science*, 240(4852), 640-642.
- Inan, M., & Meagher, M. M. (2001). Non-repressing carbon sources for alcohol oxidase (AOX1) promoter of *Pichia pastoris*. *Journal of Bioscience and Bioengineering*, 92(6), 585-589.
- Jakobsen, Ø. M., Benichou, A., Flickinger, M. C., Valla, S., Ellingsen, T. E., & Brautaset, T. (2006). Upregulated transcription of plasmid and chromosomal ribulose monophosphate pathway genes is critical for methanol assimilation rate and methanol tolerance in the methylotrophic bacterium *Bacillus methanolicus*. *Journal of Bacteriology*, 188(8), 3063-3072.
- Jorda, J., de Jesus, S. S., Peltier, S., Ferrer, P., & Albiol, J. (2014). Metabolic flux analysis of recombinant *Pichia pastoris* growing on different glycerol/methanol mixtures by iterative fitting of NMR-derived ¹³C-labelling data from proteinogenic amino acids. *New Biotechnology*, 31(1), 120-132.
- Jordà, J., Jouhten, P., Cámara, E., Maaheimo, H., Albiol, J., & Ferrer, P. (2012). Metabolic flux profiling of recombinant protein secreting *Pichia pastoris* growing on glucose:methanol mixtures. *Microbial Cell Factories*, 11(1), 57.
- Jungo, C., Schenk, J., Pasquier, M., Marison, I. W., & von Stockar, U. (2007). A quantitative analysis of the benefits of mixed feeds of sorbitol and methanol for the production of recombinant avidin with *Pichia pastoris*. *Journal of Biotechnology*, 131(1), 57-66.
- Kalyuzhnaya, M. G., Hristova, K. R., Lidstrom, M. E., & Chistoserdova, L. (2008). Characterization of a novel methanol dehydrogenase in representatives of Burkholderiales: Implications for environmental detection of methylotrophy and evidence for convergent evolution. *Journal of Bacteriology*, 190(11), 3817-3823.
- Kato, N., Higuchi, T., Sarazawa, C., Nishizawa, T., Tani, Y., & Yamada, H. (1982). Purification and properties of a transketolase responsible for formaldehyde fixation in a methanol-utilizing yeast, *Candida boidinii* (Kloekera sp.) No. 2201. *Biochimica et Biophysica Acta—General Subjects*, 715(2), 143-150.
- Keltjens, J. T., Pol, A., Reimann, J., & den Camp, H. J. O. (2014). PQQ-dependent methanol dehydrogenases: Rare-earth elements make a difference. *Applied Microbiology and Biotechnology*, 98(14), 6163-6183.
- Korman, T. P., Opgenorth, P. H., & Bowie, J. U. (2017). A synthetic biochemistry platform for cell free production of monoterpenes from glucose. *Nature Communications*, 8(1), 1-8.
- Krainer, F. W., Dietzsch, C., Hajek, T., Herwig, C., Spadiut, O., & Glieder, A. (2012). Recombinant protein expression in *Pichia pastoris* strains with an engineered methanol utilization pathway. *Microbial Cell Factories*, 11(1), 22.
- Latimer, A. A., Kakekhani, A., Kulkarni, A. R., & Nørskov, J. K. (2018). Direct methane to methanol: The selectivity-conversion limit and design strategies. *ACS Catalysis*, 8(8), 6894-6907.
- Leßmeier, L., Pfeifenschneider, J., Carnicer, M., Heux, S., Portais, J.-C., & Wendisch, V. F. (2015). Production of carbon-13-labeled cadaverine by engineered *Corynebacterium glutamicum* using carbon-13-labeled methanol as co-substrate. *Applied Microbiology and Biotechnology*, 99(23), 10163-10176.
- Leßmeier, L., & Wendisch, V. F. (2015). Identification of two mutations increasing the methanol tolerance of *Corynebacterium glutamicum*. *BMC Microbiology*, 15(1), 216.
- Linton, J. D., & Niekus, H. G. D. (1987). The potential of one-carbon compounds as fermentation feedstocks. *Antonie Van Leeuwenhoek*, 53(1), 55-63. <https://doi.org/10.1007/bf00422635>
- Liu, Z., Wang, K., Chen, Y., Tan, T., & Nielsen, J. (2020). Third-generation biorefineries as the means to produce fuels and chemicals from CO₂. *Nature Catalysis*, 3(3), 274-288.
- Long, B. M., Hee, W. Y., Sharwood, R. E., Rae, B. D., Kaines, S., Lim, Y.-L., Nguyen, N. D., Massey, B., Bala, S., von Caemmerer, S., Badger, M. R., & Price, G. D. (2018). Carboxysome encapsulation of the CO₂-fixing enzyme rubisco in tobacco chloroplasts. *Nature Communications*, 9(1), 1-14.
- Luo, X., Reiter, M. A., d'Espaux, L., Wong, J., Denby, C. M., Lechner, A., Zhang, Y., Grzybowski A. T., Harth S., Lin W., Lee H., Yu C., Shin J., Deng K., Benites V. T., Wang G., Baidoo E., Chen Y., Dev I., ... Keasling J. D. (2019). Complete biosynthesis of cannabinoids and their unnatural analogues in yeast. *Nature*, 567(7746), 123-126. <https://doi.org/10.1038/s41586-019-0978-9>
- Magdesieva, T., Zhukov, I., Kravchuk, D., Semenikhin, O., Tomilova, L., & Butin, K. (2002). Electrocatalytic CO₂ reduction in methanol catalyzed by mono-, di-, and electropolymerized phthalocyanine complexes. *Russian Chemical Bulletin*, 51(5), 805-812.
- Matsushita, K., Arents, J., Bader, R., Yamada, M., Adachi, O., & Postma, P. (1997). *Escherichia coli* is unable to produce pyrroloquinoline quinone (PQQ). *Microbiology*, 143(10), 3149-3156.
- Meyer, F., Keller, P., Hartl, J., Gröninger, O. G., Kiefer, P., & Vorholt, J. A. (2018). Methanol-essential growth of *Escherichia coli*. *Nature Communications*, 9(1), 1508.
- Molin, M., & Blomberg, A. (2006). Dihydroxyacetone detoxification in *Saccharomyces cerevisiae* involves formaldehyde dissimilation. *Molecular Microbiology*, 60(4), 925-938.
- Molin, M., Norbeck, J., & Blomberg, A. (2003). Dihydroxyacetone kinases in *Saccharomyces cerevisiae* are involved in detoxification of dihydroxyacetone. *Journal of Biological Chemistry*, 278(3), 1415-1423.
- Müller, J., Meyer, F., Litsanov, B., Kiefer, P., Potthoff, E., Heux, S., Quax, W. J., Wendisch, V. F., Brautaset, T., Portais, J. C., & Vorholt, J. A. (2015). Engineering *Escherichia coli* for methanol conversion. *Metabolic Engineering*, 28, 190-201.
- Müller, R., & Babel, W. (1980). A critical analysis of kinetic data of 3-hexulosephosphate synthases Michaelis-Menten or complex characteristics. *Zeitschrift für allgemeine Mikrobiologie*, 20(5), 325-333.
- Naik, S. N., Goud, V. V., Rout, P. K., & Dalai, A. K. (2010). Production of first and second generation biofuels: A comprehensive review. *Renewable and Sustainable Energy Reviews*, 14(2), 578-597.
- Nian, H., Meng, Q., Zhang, W., & Chen, L. (2013). Overexpression of the formaldehyde dehydrogenase gene from *Brevibacillus brevis* to enhance formaldehyde tolerance and detoxification of tobacco. *Applied Biochemistry and Biotechnology*, 169(1), 170-180.

- Nichols, C. S., & Cromartie, T. H. (1980). Irreversible inactivation of the flavoenzyme alcohol oxidase with acetylenic alcohols. *Biochemical and Biophysical Research Communications*, 97(1), 216-221.
- Nielsen, J., Fussenegger, M., Keasling, J., Lee, S. Y., Liao, J. C., Prather, K., & Palsson, B. (2014). Engineering synergy in biotechnology. *Nature Chemical Biology*, 10, 319-322. <https://doi.org/10.1038/nchembio.1519>
- Nielsen, J., & Keasling, J. D. (2016). Engineering cellular metabolism. *Cell*, 164(6), 1185-1197. <https://doi.org/10.1016/j.cell.2016.02.004>
- North, M., Gaytán, B. D., Romero, C., Jr., De La Rosa, V. Y., Loguinov, A., Smith, M. T., Zhang, L., & Vulpe, C. D. (2016). Functional toxicogenomic profiling expands insight into modulators of formaldehyde toxicity in yeast. *Frontiers in Genetics*, 7, 200.
- Ochsner, A. M., Müller, J. E., Mora, C. A., & Vorholt, J. A. (2014). In vitro activation of NAD-dependent alcohol dehydrogenases by *Nudix hydrolases* is more widespread than assumed. *FEBS Letters*, 588(17), 2993-2999.
- Papoutsakis, E., Lim, H. C., & Tsao, G. T. (1978). Role of formaldehyde in the utilization of C1 compounds via the ribulose monophosphate cycle. *Biotechnology and Bioengineering*, 20(3), 421-442.
- Paulo, P. L., Villa, G., van Lier, J. B., & Lettinga, G. (2003). The anaerobic conversion of methanol under thermophilic conditions: pH and bicarbonate dependence. *Journal of Bioscience and Bioengineering*, 96(3), 213-218.
- Pfeifenschneider, J., Brautaset, T., & Wendisch, V. F. (2017). Methanol as carbon substrate in the bio-economy: Metabolic engineering of aerobic methylotrophic bacteria for production of value-added chemicals. *Biofuels, Bioproducts and Biorefining*, 11(4), 719-731.
- Pilat, P., & Prokop, A. (1975). The effect of methanol, formaldehyde, and formic acid on growth of *Candida boidinii* 11 Bh. *Biotechnology and Bioengineering*, 17(12), 1717-1728.
- Pilat, P., & Prokop, A. (1976). Oxidation of methanol, formaldehyde and formic acid by methanol-utilizing yeast. *Folia Microbiologica*, 21(4), 306-314.
- Poljak, A., Dawes, I. W., Ingelse, B. A., Duncan, M. W., Smythe, G. A., & Grant, C. M. (2003). Oxidative damage to proteins in yeast cells exposed to adaptive levels of H₂O₂. *Redox Report*, 8(6), 371-377.
- Price, J. V., Chen, L., Whitaker, W. B., Papoutsakis, E., & Chen, W. (2016). Scaffoldless engineered enzyme assembly for enhanced methanol utilization. *Proceedings of the National Academy of Sciences of the United States of America*, 113(45), 12691-12696. <https://doi.org/10.1073/pnas.1601797113>
- Ro, Y. T., Eom, C. Y., Song, T., Cho, J. W., & Kim, Y. M. (1997). Dihydroxyacetone synthase from a methanol-utilizing carboxydobacterium, *Acinetobacter* sp. strain JC1 DSM 3803. *Journal of Bacteriology*, 179(19), 6041-6047.
- Rohllill, J., Sandoval, N. R., & Papoutsakis, E. T. (2017). Sort-seq approach to engineering a formaldehyde-inducible promoter for dynamically regulated *Escherichia coli* growth on methanol. *ACS Synthetic Biology*, 6(8), 1584-1595.
- Roth, T. B., Woolston, B. M., Stephanopoulos, G., & Liu, D. R. (2019). Phage-assisted evolution of *Bacillus methanolicus* methanol dehydrogenase 2. *ACS Synthetic Biology*, 8(4), 796-806.
- Rozova, O., But, S., Khmelenina, V., Reshetnikov, A., Mustakhimov, I., & Trotsenko, Y. (2017). Characterization of two recombinant 3-hexulose-6-phosphate synthases from the halotolerant obligate methanotroph *Methylomicrobium alcaliphilum* 20Z. *Biochemistry*, 82(2), 176-185.
- Rußmayer, H., Buchetics, M., Gruber, C., Valli, M., Grillitsch, K., Modarres, G., Guerrasio, R., Klavins K., Neubauer S., Drexler H., Steiger M., Troyer C., Al Chalabi A., Krebiehl G., Sonntag D., Zellnig G., Daum G., Graf A. B., Altmann F., ... Gasser B. (2015). Systems-level organization of yeast methylotrophic lifestyle. *BMC Biology*, 13(1), 80.
- Saeidi, S., Amin, N. A. S., & Rahimpour, M. R. (2014). Hydrogenation of CO₂ to value-added products—A review and potential future developments. *Journal of CO₂ utilization*, 5, 66-81.
- Santoni, E., Jakopitsch, C., Obinger, C., & Smulevich, G. (2004). Comparison between catalase-peroxidase and cytochrome c peroxidase. The role of the hydrogen-bond networks for protein stability and catalysis. *Biochemistry*, 43(19), 5792-5802.
- Savakis, P., & Hellingwerf, K. J. (2015). Engineering cyanobacteria for direct biofuel production from CO₂. *Current Opinion in Biotechnology*, 33, 8-14. <https://doi.org/10.1016/j.copbio.2014.09.007>
- Schrader, J., Schilling, M., Holtmann, D., Sell, D., Villela Filho, M., Marx, A., & Vorholt, J. A. (2009). Methanol-based industrial biotechnology: Current status and future perspectives of methylotrophic bacteria. *Trends in Biotechnology*, 27(2), 107-115.
- Schroer, K., Luef, K. P., Hartner, F. S., Glieder, A., & Pscheidt, B. (2010). Engineering the *Pichia pastoris* methanol oxidation pathway for improved NADH regeneration during whole-cell biotransformation. *Metabolic Engineering*, 12(1), 8-17.
- Sheehan, M., Bailey, C., Dowds, B., & McConnell, D. (1988). A new alcohol dehydrogenase, reactive towards methanol, from *Bacillus stearothermophilus*. *Biochemical Journal*, 252(3), 661-666.
- Songur, A., Ozen, O. A., & Sarsilmaz, M. (2010). The toxic effects of formaldehyde on the nervous system. In *Reviews of environmental contamination and toxicology* (pp. 105-118). Springer.
- Sperling, D. (2007). Beyond oil and gas: The methanol economy. *Energy Journal*, 28(1), 178-179. <http://search.ebscohost.com/login.aspx?direct=true&db=bth&AN=23789295&lang=zh-cn&site=ehost-live>
- Srivatsan, S. R., McFaline-Figueroa, J. L., Ramani, V., Saunders, L., Cao, J., Packer, J., Pliner, H. A., Jackson, D. L., Daza, R. M., Christiansen, L., Zhang, F., Steemers, F., Shendure, J., & Trapnell, C. (2020). Massively multiplex chemical transcriptomics at single-cell resolution. *Science*, 367(6473), 45-51.
- Subedi, K. P., Kim, I., Kim, J., Min, B., & Park, C. (2008). Role of GldA in dihydroxyacetone and methylglyoxal metabolism of *Escherichia coli* K12. *FEMS Microbiology Letters*, 279(2), 180-187.
- Sunga, A. J., & Cregg, J. M. (2004). The *Pichia pastoris* formaldehyde dehydrogenase gene (FLD1) as a marker for selection of multicopy expression strains of *P. pastoris*. *Gene*, 330, 39-47.
- Tuyishime, P., Wang, Y., Fan, L., Zhang, Q., Li, Q., Zheng, P., Sun, J., & Ma, Y. (2018). Engineering *Corynebacterium glutamicum* for methanol-dependent growth and glutamate production. *Metabolic Engineering*, 49, 220-231. <https://doi.org/10.1016/j.ymben.2018.07.011>
- Vartiainen, E., Blomberg, P., Ilmén, M., Andberg, M., Toivari, M., & Penttilä, M. (2019). Evaluation of synthetic formaldehyde and methanol assimilation pathways in *Yarrowia lipolytica*. *Fungal Biology and Biotechnology*, 6(1), 27.
- Velterop, J., Sellink, E., Meulenberg, J., David, S., Bulder, I., & Postma, P. (1995). Synthesis of pyrroloquinoline quinone in vivo and in vitro and detection of an intermediate in the biosynthetic pathway. *Journal of Bacteriology*, 177(17), 5088-5098.
- De Vries, G., Arfman, N., Terpstra, P., & Dijkhuizen, L. (1992). Cloning, expression, and sequence analysis of the *Bacillus methanolicus* C1 methanol dehydrogenase gene. *Journal of Bacteriology*, 174(16), 5346-5353.
- Vuilleumier, S., Chistoserdova, L., Lee, M.-C., Bringel, F., Lajus, A., Zhou, Y., Gourion, B., Barbe V., Chang J., Cruveiller S., Dossat C., Gillett W., Gruffaz C., Haugen E., Hourcade E., Levy R., Mangenot S., Muller E., Nadalig T., ... Lidstrom M. E. (2009). *Methylotrophicum* genome sequences: A reference blueprint to investigate microbial metabolism of C1 compounds from natural and industrial sources. *PLOS One*, 4(5), e5584.
- Wakayama, K., Yamaguchi, S., Takeuchi, A., Mizumura, T., Ozawa, S., Tomizuka, N., Hayakawa, T., & Nakagawa, T. (2016). Regulation of intracellular formaldehyde toxicity during methanol metabolism of

- the methylotrophic yeast *Pichia methanolica*. *Journal of Bioscience and Bioengineering*, 122(5), 545-549.
- Whitaker, W. B., Jones, J. A., Bennett, R. K., Gonzalez, J. E., Vernachio, V. R., Collins, S. M., Palmer, M. A., Schmidt, S., Antoniewicz, M. R., Koffas, M. A., & Papoutsakis, E. T. (2017). Engineering the biological conversion of methanol to specialty chemicals in *Escherichia coli*. *Metabolic Engineering*, 39, 49-59. <https://doi.org/10.1016/j.ymben.2016.10.015>
- Whitaker, W. B., Sandoval, N. R., Bennett, R. K., Fast, A. G., & Papoutsakis, E. T. (2015). Synthetic methylotrophy: engineering the production of biofuels and chemicals based on the biology of aerobic methanol utilization. *Current Opinion in Biotechnology*, 33, 165-175.
- Witthoff, S., Schmitz, K., Niefenführ, S., Nöh, K., Noack, S., Bott, M., & Marienhagen, J. (2015). Metabolic engineering of *Corynebacterium glutamicum* for methanol metabolism. *Applied and Environmental Microbiology*, 81(6), 2215-2225.
- Woolston, B. M., King, J. R., Reiter, M., Van Hove, B., & Stephanopoulos, G. (2018). Improving formaldehyde consumption drives methanol assimilation in engineered *E. coli*. *Nature Communications*, 9(1), 2387.
- Woolston, B. M., Roth, T., Kohale, I., Liu, D. R., & Stephanopoulos, G. (2018). Development of a formaldehyde biosensor with application to synthetic methylotrophy. *Biotechnology and Bioengineering*, 115(1), 206-215. <https://doi.org/10.1002/bit.26455>
- Wu, T.-Y., Chen, C.-T., Liu, J. T.-J., Bogorad, I. W., Damoiseaux, R., & Liao, J. C. (2016). Characterization and evolution of an activator-independent methanol dehydrogenase from *Cupriavidus necator* N-1. *Applied Microbiology and Biotechnology*, 100(11), 4969-4983.
- Xiaoding, X., & Moulijn, J. (1996). Mitigation of CO₂ by chemical conversion: Plausible chemical reactions and promising products. *Energy & Fuels*, 10(2), 305-325.
- Xu, Y., Ramanathan, V., & Victor, D. G. (2018). Global warming will happen faster than we think. Nature Publishing Group.
- Yang, X., Yuan, Q., Luo, H., Li, F., Mao, Y., Zhao, X., Du, J., Li, P., Ju, X., Zheng, Y., Chen, Y., Liu, Y., Jiang, H., Yao, Y., Ma, H., & Ma, Y. (2019). Systematic design and in vitro validation of novel one-carbon assimilation pathways. *Metabolic Engineering*, 56, 142-153.
- Yu, H., & Liao, J. C. (2018). A modified serine cycle in *Escherichia coli* converts methanol and CO₂ to two-carbon compounds. *Nature Communications*, 9, 9. <https://doi.org/10.1038/s41467-018-06496-4>
- Yurimoto, H. (2009). Molecular basis of methanol-inducible gene expression and its application in the methylotrophic yeast *Candida boidinii*. *Bioscience, Biotechnology, and Biochemistry*, 0903051357.
- Yurimoto, H., Kato, N., & Sakai, Y. (2005). Assimilation, dissimilation, and detoxification of formaldehyde, a central metabolic intermediate of methylotrophic metabolism. *The Chemical Record*, 5(6), 367-375. <https://doi.org/10.1002/tcr.20056>
- Yurimoto, H., Komeda, T., Lim, C. R., Nakagawa, T., Kondo, K., Kato, N., & Sakai, Y. (2000). Regulation and evaluation of five methanol-inducible promoters in the methylotrophic yeast *Candida boidinii*. *Biochimica et Biophysica Acta—Gene Structure and Expression*, 1493(1-2), 56-63.
- Zhan, C., Yang, Y., Zhang, Z., Li, X., Liu, X., & Bai, Z. (2017). Transcription factor Mxr1 promotes the expression of Aox1 by repressing glycerol transporter 1 in *Pichia pastoris*. *FEMS Yeast Research*, 17(4).
- Zhang, M., & Lidstrom, M. E. (2003). Promoters and transcripts for genes involved in methanol oxidation in *Methylobacterium extorquens* AM1. *Microbiology*, 149(4), 1033-1040.
- Zhang, W., Zhang, T., Wu, S., Wu, M., Xin, F., Dong, W., Ma, J., Zhang, M., & Jiang, M. (2017). Guidance for engineering of synthetic methylotrophy based on methanol metabolism in methylotrophy. *RSC Advances*, 7(7), 4083-4091. <https://doi.org/10.1039/C6RA27038G>
- Zhang, Y.-H. P. (2011). Simpler is better: High-yield and potential low-cost biofuels production through cell-free synthetic pathway biotransformation (SyPaB). *ACS Catalysis*, 1(9), 998-1009.

How to cite this article: Zhan, C., Li, X., Yang, Y., Nielsen, J., Bai, Z., & Chen, Y. (2021). Strategies and challenges with the microbial conversion of methanol to high-value chemicals. *Biotechnology and Bioengineering*, 1-14. <https://doi.org/10.1002/bit.27862>