



CHALMERS
UNIVERSITY OF TECHNOLOGY

Engineering *Saccharomyces cerevisiae* for the production and secretion of Affibody molecules

Downloaded from: <https://research.chalmers.se>, 2026-04-03 10:21 UTC

Citation for the original published paper (version of record):

Gast, V., Sandegren, A., Dunås, F. et al (2022). Engineering *Saccharomyces cerevisiae* for the production and secretion of Affibody molecules. *Microbial Cell Factories*, 21(1).
<http://dx.doi.org/10.1186/s12934-022-01761-0>


N.B. When citing this work, cite the original published paper.

RESEARCH

Open Access



Engineering *Saccharomyces cerevisiae* for the production and secretion of Affibody molecules

Veronica Gast¹, Anna Sandegren³, Finn Dunås³, Siri Ekblad³, Rezan Güler³, Staffan Thorén³, Marta Tous Mohedano¹, Mikael Molin¹, Martin K. M. Engqvist¹ and Verena Siewers^{1,2*} 

Abstract

Background: Affibody molecules are synthetic peptides with a variety of therapeutic and diagnostic applications. To date, Affibody molecules have mainly been produced by the bacterial production host *Escherichia coli*. There is an interest in exploring alternative production hosts to identify potential improvements in terms of yield, ease of production and purification advantages. In this study, we evaluated the feasibility of *Saccharomyces cerevisiae* as a production chassis for this group of proteins.

Results: We examined the production of three different Affibody molecules in *S. cerevisiae* and found that these Affibody molecules were partially degraded. An albumin-binding domain, which may be attached to the Affibody molecules to increase their half-life, was identified to be a substrate for several *S. cerevisiae* proteases. We tested the removal of three vacuolar proteases, proteinase A, proteinase B and carboxypeptidase Y. Removal of one of these, proteinase A, resulted in intact secretion of one of the targeted Affibody molecules. Removal of either or both of the two additional proteases, carboxypeptidase Y and proteinase B, resulted in intact secretion of the two remaining Affibody molecules. The produced Affibody molecules were verified to bind their target, human HER3, as potently as the corresponding molecules produced in *E. coli* in an in vitro surface-plasmon resonance binding assay. Finally, we performed a fed-batch fermentation with one of the engineered protease-deficient *S. cerevisiae* strains and achieved a protein titer of 530 mg Affibody molecule/L.

Conclusion: This study shows that engineered *S. cerevisiae* has a great potential as a production host for recombinant Affibody molecules, reaching a high titer, and for proteins where endotoxin removal could be challenging, the use of *S. cerevisiae* obviates the need for endotoxin removal from protein produced in *E. coli*.

Keywords: Recombinant protein production, Heterologous protein production, Secretion, Proteases, Pep4, Prc1, Prb1, Fed-batch, *Saccharomyces cerevisiae*, Yeast

Background

Affibody molecules are synthetic peptides that can be designed to bind with high affinity to other proteins. The part of the protein exhibiting binding properties

originates from the B domain of the immunoglobulin-binding region of the staphylococcal protein A (SpA) [1]. Nilsson et al. isolated the B domain of SpA and increased its stability, which led to the creation of the synthetic Z domain [2]. By randomizing 13 surface-exposed residues the specificity and affinity of the Z domain can be altered. Over the years, several Z domain variants have been generated with affinity to diagnostically relevant substrates, such as cancer markers [3–5]. A specific example of this

*Correspondence: siewers@chalmers.se

¹ Department of Biology and Biological Engineering, Chalmers University of Technology, Gothenburg, Sweden

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

is HER3. Elevated HER3 expression is associated with malignant cancer in ovarian, prostate, gastric, bladder, lung, melanoma and colorectal tissue [6]. Current applications for Affibody molecules are mostly within diagnostics and therapeutics where the main advantage of Affibody molecules compared to alternatives, like antibodies, is their small size which allows them to penetrate tissue more easily [7, 8].

Affibody molecules can be designed in a range of configurations, either containing solely Z-domains or Z-domains together with other peptide sequences. Affibody molecules composed of only the Z domain exhibit a relatively short half-life in the human body. Due to their small size, 6.7 kDa for a single Z domain, Affibody molecules are taken up by the kidneys and degraded [9]. To prolong the half-life of the Affibody molecules an albumin binding domain (ABD) can be added. The ABD originates from the GA3 module of streptococcal protein G and binds to serum albumin. Serum albumin, with a size of 67 kDa, is above the glomerular filtration barrier of the kidneys and adding ABD to Affibody molecules thus increases both their half-life and efficacy [5, 10]. Different configurations of the Z domain and the ABD domain, like the presence of multiple Z domains or an alternative localization of the ABD within the molecule, have shown to be effective in a range applications [7, 10, 11].

Currently, Affibody molecules are produced in the bacterial production host *Escherichia coli*. Bacteria are popular hosts for recombinant production due to rapid growth and high yields of the recombinant proteins [12]. However, recombinant protein production in *E. coli* leads to contamination with lipopolysaccharides from the bacterial cell wall, most of which exhibit endotoxin properties and can be difficult to remove during product purification [13, 14]. It is therefore of interest to explore other established and competitive production hosts like yeasts that do not have toxic components in their cell wall.

Yeasts are eukaryotic unicellular organisms, which secrete recombinant proteins and have been implemented as industrial production hosts for several pharmaceutical proteins [15, 16]. The most widely used yeast species for the industrial production of recombinant pharmaceutical proteins are *Saccharomyces cerevisiae* and *Komagataella phaffi* [16]. *K. phaffi* has been used as production host of a fusion construct of human serum albumin and a HER2 Affibody molecule; however, *S. cerevisiae* has not been tested as an Affibody molecule production host yet [17]. *S. cerevisiae* on the other hand is particularly known for its industrial production of insulin, human serum albumin, and hepatitis vaccines [18].

In this study, we tested an engineered *S. cerevisiae* strain for the production and secretion of three Affibody

molecules that bind to the cancer marker protein HER3 [6]. Initially, we found that *S. cerevisiae* degraded the produced Affibody molecules. We were able to identify several proteases responsible for this degradation, and upon removal of these, all three Affibody molecules were secreted in an intact state. We verified high-affinity HER3 binding by one of the secreted Affibody molecules and performed a fed-batch cultivation where a high final titer was reached thus demonstrating that *S. cerevisiae* is a competitive host for Affibody molecule production.

Results

The albumin-binding domain is degraded by *S. cerevisiae*

We first tested whether Affibody molecules could be produced and secreted by *S. cerevisiae*. We included three different Affibody molecules in this study. These three molecules have different configurations of either one or two Z_{HER3_1} domains, the Z domain variant that binds to HER3, and one albumin binding domain (ABD) (Fig. 1). *S. cerevisiae* strain B184k, previously evolved for high levels of protein secretion, was used as the host [19, 20]. Genes encoding the three different Affibody molecules shown in Fig. 1 were cloned into the backbone of a CPOT plasmid, thus generating the three plasmids pNatZACPOT (Z_{HER3_1} -ABD), pNatZ-ZACPOT (Z_{HER3_1} - Z_{HER3_1} -ABD), and pNatZAZCPOT (Z_{HER3_1} -ABD- Z_{HER3_1}) [21]. CPOT is a recombinant protein expression plasmid for use in *S. cerevisiae* *tpi1Δ* strains. The CPOT plasmid contains the *POT1* gene encoding triose phosphate isomerase from *Schizosaccharomyces pombe*, which partially complements the removal of the *TPI1* gene and restores the ability to grow on glucose as a carbon source. The partial complementation will lead to a high abundance of the plasmid in the cell, which is combined with the expression regulation of the recombinant gene by the native *TPI1* promoter and terminator [21]. The combination of the CPOT expression system and B184 has shown effective for high-level production of several

1. Z_{HER3_1} -ABD

2. Z_{HER3_1} - Z_{HER3_1} -ABD

3. Z_{HER3_1} -ABD- Z_{HER3_1}




Fig. 1 Schematic representation of domain configurations in the three Affibody molecules produced in yeast. The Z_{HER3_1} domain is the black peptide and the albumin binding domain (ABD) the colored peptide

recombinant proteins [19, 20]. The plasmids pNat-ZACPOT, pNatZZACPOT and pNatZAZCPOT were used to transform B184k and the positive transformants were grown for 48 h in liquid SD2xSCAA media [22]. The supernatant was analyzed by a reducing SDS-PAGE followed by western blot using antibodies against both the ABD and the Z domain (Fig. 2A–C). Although all three Affibody molecules were secreted the bands did not appear with the expected sizes on the blot. We expected a 12 kDa band for Z_{HER3_1} -ABD and 18.9 kDa for Z_{HER3_1} -ABD- Z_{HER3_1} and Z_{HER3_1} - Z_{HER3_1} -ABD. The supernatant derived from the Z_{HER3_1} -ABD- Z_{HER3_1} expressing strain showed three separate smaller bands on the anti-Z-domain blot with sizes compared to the Z_{HER3_1} -ABD standard around 12 kDa instead of one intact band seen in the Affibody standard around 18.9 kDa (Fig. 2B). The supernatant of the Z_{HER3_1} -ABD expressing strain showed two bands instead of one on the anti-Z-domain blot, which appeared around the expected size of 12 kDa (Fig. 2B). The anti-Z-domain blot for Z_{HER3_1} - Z_{HER3_1} -ABD showed an apparently intact band around the correct size of 18.9 kDa (Fig. 2B). Finally, the ABD domain showed none or little signal on the anti-ABD blot for any of the Affibody molecules (Fig. 2C). The absence of bands on the anti-ABD blots indicated an absence of intact ABD in the secreted Affibody molecules.

We hypothesized that the secreted molecules were degraded by *S. cerevisiae* proteases leading to fragmentation (Fig. 2B and C). We also checked if the presence of BSA might have an influence. Since the albumin-binding domain binds to serum albumin [5], and one of the components of the SD2xSCAA media that was used for cultivation was bovine serum albumin (BSA), we wanted to exclude any BSA-related interference. We tested our hypotheses by incubating one Affibody molecule standard for the Z_{HER3_1} -ABD in four different solutions. The solutions were fresh sterile medium, spent supernatant, demineralized water with 0.1% BSA and demineralized water. The spent supernatant is the supernatant of cultures in which yeast cells had been cultivated and were removed by centrifugation. If the absence of the intact ABD was due to proteolytic activity, we expected to see degradation by incubating the Affibody molecule standard in the spent supernatant. The spent supernatant for this experiment was derived from a 24-h culture of B184k producing α -amylase. After the incubation, the proteins were separated by reducing SDS-PAGE and analysed by western blotting against the ABD (Fig. 2D and E). The western blot shows intact bands for the fresh sterile medium, demineralized water with 0.1% BSA,




Fig. 2 SDS-PAGE and western blot analysis of supernatant of B184k expressing different Affibody molecules. **A** SDS-PAGE of supernatant after 48 h of cultivation of B184k producing three different Affibody molecules (cropped). The lane marked with ZA sup contains supernatant of B184k expressing Z_{HER3_1} -ABD, the lane marked with ZZA sup contains supernatant of B184k expressing Z_{HER3_1} - Z_{HER3_1} -ABD, the lane marked with ZAZ sup contains supernatant of B184k expressing Z_{HER3_1} -ABD- Z_{HER3_1} , the lane marked with ep contains supernatant of B184k with empty plasmid, the lane marked with ZA and st1 contains supernatant of Z_{HER3_1} -ABD standard (12 kDa) dissolved in water, the lane marked with ZAZ and st2 contains supernatant of Z_{HER3_1} -ABD- Z_{HER3_1} standard (18.9 kDa) dissolved in water. **B** Western blot against the Z-domain (cropped), the membrane was blotted with anti-Z-domain antibody (1:1000) followed by an anti-Mouse (1:5000) secondary antibody. **C** Western blot against the ABD (cropped), the membrane was blotted with anti-ABD antibody (3:1000) followed by an anti-Rabbit (1:5000) secondary antibody. **D** SDS-PAGE of supernatant of dissolved Z_{HER3_1} -ABD in different solutions (cropped). The lane marked with media contains fresh SD2xSCAA media with (+) and without (–) Z_{HER3_1} -ABD standard added, The lane marked with sup contains spent supernatant of a 24-h old culture of B184k in SD2xSCAA medium with (+) and without (–) Z_{HER3_1} -ABD standard added, the lane marked with BSA contains water with 0.1% BSA with (+) and without (–) Z_{HER3_1} -ABD standard added and the lane marked with H_2O , + contains Z_{HER3_1} -ABD standard in water. **E** Western blot against the ABD (cropped), the membrane was blotted with anti-ABD antibody (3:1000) followed by an anti-Rabbit (1:5000) secondary antibody

and demineralized water. For the spent supernatant, the stained SDS-PAGE shows two bands instead of one and there is no band visible on the western blot against the

ABD, which indicates degradation of the Affibody molecule in the spent supernatant.

Affibody molecules are degraded by aspartyl protease(s)

Proteolytic activity can drastically decrease recombinant protein yield; however, this can often at least partially be prohibited by the identification and removal of the responsible proteases [23]. First, we wanted to check if the degradation was a characteristic of the engineered yeast strain B184k, since a potential solution for the degradation would be to switch to another strain. We checked the supernatant of the parental strain of B184k, AAC [19]. The influence of expression of a recombinant protein on the degradation was also tested by the expression of α -amylase by AAC. Both B184k and AAC containing either pNatAmyCPOT or an empty CPOT plasmid were cultivated. The Affibody standard of $Z_{\text{HER3}_1\text{-ABD-Z}_{\text{HER3}_1}}$ was incubated in spent medium from those cultivations, and the mixture with the proteins was separated by reducing SDS-PAGE (Additional file 1: Fig. S1). We included supernatant after 24, 48, 72, and 96 h of cultivation and tested the presence of a protease inhibitor cocktail to confirm that the degradation was due to proteases. Degradation of the Affibody molecules was also observed in medium spent by the strain AAC (Additional file 1: Fig. S1). In the supernatant containing protease inhibitor cocktail, we did not observe degradation, confirming that the fragmentation of the Affibody molecules was a result of proteolytic activity. We observed a similar degradation pattern as for B184k. Degradation, however, seemed slightly elevated in supernatant derived from strains with recombinant α -amylase production, which could potentially indicate an induction of protease expression or activity upon recombinant protein production. Since the degradation was observed also in the parental strain of B184k we concluded that the degradation is not a characteristic of B184k. Therefore, we continued with B184k as the production host.

The next step was to identify the proteases responsible for the degradation. Protease inhibitors mostly block the activity of a specific class of proteases. Therefore, a mixture of different protease inhibitors is combined in protease inhibitor cocktails to ensure inhibition of all sort of proteases. Since in the experiment with AAC (Additional file 1: Fig. S1) a protease inhibitor cocktail showed effective against the degradation we tested the isolated inhibitors from that specific cocktail. The spent medium of B184k expressing α -amylase was incubated overnight with Affibody molecule standards and the individual components of the protease inhibitor cocktail. The mixture was subsequently analyzed by a reducing SDS-PAGE. The results of the SDS-PAGE show that the Affibody





Fig. 3 Protease inhibitor assay with spent supernatant of B184 and Affibody molecule standards. SDS-PAGE of spent supernatant with the indicated Affibody molecules and addition of different protease inhibitors (cropped). The supernatant is derived from a cultivation of B184k producing α -amylase grown for 24 h at 30 °C at 220 rpm. The Affibody molecule standards were incubated overnight at 30 °C at 220 rpm. The lane marked with pc contains the positive control with spent supernatant with the respective Affibody molecule and the complete protease inhibitor cocktail, the lane marked with nc contains the negative control with spent supernatant with the respective Affibody molecule, the lane marked with st contains the standard of the respective Affibody molecule in water, the lane marked with AEBSF contains spent supernatant with the respective Affibody molecule with 4-(2-aminoethyl) benzenesulfonyl fluoride hydrochloride 1 mM, the lane marked with aprotinin contains spent supernatant with the respective Affibody molecule with aprotinin 6.5 μ g/mL, the lane marked with bestatin contains spent supernatant with the respective Affibody molecule with bestatin hydrochloride 50 μ M, the lane marked with E-64 contains spent supernatant with the respective Affibody molecule with E-64 15 μ M, the lane marked with leupeptin contains spent supernatant with the respective Affibody molecule with leupeptin 20 μ M and the lane marked with pepstatin A contains spent supernatant with the respective Affibody molecule with pepstatin A 10 μ M

molecules were degraded in the presence of AEBSF, aprotinin, bestatin, E-64 or leupeptin (Fig. 3). Only incubation with pepstatin A prohibited degradation and resulted in Affibody molecules with the correct size (Fig. 3). Both the Affibody molecules $Z_{\text{HER3}_1\text{-Z}_{\text{HER3}_1\text{-ABD}}$ and $Z_{\text{HER3}_1\text{-ABD-Z}_{\text{HER3}_1}}$ clearly showed the absence of proteolytic processing on the SDS-PAGE in the presence of pepstatin A. For $Z_{\text{HER3}_1\text{-ABD}}$, the difference was less explicit which could be a result of the small size difference between the intact and degraded molecule. Pepstatin A inhibits aspartyl proteases, indicating that the protease(s) responsible for Affibody molecule processing are of this type.

The removal of several proteases results in the secretion of intact Affibody molecules

One of the major proteases in *S. cerevisiae* and a proven target for improving recombinant protein production yield is a vacuolar aspartyl protease, proteinase A [23–25]. We removed the *PEP4* gene encoding proteinase A in B184 using CRISPR/Cas9-based gene deletion




and expressed all three Affibody molecules in B184 *pep4Δ*. Z_{HER3_1} -ABD- Z_{HER3_1} produced by this strain was intact and showed a signal on western blots using antibodies against either the Z-domain or ABD (Fig. 4), indicating that proteinase A was indeed responsible for its cleavage in B184. In contrast, for Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD, the western blot against the Z-domain showed a small shift of the bands upon removal of *PEP4* but there was no band visible on the blot against the ABD (Fig. 4A). We suspected that at least one other protease was involved in cleaving the ABD domain of Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD in B184 *pep4Δ*.

We therefore removed two additional proteases in B184 *pep4Δ* in the hope of producing intact Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD, namely carboxypeptidase Y and proteinase B. Carboxypeptidase Y is a serine-based exopeptidase active in the vacuole and proteinase B, a serine-based endopeptidase also located to the vacuole. Proteinase B, together with proteinase A, plays an essential role in the activation of several vacuolar proteases including itself and carboxypeptidase Y. The *PRC1* gene encoding carboxypeptidase Y and the *PRB1* gene encoding proteinase B were removed in B184 and B184 *pep4Δ* (*pep4Δprc1Δ* and *pep4Δprb1Δ*) and a triple deletion strain (*pep4Δprc1Δprb1Δ*) was constructed. The single, the two double and the triple deletion strains were grown for 48 h in an aerated 24-well plate and the supernatant in which triple deletion strains expressing

the Affibody molecules were grown was analyzed for the presence of intact Affibody molecules. In the B184 *pep4Δprc1Δ* strain we observed intact Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD (Fig. 5A and B). Interestingly, the removal of solely *PEP4* during cultivation for 48 h instead of 96 h, resulted in very weak bands for the ABD for both Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD on the blot against the ABD. This indicates that the degradation in B184 *pep4Δ* could primarily occur at later cultivation stages. The single removal of either *PRC1* or *PRB1* did not result in a band on the western blot using antibody against the ABD. However, the combination of *pep4Δ* (proteinase A) with either *prb1Δ* (proteinase B) or *prc1Δ* (carboxypeptidase Y) resulted in the production of intact Z_{HER3_1} -ABD as well as intact Z_{HER3_1} -ABD- Z_{HER3_1} . The deletion of all three proteases (*pep4Δprc1Δprb1Δ*) resulted in titers comparable to the two double deletion strains (Fig. 5A and B).

Proteases fulfill a crucial role in the degradation of macromolecules in the vacuole and have even been reported to benefit aging [26, 27]. Therefore, the removal of one to three major proteases can have an impact on vacuolar homeostasis and cellular growth. We monitored the growth profiles of the protease-deficient strains in aerated 96-well plates. The protease deficiency led to a slight reduction of final OD₆₀₀ in the later stages of the growth, from 50 to 96 h, but overall growth of the deletion strains, even the triple deletion strain, was similar to B184 with the proteases intact. This was observed for both Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD (Fig. 5C and D).

Additionally, we were interested in identifying the cleavage site of the endopeptidase proteinase A in the ABD. It is known that proteinase A has a broad and variable activity for different ligands and seems to favor cut sites between adjacent hydrophobic residues [28, 29]. We incubated the Z_{HER3_1} -ABD- Z_{HER3_1} standard in spent supernatant for 24 h and analysed the mixture with mass spectrometry. The total ion chromatogram (TIC) showed four peaks, two large peaks and two smaller ones (Fig. 6A). The peptides within the two larger peaks were analysed. In the first peak, several fragments of Z_{HER3_1} -ABD- Z_{HER3_1} were identified where either the N- or C-terminus was intact and the other terminus was located within the ABD indicating a cut site of an endopeptidase (Fig. 6B). The peptides in the fourth peak were fragments which had the N-terminus in the ABD and the C-terminus in the second Z_{HER3_1} sequence (Fig. 6B). Based on these fragments it seems that proteinase A cleaves at several sites within the ABD and interestingly the C-terminal Z_{HER3_1} domain was also not intact.



Z_{HER3_1} -ABD- Z_{HER3_1} molecule produced by B184 *pep4* Δ shows similar binding kinetics compared to the molecule produced by *E. coli*

We next aimed to test whether the engineered *S. cerevisiae* strain is a competitive host for the Affibody molecule production with regards to product quality. Since the three Affibody molecules consist of the same two subunits, Z_{HER3_1} and ABD, we decided to include only one version in the subsequent studies as a proof of

concept. The Z_{HER3_1} -ABD- Z_{HER3_1} molecule produced and secreted by B184 *pep4* Δ was tested in a surface plasmon resonance binding assay with the substrate of Z_{HER3_1} , HER3. The Z_{HER3_1} -ABD- Z_{HER3_1} Affibody molecule has two binding sites for HER3; therefore, the bivalent binding kinetics were determined and presented as average value of duplicates in Table 1. The equilibrium dissociation constant of the first site has a value in the nano molar range, which indicates high




Table 1 Bivalent binding kinetics assay parameters of Z_{HER3_1} -ABD- Z_{HER3_1} produced by B184 *pep4Δ* and by the original host *E. coli*

Production host	k_a1 (1/Ms)	k_a2 (1/RUs)	k_d1 (1/s)	k_d2 (1/s)	K_D1 (M)	K_D2 (M)
<i>E. coli</i>	5.21×10^5	1.23×10^{-3}	1.43×10^{-2}	2.29×10^{-5}	2.75×10^{-8}	1.85×10^{-2}
<i>S. cerevisiae</i>	3.83×10^5	1.71×10^{-3}	1.79×10^{-2}	2.76×10^{-5}	4.68×10^{-8}	1.62×10^{-2}


k_a1 = association rate constant of the first site; k_a2 = association rate constant of the second site; k_d1 = dissociation rate constant of the first site; k_d2 = dissociation rate constant of the second site; K_D1 = equilibrium dissociation constant for the first site, K_D2 = equilibrium dissociation constant for the second site

affinity. The values of the Z_{HER3_1} -ABD- Z_{HER3_1} molecules secreted by *E. coli* and *S. cerevisiae*, respectively, show comparable kinetics which shows that the molecules produced by *S. cerevisiae* are fully functional (Table 1, Additional file 1: Fig. S2).

B184 *pep4Δ* produces a high titer of Z_{HER3_1} -ABD- Z_{HER3_1} in a fed-batch bioreactor

For *S. cerevisiae* to be a suitable host for Affibody molecule production it must be able to produce competitive titers of Affibody molecules. So next, we wanted to assess the productivity of *S. cerevisiae* in bioreactors. We decided to use B184 *pep4Δ* producing Z_{HER3_1} -ABD- Z_{HER3_1} as a proof of concept. To exclude any major impact of the deletion of *pep4Δ* on cellular growth of B184 while producing Z_{HER3_1} -ABD- Z_{HER3_1} we tested B184 and B184 *pep4Δ* in a micro-cultivation experiment in an aerated 96-well plate (Additional file 1: Fig. S3). Deletion of *pep4Δ* conferred only a minor impact on growth (Additional file 1: Fig. S3). After this

confirmation the productivity for Z_{HER3_1} -ABD- Z_{HER3_1} of B184 *pep4Δ* was tested in a bioreactor experiment. The cultivation set-up was a batch fermentation followed by fed-batch fermentation. During the batch phase, a specific growth rate of 0.31 h^{-1} was measured (Additional file 1: Fig. S4) which is the same growth rate as for B184 carrying an intact *PEP4* gene and producing α -amylase which confirmed that B184 *pep4Δ* did not suffer from a growth impairment [20]. The feeding of a low-glucose feed into the bioreactors was started after 36 h and was switched to a high-glucose feed after 130 h for the remaining duration of the fermentation until 180 h. At the end of the fed-batch cultivation, the biomass concentration exceeded 100 g/L (Fig. 7A). We also observed an increasing trend of Affibody molecule titer with the duration of the experiment especially after the switch to the high-glucose feed (Fig. 7A). During the cultivation, minor production of byproducts was observed. Ethanol and glycerol were present in the medium after the batch phase



but were rapidly consumed after the feed was started (Fig. 7B). At the end of the fed-batch, residual glycerol increased again and reached a final concentration of 10 g/L (Fig. 7B). We determined the exact quantity of Z_{HER3_1} -ABD- Z_{HER3_1} at three timepoints using a BLI based method. The results showed a final titer of 530 mg/L (Fig. 7C).

Discussion

In this study we examined *S. cerevisiae* as a host for Affibody molecule production. Currently, the molecules are mainly produced in *E. coli*. In the current study, we present engineered protease-deficient *S. cerevisiae* as a suitable alternative.

Our results show that all three Affibody molecules were produced and secreted by *S. cerevisiae* after the removal of only proteinase A or with the additional removal of proteinase B or carboxypeptidase Y. Removal of only proteinase A was sufficient for the production of intact Z_{HER3_1} -ABD- Z_{HER3_1} and a small amount of both Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD in the case of a shorter cultivation of 48 h instead of 96 h. We suspect that a longer duration of the experiment would lead to more degradation. The additional removal of carboxypeptidase Y and/or proteinase B is required for the intact production of Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD, respectively. The ABD is in Z_{HER3_1} -ABD- Z_{HER3_1} flanked by the two Z_{HER3_1} domains, which potentially protects the ABD from exopeptidase activity from proteases like carboxypeptidase Y. This would explain why we observed intact Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD only upon removal of carboxypeptidase Y in proteinase A deficient B184 *pep4Δ*. Additionally, we confirmed that the removal of carboxypeptidase Y does not further improve the production of Z_{HER3_1} -ABD- Z_{HER3_1} in proteinase A-deficient B184 *pep4Δ* (Additional file 1: Fig. S5). Interestingly, we also observed production of intact Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD upon removal of proteinase B, which is an endopeptidase, in B184 *pep4Δ* and the additional deletion of proteinase B in B184 *pep4Δprc1Δ* (lacking proteinase A and carboxypeptidase Y) or carboxypeptidase Y in B184 *pep4Δprb1Δ* (lacking proteinase A and proteinase B) did not influence the titer of intact Z_{HER3_1} -ABD and Z_{HER3_1} - Z_{HER3_1} -ABD. This observation can be explained by the fact that proteinase A, proteinase B and carboxypeptidase Y are all three vacuolar proteases that before reaching their final destination, the vacuole, pass through the secretory pathway as inactive precursors with a prepro-signalpeptide [30]. The pre-signal peptide is removed after arrival in the ER and the pro-signal peptide during trafficking to or upon arrival in the

vacuole [30]. Proteases can have multiple pro-signal peptides, which prohibits the proteases to reach their active form. Proteinase A and B are responsible for the removal of pro-signal peptide(s) on other vacuolar proteases and are therefore a central part of the activation cascade of vacuolar proteases.

For maturation of our deletion targets proteinase A, proteinase B and carboxypeptidase Y, activity of both proteinase A and B is required to reach their activate form. The maturation of proteinase A starts with an auto activation step to the pseudo form upon entering the vacuole, which is followed by removal of a second pro-signal peptide by proteinase B [31, 32]. The intermediate form of the protease with one pro-signal peptide and one already removed is called the pseudo-form. The exact activation trigger for the proteases remains unknown but has been linked to the acidic pH in the vacuole [33, 34]. Proteinase B has two pro-peptides, one at the N-terminus and another at the C-terminus. The removal of the N-terminal pro-peptide is autocatalytic and occurs in the Golgi apparatus, and the C-terminal pro-peptide is removed by proteinase A [35]. Carboxypeptidase Y has an N-terminal pro-peptide which is partly removed by proteinase A and the other part by proteinase B upon arrival in the vacuole.

As explained here, proteinase B catalyzes the activation of carboxypeptidase Y and based on our results we expect that proteinase B fulfills a role in the activation of proteinase A and carboxypeptidase Y rather than to recognize Affibody molecules as a substrate. We expect, based on our results, that in a *pep4Δprb1Δ* mutant carboxypeptidase Y remains in an inactive pro-form, which limits the capacity of carboxypeptidase Y to degrade the ABD [36]. We, however, observe digestion of the ABD by strain B184 *pep4Δ*. It has been shown that in a *PEP4* deficient *S. cerevisiae* strain pro-proteases of both carboxypeptidase Y and proteinase B accumulate, alongside the reduction of their activity [37]. The combination of deleting proteinase A and proteinase B for improved recombinant protein production has been reported for several microbial hosts, including *S. cerevisiae* [25, 38, 39]. Removal of solely *PEP4* has however also proven effective for production of several proteins [25, 41, 42].

The proteases travel through the secretory pathway to reach their destination, the vacuole. The initiation of the activation should start in the proximity of the vacuole or upon transport into this organelle [40]. Proteinase A and B are trafficked directly from the Golgi apparatus to the vacuole, but carboxypeptidase Y travels through a multivesicular body intermediate [26]. Even though the proteases are targeted for the vacuole, residual secretion

has been reported and linked to the glucose concentration in the media. In the experiments performed in this study 2% glucose concentration in the media was used, a concentration that previously was shown to result in extracellular activity of proteinase A [34]. This supports the observations of proteinase A activity in the spent supernatant.

Another interesting aspect is the propensity of the proteases for cleavage of ABD. Since potentially even pro-Prc1 and pseudo-Prc1 seem to digest ABD, it could well be that even partly active yeast proteases retain a high propensity for cleaving the ABD as a defense mechanism against bacteria. Proteolytic activity targeting the ABD was previously found in bacterial hosts [43].

The mass spectrometry performed for the analysis of Z_{HER3_1} -ABD- Z_{HER3_1} showed that the molecular weight of the molecule produced by *S. cerevisiae* consisted of an extra 700 Da compared to the molecule produced by *E. coli*. This is visible on several SDS-PAGEs and western blots (Figs. 4, 5, Additional file 1: Figs. S1 and S5). In the construct expressed by *S. cerevisiae*, a Kex2 recognition site is located behind the α -leader sequence and is processed in the ER. Between the Kex2 site and the start of the protein, a linker peptide with the amino acid sequence Glu-Glu-Gly-Glu-Gly-Ser-Met was inserted, which exhibits a molecular weight of 737 Da and thus explains the additional mass of the produced Z_{HER3_1} -ABD- Z_{HER3_1} . Since the binding kinetics indicate no interference of this additional peptide, we did not include the removal of the linker peptide in the scope of this study, but for future applications its removal may be desirable.

As a final experiment to show the potential of *S. cerevisiae* as interesting production host for Affibody molecules, B184 *pep4Δ* producing Z_{HER3_1} -ABD- Z_{HER3_1} was grown in fed-batch bioreactors. During the fed-batch cultures, buildup of residual glycerol was observed. The quantity of glycerol accumulating showed high variation between the reactors. Glycerol is a by-product of fermentation to reoxidize NADH to counteract redox imbalances. However, we do not expect this to be the cause of the buildup [44]. During later stages of the fed-batch, the feed was controlled by the dissolved oxygen in the reactor leading to pulse feed to ensure respiratory growth. Besides, in the case of alcoholic fermentation, there should be buildup of ethanol as well [45]. Therefore, the oxygen shortage does not seem to be the source of the potential redox imbalance. Besides redox imbalance, another explanation could be the expression platform used. *TPII* deficiency, which is the deficiency complemented by the CPOT plasmid, is known to result in the

accumulation of high levels of glycerol [46]. During the later stages of the experiment, this partial complementation could become limiting leading to the accumulation of excess glycerol. Nevertheless, high biomass concentration and Affibody molecule titer were achieved during the fermentation. The titer that was reached was 0.530 g/L. For *E. coli* expression titers for different recombinant proteins are reported in the range from 0.25 g/L to 8.5 g/L [47–49]. The titer of Z_{HER3_1} -ABD- Z_{HER3_1} is within this range and can be the starting point for improvement by further engineering *S. cerevisiae* or by optimizing the cultivation method in a sequential study.

The results of this study indicate that *S. cerevisiae* shows potential as a host for Affibody molecule production. Here, we tested only three different Affibody molecules, and for the future it would be interesting to expand the selection of Affibody molecules and potentially test more complex Affibody molecule based proteins like AffiMabs [7]. Secondly, our experiments show that the removal of proteases can be a powerful approach to increase productivity of heterologous proteins in *S. cerevisiae*.

Materials and methods

Strains and plasmids

Two previously constructed *S. cerevisiae* strains were used in this study, CEN.PK 530.1C [*MAT α* *URA3 HIS3 LAU2 TRP1 SUC2 MAL2-8^c tpi1(41-707)*] (AAC) and B184k. In a previous study, AAC was evolved for improved protein production by UV-mutagenesis which led to the construction of strain B184k [19]. B184k was shown to be an effective host for the production of several recombinant proteins in combination with the CPOT plasmid [19]. The CPOT plasmid contains the *Schizosaccharomyces pombe* gene *POT1*, which partially complements the *TPII* deficiency. The recombinant protein expression cassette contains the native *TPII* promoter and terminator and an α -leader sequence. The empty CPOT is a CPOT plasmid without the recombinant protein gene. The previously constructed pNatAmyCPOT was cut by restriction digest with enzymes XhoI and KpnI to remove only the α -amylase gene. Genes for Z_{HER3_1} -ABD, Z_{HER3_1} - Z_{HER3_1} -ABD and Z_{HER3_1} -ABD- Z_{HER3_1} were synthesized by GenScript. The Affibody molecule genes were codon optimized for expression in *S. cerevisiae* but without repetitive DNA sequences to reduce the risk of homologous recombination within the ORFs. The three Affibody molecule genes were amplified with homologous overhangs of the α -leader and the CPOT backbone using primers presented in Additional file 1: Table S1. The backbone and

Affibody molecule genes were assembled by Gibson assembly and the final constructs verified by sequencing using primer #7 that binds in the α -leader sequence. All primers for the construction of the plasmids can be found in Additional file 1: Table S1. After the plasmid construction, the newly assembled CPOT plasmids, pNatZACPOT, pNatZZACPOT and pNatZAZCPOT were used to transform AAC and B184k. B184k still contained a kanamycin resistance marker in the *TPII* gene. In AAC, the marker had been removed previously. The marker has two flanking *loxP* sites. pSH66 from the Euroscarf deletion marker set was used to remove the kanamycin resistance marker [50]. pSH66 contains an expression cassette for Cre recombinase under control of the *GALI* promoter and a nourseothricin resistance gene. After the strains had been transformed with the pSH66 plasmid with selection for nourseothricin, the positive transformants were streaked out on solid media with YPGal with nourseothricin to activate Cre recombinase expression. Removal of the kanamycin marker was confirmed by absence of growth on solid media with G418. *PEP4*, *PRC1* and *PRB1* were deleted by using plasmids pECAS9-gRNA-kanMX-tPEP4, pECAS9-gRNA-kanMX-tPRC1, pECAS9-gRNA-kanMX-tPRB1, which contain both a *cas9* gene and a gRNA expression cassette [51]. The plasmids pECAS9-gRNA-kanMX-tPEP4, pECAS9-gRNA-kanMX-tPRC1 and pECAS9-gRNA-kanMX-tPRB1 were constructed using pECAS9-gRNA-kanMX-tHFD1 as template [51]. First, the backbone of pECAS9-gRNA-kanMX was obtained by linearizing pECAS9-gRNA-kanMX-tHFD1 by digestion with MunI and EcoRI. The 'left' fragment was constructed with primer #14 in combination with either #9 (*PEP4*), #11 (*PRC1*) or #13 (*PRB1*) and the 'right' fragment was constructed with primer #15 in combination with either #8 (*PEP4*), #10 (*PRC1*) or #12 (*PRB1*). The correct assembly of the plasmids was confirmed by sequencing using primer #16. The genes were deleted in B184 pNatZACPOT, B184 pNatZZACPOT and B184 pNatZAZCPOT. The genomic deletions were verified using forward primers #17 and #18 with reverse primer #19 for *PEP4*, forward primers #20, #21 and reverse primer #22 for *PRC1*, and forward primers #23, #24 and reverse primer #25 for *PRB1*. The CRISPR plasmids were removed by subsequent cultivation in liquid YPD confirmed by absence of growth on solid media with G418. All the primers are presented in Additional file 1: Table S1 and repair fragments for CRISPR in Additional file 1: Table S2. The oligonucleotides were aligned by heating the two corresponding oligonucleotides to 98 degrees for 5 min in equimolar amounts and let to mixture cooldown to room temperature. *E. coli* DH5 α was

Table 2 Yeast strains used in this study

Strain	Genotype	References
AAC	CEN.PK113-7D (<i>MATa URA3 HIS3 LEU2 TRP1 MAL2-8^c SUC2 tpi1(41-707)::loxP-loxP</i>)	[53]
B184k	UV-mutant of AACK (<i>tpi1(41-707)::loxP-kanMX4-loxP</i>)	[19]
B184	UV-mutant of AACK (<i>tpi1(41-707)::loxP-loxP</i>)	This study
B184 <i>pep4Δ</i>	B184 <i>pep4Δ</i>	This study
B184 <i>prc1Δ</i>	B184 <i>prc1Δ</i>	This study
B184 <i>prb1Δ</i>	B184 <i>prb1Δ</i>	This study
B184 <i>pep4Δ prc1Δ</i>	B184 <i>pep4Δ prc1Δ</i>	This study
B184 <i>pep4Δ prb1Δ</i>	B184 <i>pep4Δ prb1Δ</i>	This study

Table 3 Plasmids used in this study

Plasmid	Description	References
pECAS9-gRNA-kanMX-tHFD1	2μ vector with <i>kanMX</i> marker expressing eCas9 under the <i>TEF1</i> promoter and <i>CYC1</i> terminator and the gRNA targeting <i>HFD1</i> under the <i>SNR52</i> promoter	[51]
pECAS9-gRNA-kanMX-tPEP4	2μ vector with <i>kanMX</i> marker expressing eCas9 under the <i>TEF1</i> promoter and <i>CYC1</i> terminator and the gRNA targeting <i>PEP4</i> under the <i>SNR52</i> promoter	This study
pECAS9-gRNA-kanMX-tPRC1	2μ vector with <i>kanMX</i> marker expressing eCas9 under the <i>TEF1</i> promoter and <i>CYC1</i> terminator and the gRNA targeting <i>PRC1</i> under the <i>SNR52</i> promoter	This study
pECAS9-gRNA-kanMX-tPRB1	2μ vector with <i>kanMX</i> marker expressing eCas9 under the <i>TEF1</i> promoter and <i>CYC1</i> terminator and the gRNA targeting <i>PRB1</i> under the <i>SNR52</i> promoter	This study
pNatAmyCPOT	2μ vector with cassette expressing <i>POT1</i> gene from <i>S. pombe</i> and an expression cassette with α-leader sequence and α-amylase gene under native <i>TPI1</i> promoter and terminator	[21]
pCPOT	2μ vector with cassette expressing <i>POT1</i> gene from <i>S. pombe</i> and an expression cassette with native <i>TPI1</i> promoter and terminator without recombinant protein gene	[21]
pNatZACPOT	2μ vector with cassette expressing <i>POT1</i> gene from <i>S. pombe</i> and an expression cassette with α-leader sequence and <i>Z_{HER3_1}-ABD</i> gene under native <i>TPI1</i> promoter and terminator	This study
pNatZZACPOT	2μ vector with cassette expressing <i>POT1</i> gene from <i>S. pombe</i> and an expression cassette with α-leader sequence and <i>Z_{HER3_1}-Z_{HER3_1}-ABD</i> gene under native <i>TPI1</i> promoter and terminator	This study
pNatZAZCPOT	2μ vector with cassette expressing <i>POT1</i> gene from <i>S. pombe</i> and an expression cassette with α-leader sequence and <i>Z_{HER3_1}-ABD-Z_{HER3_1}</i> gene under native <i>TPI1</i> promoter and terminator	This study

used for plasmid amplification. The transformation protocol used for *E. coli* was according to a known protocol [52]. The strains and plasmids used and constructed in this study are listed in Tables 2 and 3.

Media and culture conditions

The media used for *S. cerevisiae* strain construction were YPD, YPGal, YPE and YPEG. The experiments were always performed at 30 °C and for liquid cultures at 220 rpm. YPD medium contained 10 g/L yeast extract, 20 g/L peptone, and 20 g/L glucose and was used for regular cultures. YPGal media contained 10 g/L yeast extract, 20 g/L peptone, and 20 g/L galactose and was used for induction of the Cre recombinase gene on pSH66. For the selection of the *kanMX* marker on the CRISPR plasmid, 200 mg/L G418 was added to the YPD medium. For the selection of pSH66-containing cells,

100 mg/L nourseothricin sulfate was added to YPD and YPGal medium. The YPE medium contained 10 g/L yeast extract, 20 g/L peptone, 20 g/L absolute ethanol and was solely used as a solid medium. For liquid cultivations, 30 g/L glycerol was added, and the medium referred to as YPEG. Both YPE and YPEG were only used for *S. cerevisiae* strains without CPOT plasmids since those are unable to ferment glucose as the sole carbon source [54]. To solidify the media 20 g/L agar (Merck Millipore) was added. The protein expression and physiological experiments were performed in SD2xSCAA media at 30 °C and 220 rpm. SD2xSCAA medium contained 20 g/L glucose, 6.9 g/L yeast nitrogen base without amino acids, 190 mg/L Arg, 400 mg/L Asp, 1260 mg/L Glu, 130 mg/L Gly, 140 mg/L His, 290 mg/L Ile, 400 mg/L Leu, 440 mg/L Lys, 108 mg/L Met, 200 mg/L Phe, 220 mg/L Thr, 40 mg/L Trp, 52 mg/L Tyr, 380 mg/L Val, 1 g/L BSA,

5.4 g/L Na_2HPO_4 , and 8.56 g/L $\text{NaH}_2\text{PO}_4 \cdot \text{H}_2\text{O}$ with a pH of 6.4. Cells for protein production experiments were grown at 30 °C at 220 rpm in aerated 24-well plates CR1224 (Bioscreen) with a volume of 2.5 mL and a start OD_{600} of 0.1 or in 14 mL-cultivation tubes with a volume of 2 mL. For the binding assay experiment, a volume of 200 mL of SD2xSCAA, inoculated with an overnight culture to an OD of 0.1, was cultivated for 96 h at 30 °C at 220 rpm. The supernatant was stored at -80 °C before being sent to Affibody AB on dry ice. The *E. coli* cells were grown in Luria-Bertani (LB) media at 37 °C and 200 rpm. Selection media contained 80 mg/L ampicillin.

Molecular biology techniques

Saccharomyces cerevisiae strains were transformed using the LiAc/SS carrier method [55]. One μg of DNA was used for the transformation of plasmids and an additional 1 to 2 μg repair fragment if required. To verify deletions or test for the presence of the CPOT plasmids, colony PCR using SapphireAmp fast PCR mix (TaKaRa Bio) was performed. For DNA construction, Phusion High Fidelity DNA polymerase (Thermo Scientific) or Herculase II Fusion DNA polymerase (Aligent) was used. Restriction digestion was done using FastDigest (Thermo Scientific) products. All techniques were used according to the manufacturers' protocols unless stated otherwise.

Growth profiler experiments

Three independent transformation colonies per strain were grown for 24 h in 1 mL SD2xSCAA media in a 14 mL-cultivation tube. Those precultures were used to inoculate the main cultures of the growth experiment in technical triplicates with a starting OD of 0.01. The *S. cerevisiae* strains were cultivated for 96 h in 250 μL SD2xSCAA media at 30 °C and 1200 rpm in 96-well plates (EnzyScreen CR1496d). The growth curves were measured using a Growth Profiler 960 (EnzyScreen).

Protease activity experiments

For the incubation experiments, supernatants of cultivations were used. The initial cultivation was a 24 h-cultivation, if not mentioned otherwise, of a single colony in 2 mL of SD2xSCAA media. After the incubation, the culture was harvested by centrifugation at 6000 rpm for 3 min. The supernatant was kept on ice or frozen at -20 °C. The incubation was performed in 1.5-mL Eppendorf tubes with 200 μL volume of supernatant. As positive controls, standards of purified Z_{HER3_1} -ABD (1.97 mg/mL), Z_{HER3_1} - Z_{HER3_1} -ABD (0.77 mg/mL) and Z_{HER3_1} -ABD- Z_{HER3_1} (1.34 mg/mL) provided by Affibody AB were used with a concentration of 0.01 g/L or stated

otherwise. The experiments with the protease inhibitors were done by adding the protease inhibitor before adding the Affibody standard at the concentration suggested by the supplier. We used the Halt Protease Inhibitor Cocktail EDTA-free (100x) (Thermo Fisher) and 4-(2-aminoeethyl) benzenesulfonyl fluoride hydrochloride (AEBSF) (final concentration 1 mM), aprotinin (final concentration 6.5 $\mu\text{g}/\text{mL}$), bestatin hydrochloride (final concentration 50 μM), leupeptin (final concentration 20 μM), E-64 (final concentration 15 μM), and pepstatin A (final concentration 10 μM). All these chemicals were purchased from Merck.

SDS-PAGE and western blotting

The samples and controls were loaded and separated with reducing SDS-PAGE. Stain free 4–20% gels were used (Bio-rad). The proteins were transferred to 0.45-micron PVDF membranes (Bio-rad) using the Trans-Blot Turbo transfer system (Bio-rad). The blot was blocked using the Western blocker solution (Sigma Aldrich) and incubated in with either anti-Z-domain (2.87 mg/mL) (1:1000) or anti-ABD (1 mg/mL) (3:1000) antibodies, both obtained from Affibody AB followed by incubation with either anti-mouse antibody (1:5000) for anti-Z-domain or anti-rabbit (1:5000) antibody for anti-ABD antibodies, respectively. Both secondary antibodies were HRP-conjugated and visualized by using West Pico Plus HRP substrate (Thermo Fischer) and measured with a ChemidoC XRS image analyzer (Bio-Rad). As positive controls, standards of purified Z_{HER3_1} -ABD (1.97 mg/mL), Z_{HER3_1} - Z_{HER3_1} -ABD (0.77 mg/mL) and Z_{HER3_1} -ABD- Z_{HER3_1} (1.34 mg/mL) provided by Affibody AB were used with a concentration of 0.01 g/L or stated otherwise.

Mass spectrometry analysis

Intact mass analysis was performed on an Agilent 1200 LC system equipped with a C8 RP-HPLC column (Poroshell 300SB-C8, 5 μm , 2.1 \times 75 mm, Agilent) and coupled to a high-resolution Q-TOF (Bruker Maxis Impact). Buffer A was 0.1% formic acid in 10% acetonitrile, buffer B was 0.1% formic acid in 90% acetonitrile and the column temperature was 80 °C. Proteins were eluted from the LC column by using a linear gradient of solvent B from 5 to 75% over 5 min at a flow rate of 0.3 mL/min. The MS analysis was performed with an ESI source and in positive mode. MS was set in a MS only mode. The following MS parameters were used: mass range 500–4500, capillary 4500 V, nebulizer 1.2 Bar, drying gas 8.0 L/min and temperature 200 °C. The MS data were deconvoluted in Compass DataAnalysis (Bruker,

version 4.4). The compound spectra were first smoothed using the Savitzky Golay algorithm with a smoothing width of 0.2 Da and 3 cycles. Then the spectrum baseline was subtracted with 0.8 flatness. Afterwards, the raw spectra were deconvoluted using the MaxEnt algorithm. For the native proteins, the spectrums were deconvoluted within the mass range of 1000–25,000 and a resolving power of 1250. The observed masses were compared to theoretical masses of different fragment sequences with GPMW version 12.50 with a precision in Da at 0.5. Only fragment sequences with a mass accuracy ≤ 2 Da were reported.

Binding kinetics analysis

Surface plasmon resonance (SPR) analysis by Biacore 8 K (Cytiva, Marlborough, MA) was performed at 25 °C in a run buffer of HBS-EP (0.01 M HEPES pH 7.4, 0.15 M NaCl, 3 mM EDTA, 0.005% v/v Surfactant P20, Cytiva) and with 15 mM HCl as regeneration solution. Recombinant human ErbB3/Her3 Fc (R&D Systems, Minneapolis, MN) was immobilized by standard amine coupling (1-ethyl-3-(3-dimethylaminopropyl)carbodiimide (EDC) and N-hydroxysuccinimide (NHS)) on a CM5 sensor chip (Cytiva, Marlborough, MA) at ~ 1000 RU. The coated chip was preconditioned by three regeneration rounds to stabilize surfaces prior to injection of analyte. Binding of Affibody molecules to HER3 was analyzed by single cycle kinetics via injection of analyte at five different concentrations of purified Affibody molecule (1.25, 2.5, 5, 10 and 20 nM) over immobilized HER3/Fc. The experiment was performed in duplicates. Biacore Insight Evaluation Software was used to process, analyze and fit data.

Fed-batch bioreactor cultivations

For the Bioreactor experiments, the Na_2HPO_4 and $\text{NaH}_2\text{PO}_4 \cdot 2\text{H}_2\text{O}$ were replaced by KH_2PO_4 with a concentration of 2 g/L in the SD2xSCAA media. The batch fermentations were performed in 1-L bioreactors (DasGip) with a start OD_{600} of 0.01 in 500 mL of SD2xSCAA media. The conditions in the reactors were controlled and maintained at a pH of 6.0 by 4 M KOH, agitation at 600 rpm, temperature at 30 °C degrees and airflow of 30 L/h. The batch experiments were performed in biological quadruplicates. After all the glucose was consumed, culture broth was removed until a start volume of 250 mL. The conditions of the fed-batch were set to a pH of 6.0, maintained by 4 M KOH and 3 M HCl, agitation was set to 600 rpm, the temperature was set to 30 °C degrees, a constant airflow of 18 L/h and an exponentially increasing feed rate of 0.05 h^{-1} . When the dissolved oxygen in the bioreactor would decrease below

30% the airflow and agitation would increase. The airflow would increase to 48 L/h and agitation would increase to 1000 rpm. When both the agitation and airflow were at their maximum levels the feeding was changed to a pulse feed controlled by the dissolved oxygen level. The feed would stop if the dissolved oxygen was below 25% which resulted in a pulsed feeding rate around 8 mL/h. Two types of media were used for the fed-batch experiments. The first medium was the low-glucose feed with 200 g/L, 69 g/L yeast nitrogen base without amino acids, 50 g/L casamino acids (Formedium), 1 g/L BSA, and 20 g/L KH_2PO_4 . After adding an average of 320 mL of the low glucose feed the low-glucose feed was replaced by a high-glucose feed with the same composition as the low-glucose feed except for a glucose concentration of 400 g/L. The fermentation was ended after addition of 230 mL of the high-glucose feed. The fed-batch experiments were conducted in biological triplicates.

Affibody molecule quantification by Bio-Layer Interferometry (BLI)

The broth samples were centrifugated for 20 min at 4000 rpm to remove the biomass and the supernatant was stored at -80 °C. After thawing some precipitation appeared in the samples. To redissolve the precipitation the samples were 1:1 diluted in 1 NaCl in two duplicates per fermenter per sample point and transported on dry ice. Upon arrival the sample were thawed and either not diluted or diluted 1:3.75, 1:7.5 in sample diluent (1 \times PBS, 0.1% BSA, 0.02% Tween 20) based on the expected concentration. 225 μL of the pre-dilution was mixed with either 75 μL sample diluent (unspiked) or 75 μL of 100 $\mu\text{g}/\text{mL}$ $Z_{\text{HER3}_1\text{-ABD}}\text{-}Z_{\text{HER3}_1}$ (produced by *E. coli*) (spiked) was added resulting in a 1:5, 1:10 or 1:1.33 final dilution. 200 μL of the spiked and unspiked sample was moved to a black flat-bottomed 96-well plate (Greiner, #655209) together with 200 μL of the samples for the standard curve, 200 μL sample diluent (reference well) and 200 μL cultivation medium (negative control). High precision streptavidin (SAX) biosensors (ForteBio, #18–0037) which were previously loaded offline (Technical Note #10, ForteBio) with 50 $\mu\text{g}/\text{mL}$ biotinylated HSA (Recombunin Prime, Albumedix; Thermo Fisher #A39259) (MCR 1:1) were placed for 10 min in sample diluent before the measurements. The data was acquired using an Octet HTX system and Octet Data Acquisition software (ForteBio, ver. 12). The measurement settings were as follows: 8 channel read head, 5.0 Hz acquisition rate, quantitation step: 120 s, 400 rpm, plate temperature 30 °C. The quantity was determined based on the six-point standard curve with $Z_{\text{HER3}_1\text{-ABD}}\text{-}Z_{\text{HER3}_1}$

produced by *E. coli* (100 µg/mL–3.13 µg/mL). Data Analysis HT (ForteBio, ver. 12) was used to process the data. The reference well signal was subtracted from the samples and standard curve, and results with 70–143% spike recovery were included in the calculations. The assay was run twice for each sample.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12934-022-01761-0>.

Additional file 1: Figure S1. SDS PAGE (cropped) of degradation experiment with AAC and B184k. **Figure S2.** Binding assay kinetics of Z_{HER3_1}-ABD-Z_{HER3_1}. **Figure S3.** Growth profiles of B184 and B184 *pep4Δ* while producing Z_{HER3_1}-ABD-Z_{HER3_1}. **Figure S4.** Semi-log plot of cell dry weight measurements during batch fermentation. **Figure S5.** Western blot against the ABD (cropped) of Z_{HER3_1}-ABD-Z_{HER3_1} produced by B184, B184 *pep4Δ* and B184 *pep4Δprc1Δ*. **Table S1.** Primers used in this study. **Table S2.** Repair fragments for CRISPR used in this study.

Acknowledgements

The authors would like to thank Qi Qi and Yun Chen for help and support with the fed-batch fermentations.

Authors' contributions

VG, VS, FD, AS and ME conceived and designed the study. VG performed most of the experiments. MTM helped with the fermentation experiments. SE performed the BLI experiment, RG performed the binding kinetics assay, ST performed the mass spec experiment and data analysis, VG, VS and MM have written or revised the manuscript. All the authors have approved the submitted manuscript. All authors read and approved the final manuscript.

Funding

Open access funding provided by Chalmers University of Technology. The work was funded by the VINNOVA center CellNova (2017-02105) and a grant from the ÅForsk foundation (2019-649). The funding agencies did not partake in designing the study, collecting, analyzing or interpreting the data or writing the manuscript.

Availability of data and materials

Materials are available from the corresponding author upon reasonable request.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Biology and Biological Engineering, Chalmers University of Technology, Gothenburg, Sweden. ²Novo Nordisk Foundation Center for Biosustainability, Chalmers University of Technology, Gothenburg, Sweden. ³Affibody AB, Stockholm, Sweden.

Received: 22 October 2021 Accepted: 22 February 2022

Published online: 09 March 2022

References

- Moks T, Abramsén L, Nilsson B, Hellman U, Sjöquist J, Uhlén M. Staphylococcal protein A consists of five IgG-binding domains. *Eur J Biochem*. 1986;156:637–43.
- Nilsson B, Moks T, Jansson B, Abrahmsén L, Elmlblad A, Holmgren E, et al. A synthetic IgG-binding domain based on staphylococcal protein a. *Protein Eng Des Sel*. 1987;1:107–13.
- Nord K, Gunneriusson E, Ringdahl J, Stahl S, Uhlén M, Nygren A. Binding proteins selected from combinatorial libraries of an a-helical bacterial receptor domain. *Nat Biotechnol*. 1997;15:722–77.
- Rönmark J, Grönlund H, Uhlén M, Nygren PÅ. Human immunoglobulin A (IgA)-specific ligands from combinatorial engineering of protein A. *Eur J Biochem*. 2002;269(11):2647–55.
- Tove Boström JN and SH. Purification systems based on bacterial surface proteins. In: *Protein Purification*. Stockholm; 2012. p. 89–136.
- Mishra R, Patel H, Alanazi S, Yuan L, Garrett JT. HER3 signaling and targeted therapy in cancer. *Oncol Rev*. 2018;12:45–62.
- Frejd FY, Kim KT. Affibody molecules as engineered protein drugs. *Exp Mol Med*. 2017;49:e306.
- Orlova A, Magnusson M, Eriksson TLJ, Nilsson M, Larsson B, Höiden-Guthenberg I, et al. Tumor imaging using a picomolar affinity HER2 binding Affibody molecule. *Cancer Res*. 2006;66:4339–48.
- Sörensen J, Velikyan I, Sandberg D, Wennborg A, Feldwisch J, Tolmachev V, et al. Measuring HER2-receptor expression in metastatic breast cancer using [68Ga]ABY-025 Affibody PET/CT. *Theranostics*. 2016;6:262–71.
- Tolmachev V, Nilsson FY, Widström C, Andersson K, Rosik D, Gedda L, et al. 111In-Benzyl-DTPA-ZHER2:342, an affibody-based conjugate for in vivo imaging of HER2 expression in malignant tumors. *J Nucl Med*. 2006;47:846–53.
- Zielinski R, Lyakhov I, Hassan M, Kuban M, Shafer-Weaver K, Gandjbakhche A, et al. HER2-affitoxin: a potent therapeutic agent for the treatment of HER2-overexpressing tumors. *Clin Cancer Res*. 2011;17:5071–81.
- Huang CJ, Lin H, Yang X. Industrial production of recombinant therapeutics in *Escherichia coli* and its recent advancements. *J Ind Microbiol Biotechnol*. 2012;39:383–99.
- de Oliveira MP, Lopes AM, Mazzola PG, Rangel-Yagui C, Penna TCV, Pessoa A. Methods of endotoxin removal from biological preparations: a review. *J Pharm Pharm Sci*. 2007;10:388–404.
- Mamat U, Wilke K, Bramhill D, Schromm AB, Lindner B, Kohl TA, et al. Detoxifying *Escherichia coli* for endotoxin-free production of recombinant proteins. *Microb Cell Fact*. 2015;14:1–15. <https://doi.org/10.1186/s12934-015-0241-5>.
- Wang G, Huang M, Nielsen J. Exploring the potential of *Saccharomyces cerevisiae* for biopharmaceutical protein production. *Curr Opin Biotechnol*. 2017;48:77–84.
- Walsh G. Biopharmaceutical benchmarks 2014. *Nat Biotechnol*. 2014;32(10):992–1000.
- Dong D, Xia G, Li Z, Li Z. Human serum albumin and HER2-binding Affibody fusion proteins for targeted delivery of fatty acid-modified molecules and therapy. *Mol Pharm*. 2016;13:3370–80.
- Nielsen J. Production of biopharmaceutical proteins by yeast. *Bioengineered*. 2013;4:207–11.
- Huang M, Bai Y, Sjöström SL, Hallström BM, Liu Z, Petranovic D, et al. Microfluidic screening and whole-genome sequencing identifies mutations associated with improved protein secretion by yeast. *Proc Natl Acad Sci U S A*. 2015;112:E4689–96.
- Huang M, Bao J, Hallström BM, Petranovic D, Nielsen J. Efficient protein production by yeast requires global tuning of metabolism. *Nat Commun*. 2017;8:1131.
- Liu Z, Tyo KEJ, Martínez JL, Petranovic D, Nielsen J. Different expression systems for production of recombinant proteins in *Saccharomyces cerevisiae*. *Biotechnol Bioeng*. 2012;109:1259–68.
- Wittrup KD, Benig V. Optimization of amino acid supplements for heterologous protein secretion in *Saccharomyces cerevisiae*. *Biotechnol Tech*. 1994;8:161–6.

23. Wu M, Shen Q, Yang Y, Zhang S, Qu W, Chen J, et al. Disruption of *YPS1* and *PEP4* genes reduces proteolytic degradation of secreted HSA/PTH in *Pichia pastoris* GS115. *J Ind Microbiol Biotechnol*. 2013;40:589–99.
24. Parr CL, Keates RAB, Bryksa BC, Ogawa M, Yada RY. The structure and function of *Saccharomyces cerevisiae* proteinase A. *Yeast*. 2007;24:467–80.
25. Kang HA, Choi ES, Hong WK, Kim JY, Ko SM, Sohn JH, et al. Proteolytic stability of recombinant human serum albumin secreted in the yeast *Saccharomyces cerevisiae*. *Appl Microbiol Biotechnol*. 2000;53:575–82.
26. Hecht KA, O'Donnell AF, Brodsky JL. The proteolytic landscape of the yeast vacuole. *Cell Logist*. 2014;4:e28023.
27. Carmona-Gutiérrez D, Bauer MA, Ring J, Knaue HR, Eisenberg T, Büttner S, et al. The propeptide of yeast cathepsin D inhibits programmed necrosis. *Cell Death Dis*. 2011;2:1–10.
28. Dreyer T. Substrate specificity of proteinase yscA from *Saccharomyces cerevisiae*. *Carlsberg Res Commun*. 1989;54:85–97.
29. Kondo H, Shibano Y, Amachi T, Cronin N, Oda K, Dunn BM. Substrate specificities and kinetic properties of proteinase A from the yeast *Saccharomyces cerevisiae* and the development of a novel substrate. *J Biochem*. 1998;124:141–7.
30. Van Den Hazel HB, Kielland-Brandt MC, Winther JR. Review: Biosynthesis and function of yeast vacuolar proteases. *Yeast*. 1996;12:1–16.
31. Rupp S, Hirsch HH, Wolf DH. Biogenesis of the yeast vacuole (lysosome) Active site mutation in the vacuolar aspartate proteinase yscA blocks maturation of vacuolar proteinases. *FEBS Lett*. 1991;293:62–6.
32. van den Hazel HB, Kielland-Brandt MC, Winther JR. Autoactivation of proteinase A initiates activation of yeast vacuolar zymogens. *Eur J Biochem*. 1992;207:277–83.
33. Woolford CA, Daniels LB, Park FJ, Jones EW, Van Arsdel JN, Innis MA. The *PEP4* gene encodes an aspartyl protease implicated in the posttranslational regulation of *Saccharomyces cerevisiae* vacuolar hydrolases. *Mol Cell Biol*. 1986;6:2500–10.
34. Dong L, Li F, Piao Y, Sun D, Zhao R, Li C, et al. Characterization of proteinase A excretion from *Saccharomyces cerevisiae* in high sugar stress conditions. *J Korean Soc Appl Biol Chem*. 2015;58:203–8.
35. Nebes VL, Jones EW. Activation of the proteinase B precursor of the yeast *Saccharomyces cerevisiae* by autocatalysis and by an internal sequence. *J Biol Chem*. 1991;266:22851–7.
36. Mechler B, Müller H, Wolf DH. Maturation of vacuolar (lysosomal) enzymes in yeast: proteinase yscA and proteinase yscB are catalysts of the processing and activation event of carboxypeptidase yscY. *EMBO J*. 1987;6:2157–63.
37. Hemmings BA, Zubenko GS, Hasilik A, Jones EW. Mutant defective in processing of an enzyme located in the lysosome-like vacuole of *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*. 1981;78:435–9.
38. Komeda T, Sakai Y, Kato N, Kondo K. Construction of protease-deficient candida boidinii strains useful for recombinant protein production: cloning and disruption of proteinase A gene (*PEP4*) and proteinase B gene (*PRB1*). *Biosci Biotechnol Biochem*. 2002;66:628–31.
39. Safder I, Khan S, Islam I, Ali MK, Bibi ZWM. Protein expression system: a potential candidate to express protein in industrial and biopharmaceutical domains. *J Biomed Lett*. 2018;4:1–13.
40. Klionsky DJ, Banta LM, Emr SD. Intracellular sorting and processing of a yeast vacuolar hydrolase: proteinase A propeptide contains vacuolar targeting information. *Mol Cell Biol*. 1988;8:2105–16.
41. Ishchuk OP, Frost AT, Muñoz-Paredes F, Matsumoto S, Laforge N, Eriksson NL, et al. Improved production of human hemoglobin in yeast by engineering hemoglobin degradation. *Metab Eng*. 2021;66:259–67.
42. Wang ZY, He XP, Zhang BR. Over-expression of GSH1 gene and disruption of *PEP4* gene in self-cloning industrial brewer's yeast. *Int J Food Microbiol*. 2007;119:192–9.
43. Nord K, Nilsson J, Nilsson B, Uhlén M, Nygren PÅ. A combinatorial library of an α -helical bacterial receptor domain. *Protein Eng Des Sel*. 1995;8:601–8.
44. Rigoulet M, Aguilaniu H, Avéret N, Bunoust O, Camougrand N, Grandier-Vazeille X, et al. Organization and regulation of the cytosolic NADH metabolism in the yeast *Saccharomyces cerevisiae*. *Mol Cell Biochem*. 2004;256–257:73–81.
45. Gancedo C, Gancedo JM, Sols A. Glycerol metabolism in yeasts: pathways of utilization and production. *Eur J Biochem*. 1968;5:165–72.
46. Compagno C, Boschi F, Ranzi BM. Glycerol production in a triose phosphate isomerase deficient mutant of *Saccharomyces cerevisiae*. *Biotechnol Prog*. 1996;12:591–5.
47. Joly JC, Leung WS, Swartz JR. Overexpression of *Escherichia coli* oxidoreductases increases recombinant insulin-like growth factor-I accumulation. *Proc Natl Acad Sci U S A*. 1998;95:2773–7.
48. Qian ZG, Xia XX, Choi JH, Lee SY. Proteome-based identification of fusion partner for high-level extracellular production of recombinant proteins in *Escherichia coli*. *Biotechnol Bioeng*. 2008;101:587–601.
49. Yim S, Jeong K, Chang H, Lee S. High-level secretory production of human granulocyte-colony stimulating factor by fed-batch culture of recombinant *Escherichia coli*. *Bioprocess Biosyst Eng*. 2001;24:249–54.
50. Gueldener U, Heinisch J, Koehler GJ, Voss D, Hegemann JH. A second set of loxP marker cassettes for Cre-mediated multiple gene knockouts in budding yeast. *Nucleic Acids Res*. 2002;30:23.
51. Zhu Z, Zhou YJ, Kang MK, Krivoruchko A, Buijs NA, Nielsen J. Enabling the synthesis of medium chain alkanes and 1-alkenes in yeast. *Metab Eng*. 2017;44:81–8.
52. Inoue H, Nojima H, Okayama H. High efficiency transformation of *Escherichia coli* with plasmids. *Gene*. 1990;96:23–8.
53. Overkamp KM, Bakker BM, Kötter P, Luttkik MAH, Van Dijken JP, Pronk JT. Metabolic engineering of glycerol production in *Saccharomyces cerevisiae*. *Appl Environ Microbiol*. 2002;68:2814–21.
54. Compagno C, Brambilla L, Capitanio D, Boschi F, Ranzi BM, Porro D. Alterations of the glucose metabolism in a triose phosphate isomerase-negative *Saccharomyces cerevisiae* mutant. *Yeast*. 2001;18:663–70.
55. Gietz RD, Schiestl RH. High-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. *Nat Protoc*. 2007;2:31–4.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

