



**CHALMERS**  
UNIVERSITY OF TECHNOLOGY

## **Triplet Formation in a 9,10-Bis(phenylethynyl)anthracene Dimer and Trimer Occurs by Charge Recombination Rather than Singlet Fission**

Downloaded from: <https://research.chalmers.se>, 2026-04-05 21:29 UTC

Citation for the original published paper (version of record):

Ringström, R., Schroeder, Z., Mencaroni, L. et al (2023). Triplet Formation in a 9,10-Bis(phenylethynyl)anthracene Dimer and Trimer Occurs by Charge Recombination Rather than Singlet Fission. *Journal of Physical Chemistry Letters*, 14(35): 7897-7902. <http://dx.doi.org/10.1021/acs.jpcllett.3c02050>

N.B. When citing this work, cite the original published paper.

# Triplet Formation in a 9,10-Bis(phenylethynyl)anthracene Dimer and Trimer Occurs by Charge Recombination Rather than Singlet Fission

Rasmus Ringström, Zachary W. Schroeder, Letizia Mencaroni, Pavel Chabera, Rik R. Tykwinski, and Bo Albinsson\*



Cite This: *J. Phys. Chem. Lett.* 2023, 14, 7897–7902



Read Online


ACCESS |

Metrics & More

Article Recommendations

Supporting Information

**ABSTRACT:** We present an experimental study investigating the solvent-dependent dynamics of a 9,10-bis(phenylethynyl)anthracene monomer, dimer, and trimer. Using transient absorption spectroscopy, we have discovered that triplet excited state formation in the dimer and trimer molecules in polar solvents is a consequence of charge recombination subsequent to symmetry-breaking charge separation rather than singlet fission. Total internal reflection emission measurements of the monomer demonstrate that excimer formation serves as the primary decay pathway at a high concentration. In the case of highly concentrated solutions of the trimer, we observe evidence of triplet formation without the prior formation of a charge-separated state. We postulate that this is attributed to the formation of small aggregates, suggesting that oligomers mimicking the larger chromophore counts in crystals could potentially facilitate singlet fission. Our experimental study sheds light on the intricate dynamics of the 9,10-bis(phenylethynyl)anthracene system, elucidating the role of solvent- and concentration-dependent factors for triplet formation and charge separation.




Singlet fission (SF) is a multiple exciton generation process in which a singlet excited state shares its excitation energy with a neighboring ground state molecule to produce two triplet excited states.<sup>1</sup> SF has garnered significant attention in recent years due to its potential to enhance the efficiency of photovoltaic devices by utilizing the excess energy from high-energy photons that would otherwise be lost as heat.<sup>2,3</sup> To integrate SF with conventional photovoltaics, certain essential requirements must be met. These requirements include photostability to ensure the longevity of the material under prolonged light exposure as well as energy alignment of the resulting triplet states with the bandgap of the photovoltaic device to enable efficient charge separation and collection. Meeting these criteria is crucial for harnessing the full potential of SF in practical applications. One molecule that exhibits promising characteristics to fulfill these requirements is 9,10-bis(phenylethynyl)anthracene (BPEA). BPEA is a robust and photostable industrial dye with a triplet energy of 1.2–1.3 eV,<sup>4</sup> and it has demonstrated the ability to undergo SF both in thin films and in nanoparticles.<sup>5–8</sup> However, studies using molecular dimers mimicking the slip-stacked nature of solid-state packing have suffered from low yields of SF due to competing excimer formation.<sup>9</sup> SF in BPEA and other materials has been shown to be highly sensitive to the interchromophore orientation and distance.<sup>8,10–12</sup> Because calculations suggest that the basic energetic requirement  $E(S_1) \geq 2E(T_1)$ <sup>4,13</sup> for SF is fulfilled for BPEA, it is possible that there are dimer designs that can suppress the excimer decay pathway and promote SF. Furthermore, if the calculations are underestimating the triplet energy and SF is slightly

endothermic, it might be possible to enhance the rate of SF by increasing the number of chromophores as has been seen for other endothermic SF systems.<sup>14,15</sup>

Herein, we have synthesized and investigated a BPEA dimer (BPEAdim) and trimer (BPEAtri) connected in the meta position of the central phenylene unit. The molecular structures are shown in [Scheme 1](#) together with the molecular


**Scheme 1. Molecular Structures of BPEAmono, BPEAdim, and BPEAtri**



**Received:** July 24, 2023

**Accepted:** August 25, 2023

**Published:** August 29, 2023




structure of the monomer (BPEAmono). Using fluorescence spectroscopy measurements under total internal reflection (TIR) conditions on highly concentrated solutions of BPEAmono, we found that the main decay pathway of the monomer is by excimer formation. In contrast, BPEAdim and BPEAtri show no evidence of excimer formation, and the formation of the triplet excited state is observed. The triplet excited state formation is, however, not a consequence of SF, but rather by charge recombination from the symmetry breaking charge separated state (CSS) in polar solvents.

The syntheses of BPEAdim and BPEAtri share 9,10-dibromoanthracene as a common precursor (Scheme 1; see the Supporting Information for details). Substitution at the central benzene or peripheral phenyl moieties of BPEAdim and BPEAtri, respectively, was necessary to provide solubility and facilitate purification. The *m*-phenylene-linked BPEAdim was synthesized based on a two-step procedure from known building blocks. The precursor to BPEAdim, acetoxy-protected dimer BPEA-Ac (Scheme 1), was difficult to purify, and the crude reaction mixture was thus carried through the subsequent deacetylation to enhance the polarity of the dimer and facilitate purification. Gratifyingly, BPEAdim was obtained pure through a sequence of precipitation from the hydrolysis reaction mixture, column chromatography on silica gel, and recrystallization from CH<sub>2</sub>Cl<sub>2</sub>/MeOH. Compound BPEAtri was prepared using a 3-fold Sonogashira coupling reaction with the appropriate brominated anthracene and 1,3,5-triethynylbenzene building blocks. The product was purified via a sequence of size exclusion chromatography, silica gel column chromatography, and precipitation from a saturated solution of toluene via the addition of hexanes at -30 °C. Both BPEAdim and BPEAtri are stable solids under ambient conditions. Refer to Section S8 in the Supporting Information for further details.

Figure 1 displays the absorption and emission spectra of dilute solutions (~5–10 μM) in tetrahydrofuran (THF). The absorption and emission spectra of BPEAmono, BPEAdim, and BPEAtri show only minor variations. The S<sub>1</sub> energy of BPEAdim, as determined from the intersection of the absorption and emission signals, is red-shifted 0.027 eV relative to the monomer at 474 nm, and BPEAtri is shifted by an additional 0.027 eV. The unstructured absorption spectrum, in contrast to the structured emission, is attributed to excited state planarization, which has previously been studied in detail.<sup>16</sup>


The photophysical properties of BPEAmono have been initially examined in both dilute and concentrated solutions to assess its suitability for SF. In dilute solution, BPEAmono has a fluorescence quantum yield close to unity and decays monoexponentially with a lifetime of approximately 3.4 ns (Figure 2a). Conversely, highly concentrated solutions show significantly quenched emission, as evidenced by the data presented in Figure 2a. To gain further insight into the behavior of BPEAmono in concentrated solutions, emission measurements were conducted under total internal reflection (TIR) conditions. These measurements, in conjunction with transient absorption analysis, indicate that excimer formation rather than SF serves as the primary decay pathway in the concentrated solutions. Section S2 in the Supporting Information provides a comprehensive account of the analysis, and Section S3 outlines the TIR technique employed. The formation of excimers is a plausible explanation for the poor



**Figure 1.** Steady state absorption (solid line) and emission (dotted line) of BPEAmono, BPEAdim, and BPEAtri in tetrahydrofuran (THF).


upconversion quantum yield previously reported for BPEAmono.<sup>4</sup>

The emission spectra of BPEAdim and BPEAtri closely resemble that of BPEAmono in a dilute solution (Figure 1). Notably, there is no evidence of excimer formation, as observed in other BPEA dimers<sup>9</sup> as well as for BPEAmono at high concentration in the present study. Additionally, the emission lifetime of BPEAdim and BPEAtri in nonpolar solvents is similar to that of BPEAmono, indicating no substantial quenching of the singlet excited state. In polar solvents, however, emission of BPEAdim decays significantly within the response function of our time-correlated single photon counting (TCSPC) instrument (Figure 2b). To gain a better understanding of the quenching mechanism, we employed femtosecond transient absorption (fsTA). Figure 3 presents the fsTA spectra and the species associated spectra of BPEAdim in THF, showcasing a spectral evolution comparable to that of BPEAmono (Figure S4). Initially, the S<sub>1</sub> absorption, centered at 560 nm, evolves within the first few picoseconds, corresponding to excited state planarization of the molecule. After planarization, the spectral features uniformly decay across the entire spectral range with a lifetime of 3.2 ns, corroborating well the results from TCSPC (Figure 2b and Table S3). The fsTA spectra of BPEAdim in THF have been analyzed using a two-component sequential decay model shown in the inset of Figure 3b. By employing singular value decomposition (SVD) and global analysis with the KiMoPack software,<sup>17</sup> we can extract the spectral components and kinetics, as illustrated in Figures 3b and 3c. When BPEAdim is studied in the more polar solvent acetonitrile (ACN), we observe similar planarization as in THF, which is documented by the initial growth of the signal at 560 nm (Figure 4a). Following planarization, a new species emerges concomitantly with the decay of the singlet excited state, giving rise to broad excited state absorption at 640 nm. Based on the similarity with the spectra of the radical anion obtained through spectroelec-




**Figure 2.** Time-resolved emission obtained with an excitation wavelength of 405 nm. (a) BPEAmono in THF for different concentrations ( $C$ ) monitored at 473 nm. (b) BPEAdim in acetonitrile (ACN) and THF monitored at 480 nm. Fits for lifetimes are presented in Tables S1 and S3 for BPEAmono and BPEAdim, respectively.

trochemistry (Figure 4f and additional details in Section S4) and previous reports on BPEA dimers, we assign this new species as the CSS. Note that the spectroscopic signature of the CSS is a combination of the spectra of the radical anion and cation. For BPEAdim, however, the spectrum of the radical cation has very low molar absorptivity in the visible region compared to the anion (Figure S12). As the CSS state decays, a fourth state becomes discernible at 490 nm. This state exhibits a strong resemblance to the sensitized triplet excited state and has not been reported in previous studies on BPEA dimers.<sup>18</sup> It is possible that this triplet has eluded detection due to the use of less polar solvents or, alternatively, to limitations in the length of delay stages, as the signal becomes apparent only at longer time delays. The fsTA spectra of BPEAdim in Figure 4a have been fit to the four-component model (Figure 4d). The spectral components and kinetics at selected wavelengths are displayed in Figures 4b and 4c, respectively. Previous studies on SF dimers with significant through-bond electronic coupling typically exhibit short triplet-pair lifetimes, ranging from several to hundreds of picoseconds, owing to the lack of triplet decorrelation and subsequent separation into free triplets.<sup>19–23</sup> In contrast, the microsecond lifetime of the triplet excited state observed in our analysis (measured with nanosecond transient absorption, Figures S18 and S19), coupled with the emergence of the signal after the



**Figure 3.** (a) fsTA analyses of BPEAdim in a dilute solution of THF ( $\sim 10 \mu\text{M}$ ); excitation at 420 nm. (b) Species associated spectra of the singular value decomposition (SVD) analysis using the kinetic model shown in the inset. (c) Selected kinetics of ground-state bleach and stimulated emission at 483 nm and  $S_1$  excited state absorption at 580 nm with the model data shown as a solid line.

decay of the CSS, suggests the formation of one independent triplet excited state per molecule through charge recombination via CSS. Our findings thus support the hypothesis that the triplet formation is not a result of SF.<sup>24</sup> Triplet formation within organic molecules via a CSS commonly ensues through one of two distinct mechanisms: spin–orbit charge transfer intersystem crossing (SOCT-ISC) and radical-pair intersystem crossing (RP-ISC).<sup>25–27</sup> SOCT-ISC entails direct conversion of the initial  $^1\text{CSS}$  into the  $T_1$  state, involving a reverse electron transfer and a spin inversion process. This phenomenon is



**Figure 4.** (a) fsTA analyses of a dilute solution ( $\sim 5 \mu\text{M}$ ) of BPEAdim in ACN; excitation at 420 nm. (b) Species associated spectra of the SVD analysis using the model shown in panel d. (c) Selected kinetics of ground state bleach/ $T_1$  at 483 and 490 nm,  $S_1$  at 560 nm, and the CSS at 650 nm with the model data shown as a solid line. (d) Kinetic model and associated rate constants. The energy of the CSS was estimated based on the redox potentials of BPEAmo (see Section S4 for more details). (e) Comparison of the final spectral component from the singular value decomposition (SVD) analysis and kinetic model compared to the sensitized triplet absorption spectrum (see Section S7 for more details). (f) Comparison of the CSS spectral component with the radical anion obtained via spectroelectrochemical analysis (see Section S4 for more details). Note that the CSS spectrum is a combination of the radical anion and cation spectra, while for BPEAdim the spectrum of the radical cation has very low and broad molar absorptivity in the visible region compared with the anion (Figure S12).

notably amplified when the constituent subunits assume a nearly perpendicular alignment. Such an orientation facilitates compensation for changes in electron spin angular momentum during ISC through adjustments in molecular orbital angular momentum. This intricate alignment is particularly evident in closely spaced dyads, where steric hindrance between subunits enforces an orthogonal arrangement, as observed in dyads linked by a single C–C bond.<sup>28,29</sup> In contrast, the RP-ISC mechanism involves the emergence of an intermediate triplet charge-transfer state ( $^3\text{CSS}$ ) prior to charge recombination, leading to either the triplet state or the ground state. This mechanism is more prevalent in electron donor–acceptor dyads characterized by weak electronic coupling between their donor and acceptor subunits due to considerable spatial separation (e.g.,  $>10$ – $15 \text{ \AA}$ ). This separation results in a minimal energy difference between  $^1\text{CSS}$  and  $^3\text{CSS}$ . Notably, certain studies posit the coexistence of both mechanisms.<sup>30</sup> Distinguishing between SOCT-ISC and RP-ISC through fsTA spectroscopy proves challenging because radical pairs are central to both processes, yielding closely resembling optical spectral signatures. Consequently, we can only engage in qualitative speculation regarding the mechanism underlying our observations. The rate of triplet excited state formation seems to align with previous RP-ISC studies, as spin–orbit coupled mechanisms generally exhibit larger rates.<sup>31</sup> Additionally, the interchromophore distance within BPEAdim exceeds  $10 \text{ \AA}$ , further reinforcing the alignment with the RP-ISC mechanism. The relatively slow charge recombination rate toward the ground state, in comparison to triplet formation, may be explicable through the substantial negative Gibbs free

energy shift accompanying ground state recombination. This scenario potentially gives rise to behavior consistent with the Marcus inverted region characteristics.

It is noteworthy that the CSS observed in our study has a spectroscopic signature that is similar to what was previously attributed to a shifted and broadened  $S_1$  absorption in BPEA thin films.<sup>5</sup> Based on this observation, it is possible that the triplet formation observed for BPEA thin films could originate from a similar charge recombination process as we have proposed in the present study and not from SF, as previously proposed.<sup>5</sup> The mechanism of charge recombination to form the triplet excited state has been proposed in other thin film systems, such as in the indigo derivative cibalackrot, in which case transient intermediates with charge-transfer character are also observed prior to triplet formation.<sup>32</sup> It is also worth noting that the interpretation of the spectral feature as a shifted and broadened  $S_1$  absorption in BPEA thin films could be entirely accurate, given that the excited state absorption of various compounds, including polyacenes, are known to differ significantly between solution and film states.<sup>33–37</sup>

The photophysical behavior of BPEAtri appears to be very similar to that of BPEAdim, and SF is not detected in dilute solution, regardless of solvent polarity. (Further details are provided in Section S6. BPEAtri was not investigated in ACN due to limited solubility.) Consequently, it seems that an additional BPEA moiety is insufficient to drive the seemingly slightly endothermic SF of BPEA. It is possible that a larger oligomer or a linear arrangement of chromophores could enhance the entropic driving force(s), as observed in other endothermic SF systems.<sup>14,15</sup> In fact, fsTA measurements in

highly concentrated solutions (25 mM) of BPEAtri in THF reveal a broad excited state absorption feature at 500 nm during later time delays that matches the sensitized triplet spectra (Figure S27). Although the absorption spectrum shows no evidence of significant scattering from aggregates, we posit that smaller aggregates may form, and these aggregates could facilitate SF. However, there is also evidence of excimer formation for BPEAtri as the emission lifetime depends on the probing wavelength, indicating the presence of at least two emissive species, similar to that observed for BPEAmono (Figure S23 and Table S8).

In this study, we investigated the photophysical properties of a BPEA monomer, dimer, and trimer. Our findings show that in highly concentrated solutions, BPEAmono exhibits excimer formation, and SF is not observed. The photophysics of BPEAdim and BPEAtri are primarily governed by symmetry-breaking charge separation, which outcompetes SF. Triplet excited states are, however, generated by charge recombination, where the CSS acts as an intermediate. Interestingly, there are indications of SF in highly concentrated solutions of the trimer as triplet excited state absorption is discernible without prior formation of the charge-separated state. This suggests that increasing the number of chromophores with larger oligomers, mimicking the larger chromophore count in crystals, may enhance the rate of SF for BPEA and potentially for other SF systems in solution.

## ■ ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.jpcllett.3c02050>.

Additional absorption, emission, transient absorption, spectroelectrochemical data, synthetic details, and full analysis of the compounds (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

**Bo Albinsson** – Department of Chemistry and Chemical Engineering, Chalmers University of Technology, 412 96 Gothenburg, Sweden; [orcid.org/0000-0002-5991-7863](https://orcid.org/0000-0002-5991-7863); Email: [balb@chalmers.se](mailto:balb@chalmers.se)

### Authors

**Rasmus Ringström** – Department of Chemistry and Chemical Engineering, Chalmers University of Technology, 412 96 Gothenburg, Sweden; [orcid.org/0000-0002-3340-957X](https://orcid.org/0000-0002-3340-957X)

**Zachary W. Schroeder** – Department of Chemistry, University of Alberta, Edmonton, Alberta, Canada T6G 2G2; [orcid.org/0000-0003-0590-9371](https://orcid.org/0000-0003-0590-9371)

**Letizia Mencaroni** – Department of Chemistry Biology and Biotechnology, University of Perugia, 06123 Perugia, Italy

**Pavel Chabera** – The Division of Chemical Physics and NanoLund, Lund University, 22100 Lund, Sweden; [orcid.org/0000-0002-0531-5138](https://orcid.org/0000-0002-0531-5138)

**Rik R. Tykwinski** – Department of Chemistry, University of Alberta, Edmonton, Alberta, Canada T6G 2G2; [orcid.org/0000-0002-7645-4784](https://orcid.org/0000-0002-7645-4784)

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.jpcllett.3c02050>

## Author Contributions

R.R. and B.A. designed the project. Z.W.S. synthesized and analyzed the compounds. R.R. performed most of the photophysical characterization. L.M. performed TIR emission measurements and characterization of BPEAmono. P.C. assisted with parts of the fsTA. The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

## Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

B.A. acknowledges support from the Swedish Research Council and the Swedish Energy Agency (Contract 46526-1). R.R.T. acknowledges support from NSERC (Natural Sciences and Engineering Research Council of Canada). Z.W.S. acknowledges graduate scholarship support from NSERC and Alberta Innovates.

## ■ REFERENCES

- (1) Smith, M. B.; Michl, J. Singlet Fission. *Chem. Rev.* **2010**, *110* (11), 6891–6936.
- (2) Hanna, M. C.; Nozik, A. J. Solar conversion efficiency of photovoltaic and photoelectrolysis cells with carrier multiplication absorbers. *J. Appl. Phys.* **2006**, *100* (7), 074510.
- (3) Rao, A.; Friend, R. H. Harnessing singlet exciton fission to break the Shockley-Queisser limit. *Nat. Rev. Mater.* **2017**, *2* (11), 17063.
- (4) Gray, V.; Dreos, A.; Erhart, P.; Albinsson, B.; Moth-Poulsen, K.; Abrahamsson, M. Loss channels in triplet-triplet annihilation photon upconversion: importance of annihilator singlet and triplet surface shapes. *Phys. Chem. Chem. Phys.* **2017**, *19* (17), 10931–10939.
- (5) Bae, Y. J.; Kang, G.; Malliakas, C. D.; Nelson, J. N.; Zhou, J.; Young, R. M.; Wu, Y.-L.; Van Duyne, R. P.; Schatz, G. C.; Wasielewski, M. R. Singlet Fission in 9,10-Bis(phenylethynyl) anthracene Thin Films. *J. Am. Chem. Soc.* **2018**, *140* (45), 15140–15144.
- (6) Manna, B.; Nandi, A.; Ghosh, R. Ultrafast Singlet Exciton Fission Dynamics in 9,10-Bis(phenylethynyl) anthracene Nanoaggregates and Thin Films. *J. Phys. Chem. C* **2018**, *122* (36), 21047–21055.
- (7) Fan, S.; Li, W.; Li, T.; Gao, F.; Hu, W.; Liu, S.; Wang, X.; Liu, H.; Liu, Z.; Li, Z.; Chen, Y.; Li, X. Singlet fission in colloidal nanoparticles of amphiphatic 9,10-bis(phenylethynyl) anthracene derivatives. *J. Photochem. Photobiol., A* **2022**, *427*, 113826.
- (8) Bae, Y. J.; Christensen, J. A.; Kang, G.; Zhou, J.; Young, R. M.; Wu, Y.-L.; Van Duyne, R. P.; Schatz, G. C.; Wasielewski, M. R. Substituent effects on energetics and crystal morphology modulate singlet fission in 9,10-bis(phenylethynyl) anthracenes. *J. Chem. Phys.* **2019**, *151* (4), 044501.
- (9) Bae, Y. J.; Shimizu, D.; Schultz, J. D.; Kang, G.; Zhou, J.; Schatz, G. C.; Osuka, A.; Wasielewski, M. R. Balancing Charge Transfer and Frenkel Exciton Coupling Leads to Excimer Formation in Molecular Dimers: Implications for Singlet Fission. *J. Phys. Chem. A* **2020**, *124* (41), 8478–8487.
- (10) Papadopoulos, I.; Zirzmeier, J.; Hetzer, C.; Bae, Y. J.; Krzyaniak, M. D.; Wasielewski, M. R.; Clark, T.; Tykwinski, R. R.; Guldi, D. M. Varying the inter-pentacene electronic coupling to tune singlet fission. *J. Am. Chem. Soc.* **2019**, *141*, 6191–6203.
- (11) Zirzmeier, J.; Lehnher, D.; Coto, P. B.; Chernick, E. T.; Casillas, R.; Basel, B. S.; Thoss, M.; Tykwinski, R. R.; Guldi, D. M. Singlet fission in pentacene dimers. *Proc. Natl. Acad. Sci. U. S. A.* **2015**, *112* (17), 5325–5330.
- (12) Millington, O.; Montanaro, S.; Leventis, A.; Sharma, A.; Dowland, S. A.; Sawhney, N.; Fallon, K. J.; Zeng, W.; Congrave, D. G.; Musser, A. J.; Rao, A.; Bronstein, H. Soluble Diphenylhexatriene Dimers for Intramolecular Singlet Fission with High Triplet Energy. *J. Am. Chem. Soc.* **2023**, *145* (4), 2499–2510.

- (13) Zhong, F.; Zhao, J. Phenyleneanthracene derivatives as triplet energy acceptor/emitter in red light excitable triplet-triplet-annihilation upconversion. *Dyes Pigm.* **2017**, *136*, 909–918.
- (14) Korovina, N. V.; Chang, C. H.; Johnson, J. C. Spatial separation of triplet excitons drives endothermic singlet fission. *Nat. Chem.* **2020**, *12* (4), 391–398.
- (15) Liu, H.; Wang, Z.; Wang, X.; Shen, L.; Zhang, C.; Xiao, M.; Li, X. Singlet exciton fission in a linear tetracene tetramer. *J. Mater. Chem. C* **2018**, *6* (13), 3245–3253.
- (16) Fureraj, I.; Budkina, D. S.; Vauthey, E. Torsional disorder and planarization dynamics: 9,10-bis(phenylethynyl) anthracene as a case study. *Phys. Chem. Chem. Phys.* **2022**, *24* (42), 25979–25989.
- (17) Müller, C.; Pascher, T.; Eriksson, A.; Chabera, P.; Uhlig, J. KiMoPack: A python Package for Kinetic Modeling of the Chemical Mechanism. *J. Phys. Chem. A* **2022**, *126* (25), 4087–4099.
- (18) Su, P.; Liu, H.; Shen, L.; Zhou, J.; Wang, W.; Liu, S.; Nie, X.; Li, Z.; Liu, Z.; Chen, Y.; Li, X. Linker dependent symmetry breaking charge separation in 9,10-bis(phenylethynyl) anthracene dimers. *Materials Chemistry Frontiers* **2022**, *6* (6), 707–717.
- (19) Korovina, N. V.; Pompetti, N. F.; Johnson, J. C. Lessons from intramolecular singlet fission with covalently bound chromophores. *J. Chem. Phys.* **2020**, *152* (4), 040904.
- (20) Krishnapriya, K. C.; Roy, P.; Puttaraju, B.; Salzner, U.; Musser, A. J.; Jain, M.; Dasgupta, J.; Patil, S. Spin density encodes intramolecular singlet exciton fission in pentacene dimers. *Nat. Commun.* **2019**, *10* (1), 33.
- (21) Sanders, S. N.; Pun, A. B.; Parenti, K. R.; Kumarasamy, E.; Yablon, L. M.; Sfeir, M. Y.; Campos, L. M. Understanding the Bound Triplet-Pair State in Singlet Fission. *Chem.* **2019**, *5* (8), 1988–2005.
- (22) Lukman, S.; Musser, A. J.; Chen, K.; Athanasopoulos, S.; Yong, C. K.; Zeng, Z.; Ye, Q.; Chi, C.; Hodgkiss, J. M.; Wu, J.; Friend, R. H.; Greenham, N. C. Tuneable Singlet Exciton Fission and Triplet-Triplet Annihilation in an Orthogonal Pentacene Dimer. *Adv. Funct. Mater.* **2015**, *25* (34), 5452–5461.
- (23) Korovina, N. V.; Das, S.; Nett, Z.; Feng, X.; Joy, J.; Haiges, R.; Krylov, A. I.; Bradforth, S. E.; Thompson, M. E. Singlet Fission in a Covalently Linked Cofacial Alkynyltetracene Dimer. *J. Am. Chem. Soc.* **2016**, *138* (2), 617–627.
- (24) Electron paramagnetic resonance (EPR) spectroscopy could provide further insights into the nature of the formed triplet excited state, but unfortunately, we lack access to this method in our laboratory.
- (25) Hou, Y.; Zhang, X.; Chen, K.; Liu, D.; Wang, Z.; Liu, Q.; Zhao, J.; Barbon, A. Charge separation, charge recombination, long-lived charge transfer state formation and intersystem crossing in organic electron donor/acceptor dyads. *J. Mater. Chem. C* **2019**, *7* (39), 12048–12074.
- (26) Gibbons, D. J.; Farawar, A.; Mazzella, P.; Leroy-Lhez, S.; Williams, R. M. Making triplets from photo-generated charges: observations, mechanisms and theory. *Photochemical & Photobiological Sciences* **2020**, *19* (2), 136–158.
- (27) Nguyen, V.-N.; Yan, Y.; Zhao, J.; Yoon, J. Heavy-Atom-Free Photosensitizers: From Molecular Design to Applications in the Photodynamic Therapy of Cancer. *Acc. Chem. Res.* **2021**, *54* (1), 207–220.
- (28) Filatov, M. A.; Karuthedath, S.; Polestshuk, P. M.; Savoie, H.; Flanagan, K. J.; Sy, C.; Sitte, E.; Telitchko, M.; Laquai, F.; Boyle, R. W.; Senge, M. O. Generation of Triplet Excited States via Photoinduced Electron Transfer in meso-anthra-BODIPY: Fluorogenic Response toward Singlet Oxygen in Solution and in Vitro. *J. Am. Chem. Soc.* **2017**, *139* (18), 6282–6285.
- (29) Wang, Z.; Zhao, J. Bodipy-Anthracene Dyads as Triplet Photosensitizers: Effect of Chromophore Orientation on Triplet-State Formation Efficiency and Application in Triplet-Triplet Annihilation Upconversion. *Org. Lett.* **2017**, *19* (17), 4492–4495.
- (30) Colvin, M. T.; Ricks, A. B.; Scott, A. M.; Co, D. T.; Wasielewski, M. R. Intersystem Crossing Involving Strongly Spin Exchange-Coupled Radical Ion Pairs in Donor-bridge-Acceptor Molecules. *J. Phys. Chem. A* **2012**, *116* (8), 1923–1930.
- (31) Filatov, M. A. Heavy-atom-free BODIPY photosensitizers with intersystem crossing mediated by intramolecular photoinduced electron transfer. *Organic & Biomolecular Chemistry* **2020**, *18* (1), 10–27.
- (32) Ryerson, J. L.; Zaykov, A.; Aguilar Suarez, L. E.; Havenith, R. W. A.; Stepp, B. R.; Dron, P. I.; Kaleta, J.; Akdag, A.; Teat, S. J.; Magnera, T. F.; Miller, J. R.; Havlas, Z.; Broer, R.; Faraji, S.; Michl, J.; Johnson, J. C. Structure and photophysics of indigoids for singlet fission: Cibalackrot. *J. Chem. Phys.* **2019**, *151* (18), 184903.
- (33) Dvořák, M.; Prasad, S. K. K.; Dover, C. B.; Forest, C. R.; Kaleem, A.; Macqueen, R. W.; Petty, A. J.; Forecast, R.; Beves, J. E.; Anthony, J. E.; Tayebjee, M. J. Y.; Widmer-Cooper, A.; Thordarson, P.; Schmidt, T. W. Singlet Fission in Concentrated TIPS-Pentacene Solutions: The Role of Excimers and Aggregates. *J. Am. Chem. Soc.* **2021**, *143* (34), 13749–13758.
- (34) Pensack, R. D.; Grieco, C.; Purdum, G. E.; Mazza, S. M.; Tilley, A. J.; Ostroumov, E. E.; Seferos, D. S.; Loo, Y.-L.; Asbury, J. B.; Anthony, J. E.; Scholes, G. D. Solution-processable, crystalline material for quantitative singlet fission. *Materials Horizons* **2017**, *4* (5), 915–923.
- (35) Tayebjee, M. J. Y.; Schwarz, K. N.; Macqueen, R. W.; Dvořák, M.; Lam, A. W. C.; Ghiggino, K. P.; McCamey, D. R.; Schmidt, T. W.; Conibeer, G. J. Morphological Evolution and Singlet Fission in Aqueous Suspensions of TIPS-Pentacene Nanoparticles. *J. Phys. Chem. C* **2016**, *120* (1), 157–165.
- (36) Pensack, R. D.; Ostroumov, E. E.; Tilley, A. J.; Mazza, S.; Grieco, C.; Thorley, K. J.; Asbury, J. B.; Seferos, D. S.; Anthony, J. E.; Scholes, G. D. Observation of Two Triplet-Pair Intermediates in Singlet Exciton Fission. *J. Phys. Chem. Lett.* **2016**, *7* (13), 2370–2375.
- (37) Nandi, A.; Manna, B.; Ghosh, R. Ultrafast Dynamics of the 9,10-Bis[(triisopropylsilyl) ethynyl]anthracene Nanoaggregate and Thin Film: Influence of the Molecular Packing and Sample Morphology on Singlet Fission and Excimer Formation. *J. Phys. Chem. C* **2023**, *127* (26), 12621–12630.

## Recommended by ACS

### Efficient Intramolecular Singlet Fission in Spiro-Linked Heterodimers

Oskar Kefer, Tiago Buckup, *et al.*

AUGUST 03, 2023  
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

### Ultrafast Dynamics of the 9,10-Bis[(triisopropylsilyl)ethynyl]anthracene Nanoaggregate and Thin Film: Influence of the Molecular Packing and Samp...

Amitabha Nandi, Rajib Ghosh, *et al.*

JUNE 26, 2023  
THE JOURNAL OF PHYSICAL CHEMISTRY C

READ 

### Direct Observation of Ultrafast Relaxation Dynamics of a Mixed Excimer State in Perylene Monoimide Dimer by Femtosecond Transient Absorption

Chaochao Qin, Yuhai Jiang, *et al.*

MARCH 03, 2023  
THE JOURNAL OF PHYSICAL CHEMISTRY LETTERS

READ 

### Triplets with a Twist: Ultrafast Intersystem Crossing in a Series of Electron Acceptor Materials Driven by Conformational Disorder

Sai Shruthi Murali, Paul A. Hume, *et al.*

DECEMBER 20, 2022  
JOURNAL OF THE AMERICAN CHEMICAL SOCIETY

READ 

Get More Suggestions >