



CHALMERS
UNIVERSITY OF TECHNOLOGY

Proton halo effects in the $8\text{B}+64\text{Zn}$ collision around the Coulomb barrier

Downloaded from: <https://research.chalmers.se>, 2026-04-03 07:16 UTC

Citation for the original published paper (version of record):

Di Pietro, A., Spartà, R., Figuera, P. et al (2023). Proton halo effects in the $8\text{B}+64\text{Zn}$ collision around the Coulomb barrier. *Journal of Physics: Conference Series*, 2586(1).
<http://dx.doi.org/10.1088/1742-6596/2586/1/012079>

N.B. When citing this work, cite the original published paper.

PAPER • OPEN ACCESS

Proton halo effects in the $^8\text{B}+^{64}\text{Zn}$ collision around the Coulomb barrier

To cite this article: A Di Pietro *et al* 2023 *J. Phys.: Conf. Ser.* **2586** 012079

View the [article online](#) for updates and enhancements.

You may also like

- [Effective field theory description of halo nuclei](#)
H-W Hammer, C Ji and D R Phillips
- [Physics with post-accelerated beams at ISOLDE: nuclear reactions](#)
A Di Pietro, K Riisager and P Van Duppen
- [Final results of an experiment to search for 2 processes in zinc and tungsten with the help of radiopure ZnWO₄ crystal scintillators](#)
P Belli, R Bernabei, F Cappella et al.

Proton halo effects in the ${}^8\text{B}+{}^{64}\text{Zn}$ collision around the Coulomb barrier

A Di Pietro¹, R Spartá^{1,2}, P Figuera¹, O Tengblad³, A Moro⁴, J Lei⁵, JP Fernandez Garcia⁴, L Acosta^{1,6}, B Jonson⁷; MJ Borge³, J Cederkäll⁸, T Davinson⁹, JD Ovejas³, L Fraile¹⁰, D Galaviz¹¹, J Halkjaer Jensen¹², M La Cognata¹, I Martel¹³, A Perea³, A Sanchez Benitez¹³, N Soić¹⁴ and S Vignales³

¹INFN, Laboratori Nazionali del Sud, via S. Sofia 62, I-95123 Catania, Italy

²Dipartimento di Fisica e Astronomia, via S. Sofia 64, I-95123 Catania, Italy

³Instituto de Estructura de la Materia, CSIC, Serrano 113 bis, E-28006 Madrid, Spain

⁴Departamento de FAMN, Universidad de Sevilla, Apartado 1065, E-41080 Seville, Spain

⁵INFN-Sezione di Pisa, Largo Bruno Pontecorvo 3, 56127 Pisa, Italy

⁶Instituto de Física, Universidad Nacional Autónoma de México, A.P. 20-364, México City 01000, México

⁶Department of Physics, Chalmers University of Technology, S-41296 Göteborg, Sweden

⁷Department of Physics, Chalmers University of Technology, S-41296 Göteborg, Sweden

⁸Physics Department, Lund University, Box 118, SE-221 00, Lund, Sweden

⁹School of Physics and Astronomy, University of Edinburgh, JCMB, Mayfield Road, Edinburgh EH9 3JZ, UK

¹⁰UCM Madrid, Plaza de Ciencias, 1 Ciudad Universitaria, 28040 - Madrid, Spain

¹¹LIP - Laboratory for Instrumentation and Experimental Particle Physics, Lisbon, Portugal

¹²Department of Physics and Astronomy, University of Aarhus, DK-8000 Aarhus, Denmark

¹³Departamento de Física Aplicada, Universidad de Huelva, Campus de El Carmen, E-21071 Huelva, Spain

¹⁴Rudjer Bošković Institute, Bijenička, 54 HR-10000 Zagreb, Croatia

E-mail: dipietro@lns.infn.it

Abstract. The ${}^8\text{B}+{}^{64}\text{Zn}$ reaction at 38.5 MeV has been studied at HIE-ISOLDE CERN to investigate proton halo effect on the reaction dynamics. For the first time it was used the only existing post-accelerated ${}^8\text{B}$ beam. The measured elastic scattering angular distribution showed a small suppression of the Coulomb-nuclear interference peak, opposite to what observed for the one-neutron halo nucleus ${}^{11}\text{Be}$ on the same target where a large suppression was observed instead. Inclusive angular and energy distributions of breakup fragments were also measured showing that, both, elastic and non-elastic breakup contribute. The presence of the additional Coulomb interactions halo-core and halo-target in ${}^8\text{B}$ makes the reaction dynamics in this proton-halo nucleus different than the neutron-halo case.

1. Introduction

${}^8\text{B}$ has a very low breakup threshold of 0.138 MeV and thus is a good candidate for having a proton-halo structure. Huge efforts have been made in the past years to understand the reaction dynamics around the Coulomb barrier with neutron-halo nuclei; on the other hand the dynamics at the barrier with proton-halo is expected to be different, but very few experimental data exist



(for a review see [1]). Many of the investigations performed with ^8B beams concern Coulomb dissociation at energies well above the Coulomb barrier in order to get indirect information on the radioactive capture reaction $^7\text{Be}(p,\alpha)$, but not many studies exist to investigate the reaction dynamics of proton-halo nuclei. All experiments have been done using in-flight ^8B beam, therefore with some limitations associated with the characteristics of such beams (purity, energy spread and beam emittance) compared to the post-accelerated ones. In the Coulomb dissociation process, proton-halo nuclei behave differently from neutron-halo, since the loosely bound proton participates in the reaction process. Namely, for a dynamic polarization effect the valence proton is displaced behind the nuclear core and shielded from the target; this effect causes a reduction of break-up probability compared to first-order perturbation theory predictions and higher-order corrections are required [e.g. [2, 3]]. As experimentally observed [4], nuclear processes are expected to have a primary role in the dissociation of ^8B [5]. Conversely for the neutron-halo nucleus ^{11}Be on the same target at a similar energy the Coulomb contribution far exceeds the nuclear one [6]. Unlike neutron-halo, in reactions with proton-halo nuclei there is an additional Coulomb interaction between the p and the core and the p and the target whose effect is to create an effective barrier which makes the proton of the halo effectively more bound [6].

In the case of reactions induced by neutron-halo nuclei, coupling to the continuum result in a suppression in the Coulomb-nuclear interference region in the elastic scattering angular distribution relative to the Rutherford [e.g. [7, 8, 9, 10, 11]], and a large total reaction cross-section is, as a consequence, observed. Moreover, a large fraction of total reaction cross-section is due to direct processes such as break-up or transfer [8, 12, 13].

Unlike the $^8\text{B}+^{208}\text{Pb}$ result at 170 MeV, the total reaction cross-section, for the low energy experiments around the Coulomb barrier [14, 15], was found to be large as in the case of neutron-halo reactions.

In this last case of $^8\text{B}+^{58}\text{Ni}$ the inclusive break-up was measured [14] by detecting the ^7Be fragments. These data showed that in order to reproduce the experimental spectra, high order effects had to be considered in the breakup process. Summarising the experimental evidence shows that the reaction dynamics for neutron-halo and proton-halo nuclei seems to be different. The availability at ISOLDE of a post accelerated ^8B beam gives the possibility to investigate the above topic with a better precision than the one so far available.

2. Experiment

The experiment was performed at HIE-ISOLDE facility at CERN with the first post-accelerated ^8B beam [16]. The ^8B beam was produced from the protons coming from the CERN PSBooster with energy of 1.4 GeV through the reaction on a multiwalled carbon nanotube target (CNT). A molecular $^8\text{BF}_2$ beam was extracted from the target; ^8B was charge bred in REX-EBIS to 3^+ and accelerated to an energy of 4.9 A MeV. The final ^8B average intensity on target was ~ 400 pps. The reaction target was a 1.021 mg/cm^2 isotopically enriched ^{64}Zn target. This was placed at an angle of 30° with respect to the beam direction, in order to allow measurements at 90° . The experimental setup consisted of six silicon telescopes made of two stages of detectors, $40\ \mu\text{m}$ and $1000\ \mu\text{m}$ thick respectively. Each detector was a $50\times 50\text{ mm}^2$ Double Sided Silicon Strip detector (DSSSD) segmented into $16+16$ strips [17]. The large segmentation allowed for high angular resolution for the angular distribution measurement which was, in fact, limited by the statistics.

3. Results

The $^8\text{B}+^{64}\text{Zn}$ elastic scattering angular distribution is shown in fig. 1 as a ratio to the Rutherford cross-section. Unlike the neutron-halo nucleus ^{11}Be at similar $E_{\text{c.m.}}/V_{\text{C}}$ [10, 11], a clear Coulomb-nuclear interference peak is visible at around 35° .

The measured angular distribution is compared with continuum-discretized coupled-channels (CDCC) calculations in which the ^8B is described as a two-cluster system, a ^7Be and a proton. Details of the calculations are reported in [16]. Fig. 1 shows the result of these calculations.




Figure 1. Experimental elastic scattering angular distribution for $^8\text{B}+^{64}\text{Zn}$ (symbols). CDCC calculations with (full line) and without (dashed line) coupling to the continuum.




Figure 2. Experimental ^7Be angular distribution (symbols). Continuous blue line: EBU from CDCC calculations. Dashed red line: NEB. Continuous black line: sum of EBU+NEB contributions.

The cross-section obtained from the CDCC calculations is shown in fig. 1 as a solid line. To show the effect of the ^8B elastic breakup on the elastic data, the calculation omitting the coupling to the continuum channels (dashed line) is also shown for comparison. From the figure it is possible to see that the elastic breakup produces a small suppression of the elastic cross-section, but does not suppress completely the Coulomb-nuclear interference effect. Figure 2 shows the angular distribution for ^7Be events. The present breakup data is inclusive, implying that several processes can in principle contribute to the ^7Be yield: on one side, the elastic breakup (EBU) process, in which the ^8B is dissociated into $^7\text{Be}+p$, leaving the target in its ground state; on the other side, the nonelastic breakup (NEB) processes in which the dissociated proton interacts non-elastically with the target, including non-capture breakup accompanied by target excitation, proton absorption by the target (incomplete fusion) and proton transfer.

The agreement between the experiment and the sum of EBU and NEB cross-section is fairly good, with only some underestimation of the data for $\theta_{\text{lab}} < 20^\circ$. From this comparison it can

be observed that the dominant reaction mechanism for the ${}^7\text{Be}$ production at forward angles is the EBU.

4. Summary

The ${}^8\text{B}+{}^{64}\text{Zn}$ elastic scattering angular distribution unlike the neutron-halo case, shows the presence of the Coulomb-nuclear interference peak. The comparison with CDCC calculations disclosed the evidence that reaction dynamics for the proton-halo ${}^8\text{B}$ shows only modest effects of coupling to the continuum and its total reaction cross-section is similar to that of ordinary weakly bound nuclei on the same target.

The inclusive angular and energy ${}^7\text{Be}$ distributions distinctively shows a dominance of elastic breakup at small angles, whereas non-elastic breakup becomes non negligible only at larger angles.

5. Acknowledgments

This work has been supported by the European Commission Horizon 2020 framework transnational access program of the EU project ENSAR2 (Grant no. 654002).

References

- [1] L.F. Canto, P.R.S. Gomes, R. Donangelo, J. Lubian, and M.S. Hussein. Recent developments in fusion and direct reactions with weakly bound nuclei. *Physics Reports*, 596:1 – 86, 2015. Recent developments in fusion and direct reactions with weakly bound nuclei.
- [2] A. García-Camacho, G. Blanchon, A. Bonaccorso, and D. M. Brink. All orders proton breakup from exotic nuclei. *Phys. Rev. C*, 76:014607, Jul 2007.
- [3] J.F. Liang et al. Dynamic polarization in the coulomb breakup of loosely bound ${}^{17}\text{F}$. *Phys. Lett. B*, 681:22, 2009.
- [4] B. Blank et al. Total interaction and proton-removal cross-section measurements for the proton-rich isotopes ${}^7\text{Be}$, ${}^8\text{B}$, and ${}^9\text{C}$. *Nucl. Phys. A*, 624(2):242 – 256, 1997.
- [5] C.H. Dasso, S.M. Lenzi, and A. Vitturi. Dominance of nuclear processes in the dissociation of ${}^8\text{B}$. *Nucl. Phys. A*, 639(3):635 – 653, 1998.
- [6] Angela Bonaccorso, David M. Brink, and Carlos A. Bertulani. Proton vs neutron halo breakup. *Phys. Rev. C*, 69:024615, Feb 2004.
- [7] A.M. Sánchez-Benítez et al. Study of the elastic scattering of ${}^6\text{He}$ on ${}^{208}\text{Pb}$ at energies around the Coulomb barrier. *Nucl. Phys. A*, 803(1):30 – 45, 2008.
- [8] A. Di Pietro et al. Reactions induced by the halo nucleus ${}^6\text{He}$ at energies around the Coulomb barrier. *Phys. Rev. C*, 69:044613, Apr 2004.
- [9] M. Cubero et al. Do halo nuclei follow Rutherford elastic scattering at energies below the barrier? The case of ${}^{11}\text{Li}$. *Phys. Rev. Lett.*, 109:262701, Dec 2012.
- [10] A. Di Pietro et al. Elastic scattering and reaction mechanisms of the halo nucleus ${}^{11}\text{Be}$ around the Coulomb Barrier. *Phys. Rev. Lett.*, 105:022701, Jul 2010.
- [11] A. Di Pietro et al. Experimental study of the collision ${}^{11}\text{Be} + {}^{64}\text{Zn}$ around the Coulomb barrier. *Phys. Rev. C*, 85:054607, May 2012.
- [12] R. Raabe et al. No enhancement of fusion probability by the neutron halo of ${}^6\text{He}$. *Nature*, 431:823 – 826, 2004.
- [13] J.P. Fernández-García et al. ${}^{11}\text{Li}$ breakup on ${}^{208}\text{Pb}$ at energies around the Coulomb Barrier. *Phys. Rev. Lett.*, 110:142701, Apr 2013.
- [14] V. Guimarães et al. Nuclear and Coulomb interaction in ${}^8\text{B}$ breakup at sub-Coulomb energies. *Phys. Rev. Lett.*, 84:1862–1865, Feb 2000.
- [15] M. Mazzocco et al. Elastic scattering for the ${}^8\text{B}$ and ${}^7\text{Be} + {}^{208}\text{Pb}$ systems at near-Coulomb barrier energies. *Phys. Rev. C*, 100:024602, Aug 2019.
- [16] R. Spart, A. Di Pietro, P. Figuera, O. Tengblad, A.M. Moro, I. Martel, J.P. Fernández-García, Jin Lei, L. Acosta, M.J.G. Borge, G. Bruni, J. Cederkll, T. Davinson, J.D. Ovejas, L.M. Fraile, D. Galaviz, J. Halkjaer Jensen, B. Jonson, M. La Cognata, A. Perea, A.M. Sánchez-Bentéz, N. Soi, and S. Vials. Probing proton halo effects in the $8\text{b}+64\text{zn}$ collision around the coulomb barrier. *Physics Letters B*, 820:136477, 2021.

- [17] G. Marquez-Durn, L. Acosta, R. Berjillos, J.A. Dueas, J.A. Labrador, K. Rusek, A.M. Sanchez-Bentez, and I. Martel. GLORIA: A compact detector system for studying heavy ion reactions using radioactive beams. *Nucl. Instr. and Meth. A*, 755:69 – 77, 2014.