



## Soft operators in $C^*$ -algebras

Downloaded from: <https://research.chalmers.se>, 2026-04-07 09:03 UTC

Citation for the original published paper (version of record):

Thiel, H., Vilalta Vila, E. (2024). Soft operators in  $C^*$ -algebras. *Journal of Mathematical Analysis and Applications*, 536(1). <http://dx.doi.org/10.1016/j.jmaa.2024.128167>

N.B. When citing this work, cite the original published paper.



Contents lists available at ScienceDirect

## Journal of Mathematical Analysis and Applications

journal homepage: [www.elsevier.com/locate/jmaa](http://www.elsevier.com/locate/jmaa)

## Regular Articles

Soft operators in  $C^*$ -algebras <sup>☆</sup>Hannes Thiel <sup>\*</sup>, Eduard Vilalta

Department of Mathematical Sciences, Chalmers University of Technology and University of Gothenburg,  
Gothenburg SE-412 96, Sweden



## ARTICLE INFO

## Article history:

Received 9 July 2023

Available online 27 January 2024

Submitted by M. Mathieu

## Keywords:

 $C^*$ -Algebras

Global Glimm Problem

Cuntz semigroups

Soft operators

## ABSTRACT

We say that a  $C^*$ -algebra is soft if it has no nonzero unital quotients, and we connect this property to the Hjelmberg-Rørørdam condition for stability and to property (S) of Ortega-Perera-Rørørdam. We further say that an operator in a  $C^*$ -algebra is soft if its associated hereditary subalgebra is, and we provide useful spectral characterizations of this concept. Of particular interest are  $C^*$ -algebras that have an abundance of soft elements in the sense that every hereditary subalgebra contains an almost full soft element. We show that this property is implied by the Global Glimm Property, and that every  $C^*$ -algebra with an abundance of soft elements is nowhere scattered. This sheds new light on the long-standing Global Glimm Problem of whether every nowhere scattered  $C^*$ -algebra has the Global Glimm Property.

© 2024 Published by Elsevier Inc.

## 1. Introduction

A recurring property in the study of non-simple  $C^*$ -algebras that ensures sufficient noncommutativity is the notion of having no nonzero elementary ideal-quotients; see, for example, [18,37,2]. Given the stark contrast between such  $C^*$ -algebras and those that are scattered (as defined by Jensen in [25]), the property was termed *nowhere scatteredness* in [45].

Despite this notion being the natural substitute for simplicity in a variety of scenarios, one often finds oneself working with the more technical condition of having the *Global Glimm Property*. This property, introduced by Kirchberg and Rørørdam in [27], implies nowhere scatteredness and, in fact, it is known that both notions agree for a wide class of  $C^*$ -algebras. However, whether these two conditions are the same in general remains an open problem, known as the Global Glimm Problem; see [47, Section 1] for an overview

<sup>☆</sup> The first named author was partially supported by the Knut and Alice Wallenberg Foundation (KAW 2021.0140). The second named author was partially supported by MINECO (grant No. PID2020-113047GB-I00/AEI/10.13039/501100011033 and No. PRE2018-083419), and by the Departament de Recerca i Universitats de la Generalitat de Catalunya (grant No. 2021-SGR-01015).

<sup>\*</sup> Corresponding author.

E-mail addresses: [hannes.thiel@chalmers.se](mailto:hannes.thiel@chalmers.se) (H. Thiel), [vilalta@chalmers.se](mailto:vilalta@chalmers.se) (E. Vilalta).

URLs: <https://www.hannesthiel.org> (H. Thiel), <https://www.eduardvilalta.com> (E. Vilalta).

on the subject. In essence, and as in many other problems in mathematics, the Global Glimm Problem asks if a natural condition and its more useful (albeit more technical) strengthening coincide.

A new approach to attacking this problem was developed in [47], where the statement of the Global Glimm Problem was reformulated in terms of the Cuntz semigroup, a powerful invariant for  $C^*$ -algebras introduced by Cuntz in [16] and later studied more abstractly by Coward, Elliott, and Ivanescu in [15]. The theory of Cuntz semigroups, which was further developed in [3–6,1], has seen many important applications in the structure and classification of  $C^*$ -algebras; see, for example, [48,33,49,37,42,2]. For a recent introduction to the theory we refer to [20].

In order to exploit the approach from [47], it becomes imperative to deepen our understanding of the Global Glimm Property. In this paper, we do so by characterizing the property in terms of *soft elements*.

**Definition A (3.1).** Given a  $C^*$ -algebra  $A$ , we say that a positive element  $a \in A_+$  is *soft* if no nonzero quotient of  $\overline{aAa}$  is unital.

This definition is in analogy to the notion of having a soft Cuntz class, as introduced in [3], and draws inspiration from the characterization of stability for purely infinite  $C^*$ -algebras obtained in [26, Theorem 4.24]: A  $\sigma$ -unital, purely infinite  $C^*$ -algebra is stable if and only if none of its nonzero quotients is unital. In fact, despite never having a name, the notion of softness appears throughout the literature. For example, one could restate part of [12, Theorem B] by saying that a separable, nuclear  $\mathcal{O}_\infty$ -stable  $C^*$ -algebra has finite decomposition rank if and only if every positive element in  $A$  is soft.

We begin our investigation by first developing the basic theory of softness for  $C^*$ -algebras, where we say that a  $C^*$ -algebra is *soft* if it has no nonzero, unital quotient; see Definition 2.1. We relate this property to the Hjelmberg-Rørdam condition for stability and to property (S) of Ortega-Perera-Rørdam; see Proposition 2.14. We then give a spectral characterization of softness:

**Proposition B (3.6).** *A positive element  $a$  in a  $C^*$ -algebra  $A$  is soft if and only if for every closed ideal  $I \subseteq A$ , either  $a \in I$  or the spectrum of  $a + I \in A/I$  has 0 as a limit point.*

Section 4 is devoted to the introduction of three tailored definitions of softness for Cuntz classes, which we call *functional*, *weak*, and *strong* softness; see Definition 4.3. Strong softness implies weak softness, which in turn implies functional softness, and all three notions agree in Cuntz semigroups of residually stably finite  $C^*$ -algebras; see Proposition 4.6. In that case, softness of an element in the  $C^*$ -algebra can be characterized in terms of its Cuntz class:

**Proposition C (4.17).** *Let  $A$  be a residually stably finite  $C^*$ -algebra. Then an element  $a \in A_+$  is soft if and only if its Cuntz class  $[a] \in \text{Cu}(A)$  is strongly soft (equivalently: weakly soft, functionally soft).*

In order to connect the Global Glimm Problem to soft operators, we introduce the notion of *abundance of soft elements* for a  $C^*$ -algebra  $A$ , which means that for every  $a \in A_+$  and  $\varepsilon > 0$  there exists a positive, soft operator  $b$  in the hereditary sub- $C^*$ -algebra  $\overline{aAa}$  such that the cut-down  $(a - \varepsilon)_+$  is in the closed ideal generated by  $b$ ; see Definition 5.2. This definition is in the same spirit as the Global Glimm Property, which one could phrase as an ‘abundance of nilpotent elements’; see [47, Theorem 3.6]. (We recall the definition of the Global Glimm Property at the beginning of Section 7.)

Although not every nilpotent element is soft, it is nevertheless true that the Global Glimm Property implies an abundance of soft elements; see Proposition 7.7. We also show that having an abundance of soft elements implies nowhere scatteredness (Proposition 7.4), which therefore breaks the Global Glimm Problem into two subquestions; see Question 7.8.

We introduce in Definition 5.2 the notion of having an *abundance of strongly soft elements* for Cuntz semigroups. Using that soft operators have strongly soft Cuntz classes, it follows that  $C^*$ -algebras with an

abundance of soft elements have scaled Cuntz semigroups with an abundance of soft elements. We show that the converse also holds (Theorem 5.14), although it remains unclear if every strongly soft Cuntz class can be realized by a soft element (Question 4.18).

Our notion of abundance of soft elements has close connections with the 2-splitting property, which was introduced by Kirchberg and Rørdam in [28] to study when central sequence algebras have characters. In Theorem 5.14, we show that a  $C^*$ -algebra has an abundance of soft elements if and only if it has the *hereditary 2-splitting property* (see Definition 5.7), a natural strengthening of the 2-splitting property.

Using these characterizations, we provide a new description of the Global Glimm Property:

**Theorem D (7.14).** *A  $C^*$ -algebra has the Global Glimm Property if and only if it has an abundance of soft elements and its Cuntz semigroup is ideal-filtered.*

In upcoming work with Asadi-Vasfi, [8], we will exploit the abundance of soft elements to find bounds for some numerical  $C^*$ -algebraic invariants, such as the radius of comparison and the Cuntz covering dimension introduced in [46].

**Acknowledgments** The authors thank the referee for their comments and valuable suggestions.

## 2. Soft $C^*$ -algebras

In this section, we introduce the notion of *softness* for  $C^*$ -algebras; see Definition 2.1. We provide useful characterizations of softness in Proposition 2.9, and we relate it to the Hjelmberg-Rørdam condition for stability and to property (S) of Ortega-Perera-Rørdam; see Proposition 2.14.

We also prove basic permanence properties: Softness passes to ideals, quotients and extensions (Proposition 2.16), to tensor products (Proposition 2.19) and to inductive limits (Proposition 2.18).

**Definition 2.1.** We say that a  $C^*$ -algebra  $A$  is *soft* if for every proper, closed ideal  $I \subseteq A$  the quotient  $A/I$  is nonunital.

**Remark 2.2.** Our notion of softness for  $C^*$ -algebras is not related to that introduced by Farsi in [19]. The latter is defined in terms of soft polynomial relations. Our terminology is inspired by the close connection to soft elements in Cuntz semigroups (see Proposition 4.17) and is therefore ultimately based on the notion of soft intervals introduced by Goodearl and Handelman in [22]; see also [3, Section 5.3].

The next result follows immediately from the definition. It is, in some sense, the  $C^*$ -algebraic analog of [3, Proposition 5.3.16].

**Proposition 2.3.** *A simple  $C^*$ -algebra is either unital or soft.*

**Notation 2.4.** Let  $A$  be a  $C^*$ -algebra. The annihilator of  $a \in A_+$  is

$$\text{Ann}(a) = \{x \in A : xa = ax = 0\}.$$

We write  $\text{Ann}_A(a)$  whenever we want to stress that the annihilator is considered inside the algebra  $A$ . Note that  $\text{Ann}(a)$  is a hereditary sub- $C^*$ -algebra of  $A$ .

If  $a, b \in A_+$ , then:

- we write  $a \ll b$  if  $b$  acts as a unit on  $a$ , that is, if  $a = ab$ ;
- we write  $a \triangleleft b$  to mean that  $a$  belongs to the closed ideal generated by  $b$ , that is, if  $a \in \overline{\text{span}}AbA$ .

- we write  $a \perp b$  if  $ab = 0$ .

**Lemma 2.5.** *Let  $\pi: A \rightarrow B$  be a surjective  $*$ -homomorphism between  $C^*$ -algebras, and let  $a, b \in A_+$  satisfy  $a \ll b$ . Then  $\text{Ann}_B(\pi(b)) \subseteq \pi(\text{Ann}_A(a))$ .*

**Proof.** By adjoining a unit to  $A$  if necessary, we may assume that  $A$  is unital. Let  $x \in \text{Ann}_B(\pi(b))$ . Lift  $x$  to obtain  $y \in A$  with  $\pi(y) = x$ . Set  $z = (1 - b)y(1 - b) \in A$ . Then  $\pi(z) = x$ . Further, using that  $a \ll b$ , we have  $a(1 - b) = (1 - b)a = 0$ , and therefore  $z \in \text{Ann}_A(a)$ .  $\square$

**Lemma 2.6.** *Let  $A$  be a  $C^*$ -algebra, let  $B \subseteq A$  be a soft sub- $C^*$ -algebra, and let  $p \in B$  be a projection. Then  $\text{Ann}(p)$  is full in  $A$ .*

**Proof.** Let  $I$  denote the closed ideal of  $A$  generated by the annihilator  $\text{Ann}_A(p)$ , and let  $\pi: A \rightarrow A/I$  denote the quotient map. Since  $p \ll p$ , we can apply Lemma 2.5 at the first step to obtain

$$\text{Ann}_{A/I}(\pi(p)) \subseteq \pi(\text{Ann}_A(p)) \subseteq \pi(I) = \{0\}.$$

Hence, the projection  $\pi(p)$  has zero annihilator in  $A/I$ . Assuming  $\pi(p) \neq 0$ , it follows that  $\pi(p)$  is the unit of  $A/I$  and, thus, of  $\pi(B)$ . This contradicts the softness of  $B$  and shows that  $\pi(p) = 0$ , which implies  $A/I = \{0\}$  and so  $I = A$ .  $\square$

**Notation 2.7.** Following the notation from [9], we set

$$A_c = \{a \in A_+ : a \ll b \text{ for some } b \in A_+\}.$$

The (nonclosed) ideal of  $A$  generated by  $A_c$  is called the Pedersen ideal – it is the minimal dense ideal of  $A$ ; see [10, Theorem II.5.2.4].

Note that  $A_c$  is denoted by  $F(A)$  in [29] and [23]. We do not use this notation to avoid confusion with the central sequence algebra of  $A$ . In [10],  $A_c$  is denoted by  $A_+^c$ .

**Lemma 2.8.** *Let  $A$  be a  $C^*$ -algebra, let  $B \subseteq A$  be a soft sub- $C^*$ -algebra, and let  $a \in B_c$ . Then  $\text{Ann}(a)$  is full in  $A$ .*

**Proof.** Let  $I$  denote the closed ideal of  $A$  generated by  $\text{Ann}_A(a)$ . We will show that  $I = A$ . Choose  $b \in B_+$  such that  $a \ll b$ . Let  $f, g: \mathbb{R} \rightarrow [0, 1]$  be given by

$$f(t) = \begin{cases} 0, & t \in (-\infty, \frac{1}{2}] \\ 2t - 1, & t \in [\frac{1}{2}, 1] \\ 1, & t \in [1, \infty), \end{cases} \quad \text{and} \quad g(t) = \begin{cases} 0, & t \in (-\infty, 0] \\ 2t, & t \in [0, \frac{1}{2}] \\ 1, & t \in [\frac{1}{2}, \infty). \end{cases}$$

Then  $a \ll f(b) \ll g(b)$ .

Let  $J$  denote the closed ideal generated by  $g(b) - f(b)$ , and let  $\pi_J: A \rightarrow A/J$  denote the quotient map. Note that  $g(b) - f(b) \in \text{Ann}_A(a)$ . Thus,  $J \subseteq I$ .

Then  $\pi_J(f(b)) \ll \pi_J(g(b)) = \pi_J(f(b))$ , which shows that  $\pi_J(f(b))$  is a projection. Using that  $B$  is soft, it easily follows that  $\pi_J(B)$  is soft as well. Then, by Lemma 2.6, the annihilator of  $\pi_J(f(b))$  in  $A/J$  is full. Since  $a \ll f(b)$ , we can apply Lemma 2.5 to obtain

$$\text{Ann}_{A/J}(\pi_J(f(b))) \subseteq \pi_J(\text{Ann}_A(a)).$$

It follows that  $\pi_J(I) = A/J$ . Combined with  $J \subseteq I$ , we get  $I = A$ .  $\square$

The next result is similar to the Hjelmberg-Rørdam characterization of stability from [24]; see also Proposition 2.11 and the paragraph preceding it.

**Proposition 2.9.** *Let  $A$  be a  $C^*$ -algebra. Then the following are equivalent:*

- (1)  $A$  is soft;
- (2) for every  $a \in A_c$ , the algebra  $\text{Ann}(a)$  is full in  $A$ ;
- (3) for every  $a \in A_c$ , there exist  $b \in A_+$  with  $a \perp b$  and  $a \triangleleft b$ ;
- (4) for every  $a \in A_+$  and  $\varepsilon > 0$  there exist  $b, c \in A_+$  with  $\|a - b\| < \varepsilon$ ,  $b \perp c$ , and  $b \triangleleft c$ .

**Proof.** It follows from Lemma 2.8 that (1) implies (2). To show that (2) implies (3), assume that  $\text{Ann}(a)$  is full. This implies that  $a$  belongs to the closed ideal generated by  $\text{Ann}(a)$ , which allows us to choose a sequence  $(b_n)_n$  in  $\text{Ann}(a)_+$  such that  $a$  belongs to the closed ideal generated by  $\{b_n : n \in \mathbb{N}\}$ . Set  $b = \sum_n \frac{1}{2^n \|b_n\| + 1} b_n$ . Then  $b$  belongs to  $\text{Ann}(a)$ . Further, for each  $n$ , we have  $b_n \leq (2^n \|b_n\| + 1)b$ , which implies that  $b_n$  belongs to the closed ideal generated by  $b$ . Consequently,  $a$  belongs to the closed ideal generated by  $b$ .

To show that (3) implies (4), assume that (3) holds, and let  $a \in A_+$  and  $\varepsilon > 0$ . Set  $b = (a - \frac{\varepsilon}{2})_+$ . Then  $b \in A_c$  and  $\|a - b\| = \frac{\varepsilon}{2} < \varepsilon$ . Applying (3), we obtain  $c \in A_+$  such that  $b \perp c$  and  $b \triangleleft c$ .

Finally, to show that (4) implies (1), assume that (4) holds and let  $I \subseteq A$  be a proper closed ideal. To reach a contradiction, assume that  $A/I$  is unital. Choose  $a \in A_+$  such that its image in  $A/I$  is the unit. Applying (4) with  $\varepsilon = 1$ , we obtain  $b, c \in A_+$  such that  $\|a - b\| < 1$  and  $b \triangleleft c$ . It follows that the image of  $b$  in the quotient  $A/I$  is at distance less than 1 to the unit, and is therefore strictly positive. Thus, the image of  $c$  in  $A/I$  is zero, that is,  $c \in I$ . We get  $b \in I$ , which implies that  $A/I = \{0\}$ .  $\square$

**2.10.** Given two positive elements  $a, b$  in a  $C^*$ -algebra  $A$ , we say that  $a$  is *Cuntz subequivalent* to  $b$ , written  $a \preceq b$ , if there exists a sequence  $(r_n)_n$  in  $A$  such that  $a = \lim_n r_n b r_n^*$ . Further, one says that  $a$  is *Cuntz equivalent* to  $b$ , and writes  $a \sim b$ , if  $a \preceq b \preceq a$ .

The *Cuntz semigroup*  $\text{Cu}(A)$  of a  $C^*$ -algebra  $A$  is the quotient  $(A \otimes \mathbb{K})_+ / \sim$  with the order induced by the Cuntz subequivalence and the addition induced by addition of orthogonal elements; we refer to [7,3] for details.

We let  $\text{QT}(A)$  denote the cone of  $[0, \infty]$ -valued, lower-semicontinuous 2-quasitraces on  $(A \otimes \mathbb{K})_+$ . Each element  $[a] \in \text{Cu}(A)$  induces a linear, lower semicontinuous map  $\widehat{[a]}: \text{QT}(A) \rightarrow [0, \infty]$  by  $\widehat{[a]}(\tau) = \lim_{n \rightarrow \infty} \tau(a^{1/n})$  for  $\tau \in \text{QT}(A)$ ; see [17,34,1].

In [24], Hjelmberg and Rørdam introduced a property for  $C^*$ -algebras that characterizes stability in the  $\sigma$ -unital case. We will say that a  $C^*$ -algebra has the *Hjelmberg-Rørdam property* if, for every  $a \in A_c$ , there exists  $b \in A_+$  such that  $a \perp b$ , and such that  $a = x^*x$  and  $b = xx^*$  for some  $x \in A$ .

**Proposition 2.11.** *A  $C^*$ -algebra  $A$  has the Hjelmberg-Rørdam property if and only if for every  $a \in A_c$  there exists  $b \in A_+$  such that  $a \perp b$  and  $[a] \leq [b]$  in  $\text{Cu}(A)$ .*

**Proof.** For every  $x \in A$ , we have  $[x^*x] = [xx^*]$  in  $\text{Cu}(A)$ . This shows the forward implication.

Conversely, let  $a \in A_c$  and take  $\varepsilon > 0$ . By assumption, there exists  $b \in A_+$  such that  $a \perp b$  and  $[a] \leq [b]$  in  $\text{Cu}(A)$ . Thus, it follows from [38, Proposition 2.4] that there exists  $x \in A$  satisfying

$$(a - \varepsilon)_+ = xx^*, \quad \text{and} \quad x^*x \in \overline{bAb}.$$

In particular, note that  $xx^*$  and  $x^*x$  are orthogonal. Applying [24, Proposition 2.2], we see that  $A$  has the Hjelmberg-Rørdam property.  $\square$

For elements  $[a], [b]$  in a Cuntz semigroup, one writes  $[a] <_s [b]$  if  $(n+1)[a] \leq n[b]$  for some  $n \in \mathbb{N}$ . Following [29, Definition 4.1], we say that a  $C^*$ -algebra  $A$  has *property (S)* if for every  $a \in A_c$  there exists  $b \in A_+$  such that  $a \perp b$  and  $[a] <_s [b]$  in  $\text{Cu}(A)$ .

**Proposition 2.12.** *A  $C^*$ -algebra  $A$  has property (S) if and only if for every  $a \in A_c$  there exists  $b \in A_+$  such that  $a \perp b$  and  $\widehat{[a]} \leq \widehat{[b]}$ .*

**Proof.** Let  $[a], [b]$  in  $\text{Cu}(A)$ . In general, one has that  $[a] <_s [b]$  implies  $\widehat{[a]} \leq \widehat{[b]}$ . This shows the forward implication.

For the converse, let  $a \in A_c$ . Using an argument as in the proof of Lemma 2.8, we can find  $b \in A_c$  such that  $a \ll b$ . Applying the assumption for  $b$ , we obtain  $c \in A_+$  such that  $b \perp c$  and  $\widehat{[b]} \leq \widehat{[c]}$ .

Note that  $[a]$  is way-below  $[b]$  in  $\text{Cu}(A)$ , denoted  $[a] \ll [b]$ ; see Paragraph 4.1 for details. This allows us to choose  $\varepsilon > 0$  such that  $[a] \ll [(b-\varepsilon)_+]$ . Using that  $[(b-\varepsilon)_+] \ll [b]$ , we can apply [34, Lemma 2.2.5] to deduce that  $\widehat{[(b-\varepsilon)_+]}$  is way-below  $2\widehat{[b]}$  (and hence also  $2\widehat{[c]}$ ) in the semigroup of linear, lower-semicontinuous functions  $\text{QT}(A) \rightarrow [0, \infty]$ . This allows us to choose  $\delta > 0$  such that  $\widehat{[(b-\varepsilon)_+]} \leq 2\widehat{[(c-\delta)_+]}$ .

Note that  $b + (c-\delta)_+$  belongs to  $A_c$ . Applying the assumption again, we obtain  $d \in A_+$  such that

$$b + (c-\delta)_+ \perp d, \quad \text{and} \quad \widehat{[b]} + \widehat{[(c-\delta)_+]} \leq \widehat{[d]},$$

where at the second inequality we have used that  $b \perp (c-\delta)_+$ .

Then

$$2\widehat{[(b-\varepsilon)_+]} \leq \widehat{[b]} + \widehat{[(b-\varepsilon)_+]} \leq \widehat{[b]} + 2\widehat{[(c-\delta)_+]} \leq \widehat{[(c-\delta)_+]} + \widehat{[d]}.$$

Thus,  $\widehat{[(b-\varepsilon)_+]} <_s \widehat{[(c-\delta)_+]} + \widehat{[d]}$ . Using at the first step that  $[a] \ll [(b-\varepsilon)_+]$  and applying [3, Theorem 5.2.13], we get

$$[a] <_s [(c-\delta)_+] + [d] = [(c-\delta)_+ + d].$$

Since  $a \perp (c-\delta)_+ + d$ , we see that  $(c-\delta)_+ + d$  has the desired properties.  $\square$

**Remark 2.13.** For positive elements  $a$  and  $b$  in a  $C^*$ -algebra, we have  $a \triangleleft b$  if and only if  $[a] \leq \infty[b]$  in  $\text{Cu}(A)$ ; see, for example, [3, Section 5.1].

Therefore, Propositions 2.11, 2.12 and 2.9 show that the Hjelmberg-Rørdam property, property (S), and softness for a  $C^*$ -algebra  $A$ , can be characterized in very similar ways: For every  $a \in A_c$  there exists  $b \in A_+$  such that  $a \perp b$  and

- $[a] \leq [b]$  for the Hjelmberg-Rørdam property;
- $[a] \leq [b]$  for property (S);
- $[a] \leq \infty[b]$  for softness.

For elements  $[a], [b]$  in a Cuntz semigroup, we have the following implications:

$$[a] \leq [b] \quad \Rightarrow \quad \widehat{[a]} \leq \widehat{[b]} \quad \Rightarrow \quad [a] \leq \infty[b].$$

This proves the next result.

**Proposition 2.14.** *For every  $C^*$ -algebra, the following implications hold:*

$$\text{Hjelmberg-Rørdam property} \quad \Rightarrow \quad \text{property (S)} \quad \Rightarrow \quad \text{softness}.$$

**Proposition 2.15** ([29, Proposition 4.5]). *A separable  $C^*$ -algebra has property (S) if and only if it is soft and has no bounded quasitraces.*

We now turn to permanence properties of softness. With a view towards the close connection between softness, the Global Glimm Property, and nowhere scatteredness discussed in Section 7, we remark that similar permanence properties hold for nowhere scatteredness ([45, Section 4]) and the Global Glimm Property ([47, Section 3]).

**Proposition 2.16.** *Let  $A$  be  $C^*$ -algebra, and let  $I \subseteq A$  be a closed ideal. Then  $A$  is soft if and only if  $I$  and  $A/I$  are soft.*

**Proof.** Assume first that  $A$  is soft, and let  $I$  be a closed ideal of  $A$ . Clearly,  $A/I$  is soft. To see that  $I$  is also soft, assume for the sake of contradiction that there is a proper ideal  $J \subseteq I$  such that  $I/J$  is unital, with unit  $e \in I/J$ . Then  $I/J$  is a unital ideal of  $A/J$ , which implies that  $A/J$  decomposes as the direct sum of  $I/J$  and the ideal  $K = \text{Ann}_{A/J}(e)$ . It follows that  $I/J = (A/J)/K$  is a proper, unital quotient of  $A$ , a contradiction.

Now assume that  $I$  and  $A/I$  are soft. Arguing once again by contradiction, assume that there exists a proper ideal  $J \subseteq A$  such that  $A/J$  is unital. Then,  $A/(I + J)$  is isomorphic to a quotient of both  $A/I$  and  $A/J$ . Since  $A/I$  is soft and  $A/J$  is unital, we get that  $A/(I + J)$  is both soft and unital and thus  $A = I + J$ .

It follows that  $A/J \cong I/(I \cap J)$ . Using that  $I$  is soft, we see that  $A = J$ . This contradicts the fact that  $J$  is proper.  $\square$

One says that a  $C^*$ -algebra  $A$  is *approximated* by a family  $(A_\lambda)_{\lambda \in \Lambda}$  of sub- $C^*$ -algebras  $A_\lambda \subseteq A$  if for every finite collection  $a_1, \dots, a_n \in A$  and  $\varepsilon > 0$  there exists  $\lambda \in \Lambda$  and  $b_1, \dots, b_n \in A_\lambda$  such that  $\|a_j - b_j\| < \varepsilon$  for  $j = 1, \dots, n$ .

**Proposition 2.17.** *A  $C^*$ -algebra is soft whenever it is approximated by a family of soft sub- $C^*$ -algebras.*

**Proof.** Let  $A$  be a  $C^*$ -algebra, and let  $A_\lambda \subseteq A$  be soft sub- $C^*$ -algebras that approximate  $A$ . To show that  $A$  is soft, we verify condition (4) in Proposition 2.9. Let  $a \in A_+$  and  $\varepsilon > 0$ . Choose  $\lambda$  and  $\bar{a} \in A_\lambda$  such that  $\|a - \bar{a}\| < \frac{\varepsilon}{2}$ . Taking additional care, we may assume  $\bar{a}$  to be positive; see, for example, the arguments in the proof of [44, Proposition 3.7].

Using that  $A_\lambda$  is soft, we can apply Proposition 2.9 to obtain  $b, c \in (A_\lambda)_+$  such that  $\|\bar{a} - b\| < \frac{\varepsilon}{2}$ ,  $b \perp c$ , and  $b \in \overline{\text{span}}A_\lambda c A_\lambda$ . Then

$$\|a - b\| \leq \|a - \bar{a}\| + \|\bar{a} - b\| < \varepsilon, \quad \text{and} \quad b \in \overline{\text{span}}A_\lambda c A_\lambda \subseteq \overline{\text{span}}AcA$$

which shows that  $b$  and  $c$ , viewed in  $A$ , have the desired properties.  $\square$

**Proposition 2.18.** *Softness passes to inductive limits.*

**Proof.** This follows from Propositions 2.16 and 2.17 as in the proof of [43, Proposition 2.4]. We include some details for the convenience of the reader. Let  $A$  be the inductive limit of an inductive system of soft  $C^*$ -algebras  $A_j$ . For each  $j$ , consider the natural map  $A_j \rightarrow A$  and its image  $B_j \subseteq A$ . Since  $B_j$  is a quotient of  $A_j$ , it is soft by Proposition 2.16. Now, using that  $A$  is approximated by the family  $B_j$ , it follows from Proposition 2.17 that  $A$  is soft.  $\square$

**Proposition 2.19.** *Let  $A, B$  be  $C^*$ -algebras. Then the following are equivalent:*

- (1)  $A$  or  $B$  is soft;
- (2) The maximal tensor product  $A \otimes_{\max} B$  is soft;
- (3) The tensor product  $A \otimes_{\varrho} B$  is soft for some (equivalently: any) cross-norm  $\varrho$ ;
- (4) The minimal tensor product  $A \otimes_{\min} B$  is soft.

**Proof.** For every cross-norm  $\varrho$ , the tensor product  $A \otimes_{\varrho} B$  is a quotient of  $A \otimes_{\max} B$ , and  $A \otimes_{\min} B$  is a quotient of  $A \otimes_{\varrho} B$ . Since softness passes to quotients by Proposition 2.16, we see that (2) implies (3), and that (3) implies (4).

Let us show that (4) implies (1). To reach a contradiction, assume that neither  $A$  nor  $B$  is soft. Then there are proper closed ideals  $I \subseteq A$  and  $J \subseteq B$  such that  $A/I$  and  $B/J$  are unital. The quotient maps  $A \rightarrow A/I$  and  $B \rightarrow B/J$  induce a natural surjective  $*$ -homomorphism  $A \otimes_{\min} B \rightarrow (A/I) \otimes_{\min} (B/J)$  (see, for example, [10, II.9.6.6]). Since  $(A/I) \otimes_{\min} (B/J)$  is unital, this shows that  $A \otimes_{\min} B$  is not soft.

Finally, let us show that (1) implies (2). To reach a contradiction, assume that  $A \otimes_{\max} B$  is not soft. Then there exists a  $*$ -representation  $\pi: A \otimes_{\max} B \rightarrow \mathcal{B}(H)$  on some Hilbert space  $H$  such that the identity  $1 \in \mathcal{B}(H)$  is contained in the image of  $\pi$ . By [10, Theorem II.9.2.1] there exist unique  $*$ -representations  $\pi_A: A \rightarrow \mathcal{B}(H)$  and  $\pi_B: B \rightarrow \mathcal{B}(H)$  such that  $\pi(a \otimes b) = \pi_A(a)\pi_B(b) = \pi_B(b)\pi_A(a)$  for all  $a \in A$  and  $b \in B$ .

Choose approximate units  $(a_\lambda)_\lambda$  for  $A$  and  $(b_\mu)_\mu$  for  $B$ . Then  $(a_\lambda \otimes b_\mu)_{\lambda, \mu}$  is an approximate unit for  $A \otimes_{\max} B$  and it follows that  $\pi(a_\lambda \otimes b_\mu)$  converges to  $1 \in \mathcal{B}(H)$  in norm. Thus, by setting  $a = a_\lambda$  and  $b = b_\mu$  for sufficiently large  $\lambda$  and  $\mu$ , the operator  $\pi(a \otimes b)$  is invertible. Using that  $\pi(a \otimes b) = \pi_A(a)\pi_B(b) = \pi_B(b)\pi_A(a)$ , we deduce that  $\pi_A(a)$  and  $\pi_B(b)$  are invertible. This implies that  $1$  belongs to the image of  $\pi_A$  and  $\pi_B$ . Hence,  $\ker(\pi_A)$  and  $\ker(\pi_B)$  are proper closed ideals of  $A$  and  $B$  with unital quotients, showing that neither  $A$  nor  $B$  is soft.  $\square$

The next result shows that softness is a stable property.

**Proposition 2.20.** *Let  $A$  be a  $C^*$ -algebra, and let  $n \geq 1$ . Then  $A$  is soft if and only if  $M_n(A)$  is.*

**Proof.** The forward implication follows directly from Proposition 2.19.

Conversely, assume that  $M_n(A)$  is soft. Thus, using Proposition 2.19, we know that either  $M_n(\mathbb{C})$  or  $A$  is soft. Since  $M_n(\mathbb{C})$  is not soft, we must have  $A$  soft, as desired.  $\square$

**Question 2.21.** Does softness satisfy the Löwenheim-Skolem condition? Here, a property  $\mathcal{P}$  of  $C^*$ -algebras is said to satisfy the *Löwenheim-Skolem condition* if for every  $C^*$ -algebra  $A$  satisfying  $\mathcal{P}$ , there exists a  $\sigma$ -complete, cofinal subset of separable sub- $C^*$ -algebras of  $A$  that each satisfy  $\mathcal{P}$ ; see [45, Section 4] for further details. This is closely related to Blackadar's concept of a *separably inheritable* property [10, Definition II.8.5.1]. Every separably inheritable property satisfies the Löwenheim-Skolem condition.

### 3. Soft operators

In this section, we introduce and study the notion of *softness* for operators in  $C^*$ -algebras; see Definition 3.1. We will see in Section 5 that this concept is closely related to the notion of softness for elements in a Cuntz semigroup.

**Definition 3.1.** We say that an element  $x$  in  $C^*$ -algebra  $A$  is *soft* if the hereditary sub- $C^*$ -algebra  $\overline{x^*Ax}$  is soft.

The next result shows that we could equivalently use  $\overline{xAx^*}$  instead of  $\overline{x^*Ax}$  in the above definition. We give further characterizations of softness for positive operators in Proposition 3.6 below.

**Proposition 3.2.** *Let  $A$  be a  $C^*$ -algebra, and  $x \in A$ . Then the following are equivalent:*

- (1) *The element  $x$  is soft.*
- (2) *The element  $x^*$  is soft.*
- (3) *The element  $x^*x$  is soft.*
- (4) *The element  $xx^*$  is soft.*

**Proof.** By [10, Proposition II.3.4.2], we have  $\overline{x^*Ax} = \overline{x^*xAx^*x}$ . This shows that (1) and (3) are equivalent, and similarly that (2) and (4) are equivalent. The equivalence of (1) and (2) follows using that  $\overline{x^*Ax}$  and  $\overline{xAx^*}$  are isomorphic. Indeed, if  $x = v|x|$  is the polar decomposition of  $x$  in  $A^{**}$ , then  $a \mapsto vav^*$  is a well-defined isomorphism  $\overline{x^*Ax} \rightarrow \overline{xAx^*}$ ; see also [30, Lemma 4.2].  $\square$

**Remark 3.3.** The property of softness for operators depends on the containing  $C^*$ -algebra. For example, if  $a$  is a strictly positive element in a simple, non-unital  $C^*$ -algebra  $A$ , then  $a$  is soft in  $A$  since  $\overline{aAa} = A$ , which is soft by Proposition 2.3, while  $a$  is not soft as an element of the commutative  $C^*$ -algebra  $C^*(a)$ .

We say that a  $C^*$ -algebra is *residually projectionless* if none of its quotients contains a nonzero projection. (In [35], such  $C^*$ -algebras are simply called ‘projectionless’, but we feel that ‘residually projectionless’ is more descriptive.)

**Proposition 3.4.** *A  $C^*$ -algebra  $A$  is residually projectionless if and only if every element of  $A$  is soft.*

**Proof.** Let  $A$  be a  $C^*$ -algebra. Assume first that  $A$  is residually projectionless. Given any hereditary sub- $C^*$ -algebra  $B \subseteq A$ , and a closed ideal  $I \subseteq B$ , let  $J$  be the closed ideal of  $A$  generated by  $I$ . Then the quotient  $B/I$  is naturally isomorphic to a hereditary sub- $C^*$ -algebra of  $A/J$ . Since  $A/J$  contains no nonzero projections,  $B/I$  cannot be unital and nonzero. This shows that every hereditary sub- $C^*$ -algebra of  $A$  is soft, and hence so is every element of  $A$ .

Conversely, assume that  $A$  is not residually projectionless. Choose a closed ideal  $I \subseteq A$  and a nonzero projection  $p \in A/I$ . Pick a positive lift  $a \in A$  of  $p$ . Then  $a$  is not soft since  $\overline{aAa}$  has the nonzero, unital quotient  $p(A/I)p$ .  $\square$

**Notation 3.5.** Given  $\varepsilon > 0$ , we let  $g_\varepsilon : \mathbb{R} \rightarrow \mathbb{R}$  be the continuous function given by

$$g_\varepsilon(t) = \begin{cases} 0, & \text{if } t \in (-\infty, 0] \cup [\varepsilon, \infty) \\ t, & \text{if } t \in [0, \frac{\varepsilon}{2}] \\ \varepsilon - t, & \text{if } t \in [\frac{\varepsilon}{2}, \varepsilon] \end{cases}$$

The equivalence of (1) and (3) in the next result is analogous to Bosa’s characterization of stable elements; see [11, Lemma 2.3].

**Proposition 3.6.** *Let  $A$  be a  $C^*$ -algebra, and  $a \in A_+$ . Then the following are equivalent:*

- (1) *The element  $a$  is soft.*
- (2) *For every closed ideal  $I \subseteq A$ , either  $a \in I$  or the spectrum of  $a + I \in A/I$  has 0 as a limit point.*
- (3) *We have  $a \triangleleft g_\varepsilon(a)$  for every  $\varepsilon > 0$ .*
- (4) *For every  $\varepsilon > 0$  there exists  $b \in \overline{aAa}_+$  with  $b \perp (a - \varepsilon)_+$  and  $a \triangleleft b$ .*

**Proof.** Let us show that (1) implies (2). Assuming that  $a$  is soft, let  $I \subseteq A$  be a closed ideal such that  $a \notin I$ . To reach a contradiction, assume that 0 is isolated in the spectrum of  $a + I$ . Then the characteristic

function of  $(0, \infty)$  is continuous on the spectrum of  $a + I$ , and applying functional calculus we obtain a projection in  $A/I$  that generates the same hereditary sub- $C^*$ -algebra as  $a + I$ . Thus,  $\overline{aAa}$  has a nonzero, unital quotient, which is the desired contradiction.

Let us show that (2) implies (3). Assuming (2), let  $\varepsilon > 0$ . Set  $I = \overline{\text{span}Ag_\varepsilon(a)A}$ . Then the spectrum of  $a + I$  is contained in  $\{0\} \cup [\varepsilon, \infty)$ . By assumption, this implies that  $a \in I$ , and thus  $a \triangleleft g_\varepsilon(a)$ .

It is clear that (3) implies (4). To show that (4) implies (1), let  $I \subseteq \overline{aAa}$  be a proper closed ideal, and let  $\pi: \overline{aAa} \rightarrow \overline{aAa}/I$  be the quotient map. To reach a contradiction, assume that  $\overline{aAa}/I$  is unital and  $\overline{aAa}/I \neq \{0\}$ . Since  $a$  is strictly positive,  $\pi(a)$  is invertible in  $\overline{aAa}/I$ , and we obtain  $\varepsilon > 0$  such that the spectrum of  $\pi(a)$  is contained in  $[\varepsilon, \infty)$ .

Applying the assumption for  $\frac{\varepsilon}{2}$ , we obtain a full element  $b \in \overline{aAa}_+$  such that  $b \perp (a - \frac{\varepsilon}{2})_+$ . We have  $\pi((a - \frac{\varepsilon}{2})_+) = (\pi(a) - \frac{\varepsilon}{2})_+$ , whose spectrum is contained in  $[\frac{\varepsilon}{2}, \infty)$ . Thus,  $\pi((a - \frac{\varepsilon}{2})_+)$  is strictly positive in  $\overline{aAa}/I$ , which implies that  $\pi(b) = 0$ , a contradiction.  $\square$

A completely positive map between  $C^*$ -algebras is said to have *order-zero* if it preserves zero-products among positive elements. There is a rich theory of such maps, most importantly the structure theorem of Winter-Zacharias [50], which characterizes order-zero maps as  $*$ -homomorphisms from the cone over the domain. Another characterization is given in [21, Corollary 4.8], where it is shown that the class of completely positive, order zero maps can equivalently be described as the class of positive, zero-product preserving maps.

It remains unclear if one can remove the positivity assumption for the soft element in the next result; see Question 3.9.

**Proposition 3.7.** *Let  $\varphi: A \rightarrow B$  be a completely positive, order-zero map between  $C^*$ -algebras, and let  $a \in A_+$  be soft. Then  $\varphi(a)$  is soft.*

**Proof.** We use the characterization of softness from Proposition 3.6(4). So let  $\varepsilon > 0$ . We need to find a positive element  $b \in \overline{\varphi(a)B\varphi(a)}$  orthogonal to  $(\varphi(a) - \varepsilon)_+$  and such that  $\varphi(a) \triangleleft b$ .

Using that  $a$  is soft, and applying Proposition 3.6(3), we see that  $a \triangleleft g_\varepsilon(a)$ . Set  $b = \varphi(g_\varepsilon(a))$ . We have  $g_\varepsilon(a) \leq a$ , and therefore  $b = \varphi(g_\varepsilon(a)) \leq \varphi(a)$ , and thus  $b \in \overline{\varphi(a)B\varphi(a)}$ .

Further, we have  $g_\varepsilon(a) \perp (a - \varepsilon)_+$ , and since  $\varphi$  preserves orthogonality among positive elements, we see that  $b \perp \varphi((a - \varepsilon)_+)$ . As noted in the proof of [3, Proposition 2.2.7], we have  $(\varphi(a) - \varepsilon)_+ \leq \varphi((a - \varepsilon)_+)$ , which implies that  $b$  is orthogonal to  $(\varphi(a) - \varepsilon)_+$ .

Let  $x, y \in A_+$  with  $x \triangleleft y$ . Then  $[x] \leq \infty[y]$  in  $\text{Cu}(A)$ , and since  $\varphi$  naturally induces a generalized  $\text{Cu}$ -morphism  $\text{Cu}(A) \rightarrow \text{Cu}(B)$  by [50, Corollary 4.5] (see also [3, 3.2.5]), we obtain that  $[\varphi(x)] \leq \infty[\varphi(y)]$  in  $\text{Cu}(B)$ , whence  $\varphi(x) \triangleleft \varphi(y)$ . Applying this argument for  $a \triangleleft g_\varepsilon(a)$ , we deduce that  $\varphi(a) \triangleleft b$ , which shows that  $b$  has the desired properties.  $\square$

Applying the above result for the quotient map by a closed ideal, we get:

**Proposition 3.8.** *Let  $I \subseteq A$  be a closed ideal in a  $C^*$ -algebra  $A$ , and let  $a \in A_+$  be soft. Then  $a + I \in A/I$  is soft.*

**Question 3.9.** Let  $\varphi: A \rightarrow B$  be a completely positive, order-zero map between  $C^*$ -algebras. Given a (not necessarily positive) soft element  $x \in A$ , is  $\varphi(x)$  soft?

#### 4. Shades of softness in $\text{Cu}$ -semigroups

In this section, we introduce the concepts of strong softness, weak softness and functional softness for elements in  $\text{Cu}$ -semigroups; see Definition 4.3. (We recall the definition of  $\text{Cu}$ -semigroups in Paragraph 4.1

below.) We show that every strongly soft element is weakly soft, and that every weakly soft element is functionally soft. For residually stably finite Cu-semigroups, we prove that these three notions coincide; see Proposition 4.6.

Functionally soft elements were first considered in [3, Definition 5.3.1], where they are just called ‘soft’ elements. Strongly soft elements were implicitly also considered in [3, Section 5.3]. We use the term ‘functionally soft’ to clarify the difference with the other notions.

We analyze the relation between soft elements in a C\*-algebra and (strongly) soft elements in its Cuntz semigroup in Proposition 4.17. In particular, in a stably finite C\*-algebra, a positive element is soft if and only if its Cuntz class is.

**4.1.** Given two elements  $x, y$  in a partially ordered monoid where suprema of increasing sequences exist, recall that we write  $x \ll y$  if, for every increasing sequence  $(z_n)_n$  such that  $y \leq \sup_n z_n$ , there exists  $n \in \mathbb{N}$  such that  $x \leq z_n$ .

Following [15], one says that a partially ordered, commutative monoid  $S$  is a Cu-semigroup if it satisfies the conditions below:

- (O1) Every increasing sequence in  $S$  has a supremum.
- (O2) Every element in  $S$  can be written as the supremum of a  $\ll$ -increasing sequence.
- (O3) Given  $x', x, y', y \in S$  with  $x' \ll x$  and  $y' \ll y$ , one has  $x' + y' \ll x + y$ .
- (O4) Given two increasing sequences  $(x_n)_n, (y_n)_n$  in  $S$ , one has  $\sup_n (x_n + y_n) = \sup_n x_n + \sup_n y_n$ .

As shown in [15], the Cuntz semigroup  $\text{Cu}(A)$  of any C\*-algebra  $A$  is a Cu-semigroup. Further, it was subsequently shown in [3], [34] and [1] respectively that  $\text{Cu}(A)$  also satisfies the following properties:

- (O5) Given  $x', x, y', y, z \in S$  such that  $x' \ll x, y' \ll y$  and  $x + y \leq z$ , there exists  $c \in S$  with  $x' + c \leq z \leq x + c$  and  $y' \ll c$ .
- (O6) Given  $x', x, y, z \in S$  such that  $x' \ll x \leq y + z$ , there exist elements  $e, f$  such that  $e \leq x, y, f \leq x, z$  and  $x' \leq e + f$ .
- (O7) Given  $x', x, y', y, z \in S$  such that  $x' \ll x \leq z$  and  $y' \ll y \leq z$ , there exists  $w \in S$  satisfying  $x', y' \ll w \leq z, x + y$ .

Recently, an additional property that the Cuntz semigroup of every C\*-algebra satisfies, termed (O8), has been uncovered; see [45, Theorem 7.4].

We will often use the following equivalent formulations of (O6) and (O7). This is a folklore result but, since we have been unable to locate a reference, we include a short proof here.

**Lemma 4.2.** *Let  $S$  be a Cu-semigroup. Then,*

- $S$  satisfies (O6) if and only if, whenever  $x' \ll x \leq y_1 + \dots + y_n$ , there exist  $e_1, \dots, e_n \in S$  such that  $x' \leq e_1 + \dots + e_n$ , and such that  $e_j \leq x, y_j$  for all  $j$ .
- $S$  satisfies (O7) if and only if, for given  $x'_j \ll x_j \leq z$  for  $j = 1, \dots, n$ , there exists  $w \in S$  satisfying  $x'_j \ll w \leq z, x_1 + \dots + x_n$ .

**Proof.** Note that the backwards implication is trivial in both cases. The proof of the forward implication is by induction on  $n$ . We only provide details for (i), since the proof of (ii) is analogous.

The statement follows directly from definition if  $n = 2$ . Let  $m \in \mathbb{N}$ , and assume that the forward implication is true for every  $n \leq m$ . Let  $x', x, y_1, \dots, y_{m+1} \in S$  be such that  $x' \ll x \leq y_1 + \dots + y_{m+1}$ . Take

$x'' \in S$  such that  $x' \ll x'' \ll x$ . Applying (O6) to  $x'' \ll x \leq (y_1 + \dots + y_m) + y_{m+1}$ , we obtain elements  $e, e_{m+1} \in S$  such that

$$x' \ll x'' \leq e + e_{m+1}, \quad e \leq x, y_1 + \dots + y_m, \quad \text{and} \quad e_{m+1} \leq x, y_{m+1}.$$

Now find  $e' \in S$  such that  $e' \ll e$  and  $x' \leq e' + e_{m+1}$ . Using the induction hypothesis at  $e' \ll e \leq y_1 + \dots + y_m$ , we obtain elements  $e_1, \dots, e_m$  such that  $e' \leq e_1 + \dots + e_m$ , and such that  $e_j \leq e, y_j$  for all  $j \leq m$ . It is readily checked that  $e_1, \dots, e_{m+1}$  satisfy the desired conditions.  $\square$

**Definition 4.3.** Let  $S$  be a Cu-semigroup, and let  $x \in S$ . We say that  $x$  is *strongly soft* if for every  $x' \in S$  satisfying  $x' \ll x$  there exists  $t \in S$  such that

$$x' + t \ll x, \quad \text{and} \quad x' \ll \infty t.$$

We say that  $x$  is *weakly soft* if for every  $x' \in S$  satisfying  $x' \ll x$  there exists  $n \geq 1$  and  $t_1, \dots, t_n \in S$  such that

$$x' + t_j \ll x \quad \text{for } j = 1, \dots, n, \quad \text{and} \quad x' \ll t_1 + \dots + t_n.$$

We say that  $x$  is *functionally soft* if for every  $x' \in S$  satisfying  $x' \ll x$  there exists  $n \geq 1$  such that

$$(n+1)x' \ll nx.$$

**Remark 4.4.** In [3, Definition 5.3.1], an element  $x$  in a Cu-semigroup  $S$  is defined to be ‘soft’ if for every  $x' \in S$  satisfying  $x' \ll x$  there exists  $n \in \mathbb{N}$  such that  $(n+1)x' \leq nx$ . Note that  $x$  is ‘soft’ in this sense if and only if  $x$  is functionally soft in the sense of Definition 4.3. Indeed, the backward implication is clear. Conversely, if  $x$  is ‘soft’ and  $x' \ll x$ , then choose  $x''$  satisfying  $x' \ll x'' \ll x$ . Applying the definition we get  $n$  such that  $(n+1)x'' \leq nx$ , and thus  $(n+1)x' \ll nx$ .

The notions from Definition 4.3 are closely related to the concept of *pure noncompactness* introduced in [17], and its generalization *weak pure noncompactness* from [3]. After recalling the definitions, we will see how these notions are related. We refer to [3, Section 5.1] for details on ideals and quotients of Cu-semigroups.

**Definition 4.5.** Let  $S$  be a Cu-semigroup, and let  $x \in S$ . One says that  $x$  is *purely noncompact* if for every ideal  $I \subseteq S$  such that the image  $x_I$  of  $x$  in the quotient  $S/I$  is compact, we have  $2x_I = x_I$ .

One says that  $x$  is *weakly purely noncompact* if for every ideal  $I \subseteq S$  such that  $x_I$  is compact, there exists  $n \in \mathbb{N}$  such that  $(n+1)x_I = nx_I$ .


We will say that a Cu-semigroup  $S$  is *stably finite* if for all  $x, y \in S$  with  $x + y \ll x$  we have  $y = 0$ . Note that this is more restrictive than the definition in [3], although both notions agree if  $S$  is simple. Further, we will say that  $S$  is *residually stably finite* if every quotient of  $S$  is stably finite.

A stably finite C\*-algebra has a stably finite Cuntz semigroup, and a residually stably finite C\*-algebra has a residually stably finite Cuntz semigroup.

**Proposition 4.6.** *Let  $S$  be a Cu-semigroup, and let  $x \in S$ . If  $x$  is strongly soft, then  $x$  is weakly soft, which in turn implies that  $x$  is functionally soft and purely noncompact. Further,  $x$  is weakly purely noncompact whenever it is purely noncompact or functionally soft.*

*If  $S$  satisfies (O5), then  $x$  is functionally soft if and only if  $x$  is weakly purely noncompact. If  $S$  satisfies (O5) and is residually stably finite, then all notions from Definitions 4.3 and 4.5 are equivalent.*

The implications are shown in the following diagram:



**Proof.** It is easy to see that strong softness implies weak softness, and that pure noncompactness implies weak pure noncompactness.

To show that weak softness implies functional softness, assume that  $x$  is weakly soft, and let  $x' \in S$  satisfy  $x' \ll x$ . We obtain  $n \geq 1$  and  $t_1, \dots, t_n \in S$  such that

$$x' + t_j \ll x \quad \text{for } j = 1, \dots, n, \quad \text{and} \quad x' \ll t_1 + \dots + t_n.$$

Then

$$(n + 1)x' \leq nx' + t_1 + \dots + t_n \ll nx.$$

To show that weak softness implies pure noncompactness, assume that  $x$  is weakly soft, and let  $I \subseteq S$  be an ideal such that  $\pi(x) \ll \pi(x)$ , where  $\pi: S \rightarrow S/I$  denotes the quotient map. Using that  $\pi$  preserves suprema of increasing sequences, we obtain  $x' \in S$  such that  $x' \ll x$  and  $\pi(x) \leq \pi(x')$ , which then implies  $\pi(x) = \pi(x')$ . By Definition 4.3, we obtain  $n \geq 1$  and  $t_1, \dots, t_n \in S$  such that

$$x' + t_j \ll x \quad \text{for } j = 1, \dots, n, \quad \text{and} \quad x' \ll t_1 + \dots + t_n.$$

For each  $j$ , we get  $\pi(x) + \pi(t_j) = \pi(x)$ , and therefore

$$2\pi(x) = \pi(x) + \pi(x') \leq \pi(x) + \pi(t_1) + \dots + \pi(t_n) = \pi(x),$$

as required.

It was shown in [3, Proposition 5.3.5] that functional softness implies weak pure noncompactness and that the converse implication holds if  $S$  satisfies (O5).

Lastly, assume that  $S$  satisfies (O5) and is residually stably finite, and that  $x$  is functionally soft. In this case, it was shown in [3, Lemma 5.3.8] that  $x$  is strongly soft. Note that a stronger notion of stable finiteness is used in [3], but for the argument in [3, Lemma 5.3.8] the weaker notion defined above suffices. For convenience, we include the short argument.

To verify that  $x$  is strongly soft, let  $x' \in S$  satisfy  $x' \ll x$ . Choose  $y', y \in S$  such that  $x' \ll y' \ll y \ll x$ .

Since  $x$  is functionally soft, we obtain  $n \in \mathbb{N}$  such that  $(n + 1)y \ll nx$ . Applying (O5) for  $y' \ll y \leq x$ , we obtain  $t \in S$  such that

$$y' + t \leq x \leq y + t.$$

Let  $I = \{s \in S : s \leq \infty t\}$  be the ideal generated by  $t$ , and let  $\pi: S \rightarrow S/I$  denote the quotient map. Then  $\pi(x) \leq \pi(y)$ , and therefore

$$n\pi(x) + \pi(x) = (n + 1)\pi(x) \leq (n + 1)\pi(y) \ll n\pi(x).$$

Using that  $S/I$  is stably finite, we get  $\pi(x) = 0$  and, consequently,  $x \leq \infty t$ . Using that  $x' \ll x \leq \infty t$ , we can choose  $t' \in S$  such that  $t' \ll t$  and  $x' \ll \infty t'$ . Since  $x' \ll y'$  and  $t' \ll t$ , we have  $x' + t' \ll y' + t \leq x$ , which shows that  $t'$  has the desired properties.  $\square$

Recall that a Cu-semigroup is said to satisfy *weak cancellation* if for all elements  $x, y, z$  with  $x + z \ll y + z$  we have  $x \ll y$ . It is easy to check that weak cancellation passes to quotients and implies stably finiteness. Hence, every weakly cancellative Cu-semigroup is residually stably finite.

It was shown in [40, Theorem 4.3] that the Cuntz semigroups of stable rank one  $C^*$ -algebras always satisfy weak cancellation.

**Corollary 4.7.** *Let  $A$  be a residually stably finite  $C^*$ -algebra. (For example, a  $C^*$ -algebra with stable rank one.) Then for elements in  $\text{Cu}(A)$ , all notions from Definitions 4.3 and 4.5 are equivalent.*

**Proposition 4.8.** *Let  $S$  be a Cu-semigroup, and let  $x \in S$  be compact. Then  $x$  is strongly soft if and only if  $x$  is weakly soft, if and only if  $x$  is purely noncompact, if and only if  $2x = x$ .*

*Further,  $x$  is functionally soft if and only if  $x$  is weakly purely noncompact, if and only if there exists  $n \in \mathbb{N}$  such that  $(n + 1)x = nx$ .*

**Proof.** If  $2x = x$ , then  $x$  is easily seen to be strongly soft. In general, strong softness implies weak softness, which in turn implies pure noncompactness, by Proposition 4.6. Further, if  $x$  is purely noncompact and compact, then  $2x = x$  by definition.

Similarly, if  $(n + 1)x = nx$  for some  $n$ , then  $x$  is easily seen to be functionally soft, which in turn implies that  $x$  is weakly purely noncompact by Proposition 4.6. Further, if  $x$  is weakly purely noncompact and compact, then  $(n + 1)x = nx$  for some  $n$  by definition.  $\square$

**Example 4.9.** Given  $k \geq 1$ , let  $E_k = \{0, 1, 2, \dots, k, \infty\}$ , where the order is algebraic and the sum of two elements  $x, y$  is  $x + y$  if  $x + y \leq k$  or  $\infty$  otherwise. Then  $E_k$  is a Cu-semigroup that satisfies (O5); see [3, Paragraph 5.1.16] for more details.

The element  $x = 1$  is compact and satisfies  $(k + 2)x = \infty = (k + 1)x$ , but not  $2x = x$ . Therefore,  $x$  is functionally soft and weakly purely noncompact, but it is neither strongly soft, nor weakly soft, nor purely noncompact.

**Example 4.10.** Rørdam showed in [39] that there exist unital  $C^*$ -algebras  $A$  such that the unit of  $A$  is finite, while the unit of  $M_2(A)$  is properly infinite. This implies that in  $\text{Cu}(A)$  the class  $[1_A]$  of the unit of  $A$  is a compact element satisfying  $3[1_A] = 2[1_A] \neq [1_A]$ . By Proposition 4.8,  $[1_A]$  is functionally soft and weakly purely noncompact, but it is neither strongly soft, nor weakly soft, nor purely noncompact.

Example 4.10 shows that the lower downwards implications in Proposition 4.6 cannot be reversed for elements in Cuntz semigroups. The situation for the other implications is unclear.

**Question 4.11.** For elements in the Cuntz semigroup of a  $C^*$ -algebra, does weak softness imply strong softness? Does pure noncompactness imply weak softness?

**Example 4.12.** Let  $S$  be a Cu-semigroup. An element  $x \in S$  is *idempotent* if  $2x = x$ . One says that  $S$  is *idempotent* if each of its elements is. It is easy to see that idempotent elements are strongly soft. Thus, every element in an idempotent Cu-semigroup is strongly soft.

Given a  $C^*$ -algebra  $A$ , it was noted in [3, Section 7.2] that  $\text{Cu}(A)$  is idempotent if (and only if)  $A$  is purely infinite. Thus, every element in the Cuntz semigroup of a purely infinite  $C^*$ -algebra is strongly soft.

A  $C^*$ -algebra  $A$  is said to be *weakly purely infinite* if there exists  $n \in \mathbb{N}$  such that  $(n + 1)x = nx$  for every  $x \in \text{Cu}(A)$ . Such elements are functionally soft. Thus, Cuntz semigroups of weakly purely infinite  $C^*$ -algebras satisfy that every element is functionally soft.

It is an open problem if every weakly purely infinite  $C^*$ -algebra is purely infinite; see [27, Question 9.5].

The next result provides a method of building strongly soft elements in Cu-semigroups.

**Lemma 4.13.** *Let  $S$  be a Cu-semigroup, and let  $(y_n)_n$  be a sequence in  $S$  such that  $y_n \leq \infty y_{n+1}$  for each  $n \geq 0$ . Then the element  $\sum_{n=0}^\infty y_n$  is strongly soft.*

**Proof.** To verify that  $y = \sum_{n=0}^\infty y_n$  is strongly soft, let  $y' \in S$  be such that  $y' \ll y$ . It follows that there exists some  $m \in \mathbb{N}$  such that  $y' \ll \sum_{n=0}^m y_n$ .

Using that  $y_n \leq \infty y_{n+1}$  for each  $n$ , we obtain

$$\sum_{n=0}^m y_n \leq \infty y_1 + \sum_{n=1}^m y_n \leq \infty y_2 + \sum_{n=2}^m y_n \leq \dots \leq \infty y_m \leq \infty y_{m+1}$$

and, therefore,  $y' \ll \infty y_{m+1}$ .

Let  $t \in S$  be such that  $t \ll y_{m+1}$  and  $y' \ll \infty t$ . Then,

$$y' + t \ll \left( \sum_{n=0}^m y_n \right) + y_{m+1} \leq y, \quad \text{and} \quad y' \ll \infty t,$$

as desired.  $\square$

**Proposition 4.14.** *Let  $S$  be a Cu-semigroup, and let  $x \in S$ . Then the following are equivalent:*

- (1) *The element  $x$  is strongly soft.*
- (2) *For every  $x' \in S$  satisfying  $x' \ll x$  there exists a strongly soft element  $t \in S$  such that*

$$x' + t \leq x \leq \infty t.$$

- (3) *For every  $x' \in S$  satisfying  $x' \ll x$  there exists  $t \in S$  such that*

$$x' + t \leq x, \quad \text{and} \quad x' \leq \infty t.$$

**Proof.** It is clear that (2) implies (3). To show that (1) implies (2), assume that  $x$  is strongly soft. Choose a  $\ll$ -increasing sequence  $(x_n)_n$  with supremum  $x$  and such that  $x_0 = x'$ . We inductively find  $y_n$  and  $t_n$  such that

$$y_n + t_n \ll y_{n+1} \ll x, \quad y_n \ll \infty t_n, \quad \text{and} \quad x_{n+1} \ll y_{n+1}$$

for all  $n \geq 0$ . We start by setting  $y_0 = x_0$ . Since  $x$  is strongly soft, we obtain  $t_0 \in S$  such that  $y_0 + t_0 \ll x$  and  $y_0 \ll \infty t_0$ . Now let  $n \geq 0$  and assume that we have chosen  $y_k$  and  $t_k$  for all  $k \leq n$ . Then, using that  $y_n + t_n \ll x$  and  $x_{n+1} \ll x$ , we choose  $y_{n+1} \ll x$  such that

$$y_n + t_n \ll y_{n+1}, \quad \text{and} \quad x_{n+1} \ll y_{n+1}.$$

Since  $x$  is strongly soft, we obtain  $t_{n+1} \in S$  such that  $y_{n+1} + t_{n+1} \ll x$  and  $y_{n+1} \ll \infty t_{n+1}$ , which finishes the induction argument.

For each  $n \geq 0$ , we have

$$x_0 + \sum_{k=0}^n t_k = y_0 + t_0 + t_1 + \dots + t_n \leq y_1 + t_1 + \dots + t_n \leq \dots \leq y_{n+1} \leq x.$$

Setting  $t = \sum_{k=0}^{\infty} t_k$ , it is clear that  $x' + t \leq x \leq \infty t$ . Further, it follows from Lemma 4.13 that  $t$  is strongly soft.

To show that (3) implies (1), assume that  $x$  satisfies (3), and let  $x' \in S$  satisfy  $x' \ll x$ . Choose  $x'' \in S$  such that  $x' \ll x'' \ll x$ . Applying (3) to  $x'' \ll x$ , we obtain  $t \in S$  such that

$$x'' + t \leq x, \quad \text{and} \quad x'' \leq \infty t.$$

Using that  $x' \ll x''$ , we can choose  $t' \in S$  such that

$$x' \ll \infty t', \quad \text{and} \quad t' \ll t,$$

which implies that  $x' + t' \ll x$ , as desired.  $\square$

The next result is the analog of [3, Theorem 5.3.11] for strongly soft elements.

**Theorem 4.15.** *Let  $S$  be a Cu-semigroup, and let  $S_{\text{soft}}$  denote the set of strongly soft elements in  $S$ . Then:*

- (1) *The set  $S_{\text{soft}}$  is a submonoid of  $S$  that is closed under suprema of increasing sequences.*
- (2)  *$S_{\text{soft}}$  is absorbing in the following sense: if  $x, y \in S$  satisfy  $x \leq \infty y$  and  $y$  is strongly soft, then so is  $x + y$ .*

**Proof.** To show that  $S_{\text{soft}}$  is closed under addition, let  $x, y \in S_{\text{soft}}$ . To verify that  $x + y$  is strongly soft, let  $w \in S$  satisfy  $w \ll x + y$ . Choose  $x', y' \in S$  such that

$$w \ll x' + y', \quad x' \ll x, \quad \text{and} \quad y' \ll y.$$

Since  $x$  and  $y$  are strongly soft, we obtain  $r, s \in S$  such that

$$x' + r \ll x, \quad x' \ll \infty r, \quad y' + s \ll y, \quad \text{and} \quad y' \ll \infty s.$$

Then

$$w + (r + s) \leq x' + y' + r + s \ll x + y, \quad \text{and} \quad w \leq x' + y' \ll \infty(r + s).$$

Using that 0 is strongly soft, it follows that  $S_{\text{soft}}$  is a submonoid.

To see that  $S_{\text{soft}}$  is closed under suprema of increasing sequences, let  $(x_n)_n$  be an increasing sequence in  $S_{\text{soft}}$ , and set  $x = \sup_n x_n$ . To verify that  $x$  is strongly soft, let  $x' \in S$  satisfy  $x' \ll x$ . We obtain  $n \in \mathbb{N}$  such that  $x' \ll x_n$ . Since  $x_n$  is strongly soft, we get  $r \in S$  such that

$$x' + r \ll x_n, \quad \text{and} \quad x' \ll \infty r.$$

Then  $x' + r \ll x$  and  $x' \ll \infty r$ , as desired.

Let us now prove (2). Thus, let  $x, y \in S$  satisfy  $x \leq \infty y$ , and assume that  $y$  is strongly soft. We verify condition (3) of Proposition 4.14 for  $x + y$ . Let  $w \in S$  satisfy  $w \ll x + y$ , and choose  $y' \in S$  such that

$$w \ll x + y', \quad \text{and} \quad y' \ll y.$$

Since  $y$  is strongly soft, we can apply Proposition 4.14 to obtain  $r \in S$  such that

$$y' + r \leq y \leq \infty r.$$

Using that  $x \leq \infty y$  at the third step, we get

$$w + r \leq x + y' + r \leq x + y \leq \infty y \leq \infty r,$$

as desired.  $\square$

The next result is the Cu-version of Proposition 3.7.

**Lemma 4.16.** *Let  $\varphi: S \rightarrow T$  be a generalized Cu-morphism between Cu-semigroups, and let  $x \in S$ . If  $x$  is (functionally, weakly, strongly) soft, then so is  $\varphi(x)$ .*

**Proof.** Set  $y = \varphi(x)$ . Assuming that  $x$  is strongly soft, we will apply Proposition 4.14(3) to verify that  $y$  is strongly soft. Let  $y \in T$  satisfy  $y' \ll y$ . Using that  $\varphi$  preserves suprema of increasing sequences, we obtain  $x' \in S$  such that  $x' \ll x$  and  $y' \leq \varphi(x')$ . By Proposition 4.14, we obtain  $t \in S$  such that  $x' + t \leq x \leq \infty t$ . Then

$$y' + \varphi(t) \leq \varphi(x') + \varphi(t) = \varphi(x' + t) \leq \varphi(x) = y \leq \varphi(\infty t) = \infty \varphi(t),$$

which shows that  $\varphi(t)$  has the desired properties.

Similarly, one shows that functional and weak softness pass from  $x$  to  $y$ .  $\square$

A projection  $p$  in a  $C^*$ -algebra  $A$  is said to be *properly infinite* if  $p \oplus p \precsim p \oplus 0$  in  $M_2(A)$ . By definition, the zero projection is properly infinite.

Note that, whenever  $A$  is residually stably finite, Proposition 4.17 below says that  $a \in A_+$  is soft if and only if no Cuntz representative of  $a + I$  is a nonzero projection; see [13, Theorem 3.5]. Thus, under these assumptions, one can understand softness as a generalization of pure positivity, as defined in [32, Definition 2.1].

**Proposition 4.17.** *Let  $A$  be a  $C^*$ -algebra, and let  $a \in A_+$ . Consider the following statements:*

- (1) *The element  $a$  is soft.*
- (2) *The class  $[a] \in \text{Cu}(A)$  is strongly soft.*
- (3) *For every closed ideal  $I \subseteq A$ , either the spectrum  $\text{sp}(a + I)$  has 0 as a limit point, or the characteristic function  $\chi$  of  $(0, \infty)$  is continuous on  $\text{sp}(a + I)$  and the corresponding projection  $p = \chi(a + I)$  is properly infinite.*

*Then the implications '(1) $\Rightarrow$ (2) $\Rightarrow$ (3)' hold. If  $A$  is residually stably finite, then the implication '(3) $\Rightarrow$ (1)' holds and so (1)-(3) are equivalent in this case.*

**Proof.** To show that (1) implies (2), set  $x = [a] \in \text{Cu}(A)$ , and let  $x' \in \text{Cu}(A)$  satisfy  $x' \ll x$ . Then, there exists  $\varepsilon > 0$  satisfying  $x' \leq [(a - \varepsilon)_+] \leq x$ . Using Proposition 3.6, we obtain  $b \in \overline{aAa}_+$ , orthogonal to  $(a - \varepsilon)_+$ , such that  $a \in \overline{\text{span}}AbA$ . Setting  $t = [b]$  in  $\text{Cu}(A)$ , one gets

$$x' + t \leq x \leq \infty t.$$

Now it follows from Proposition 4.14 that  $x = [a]$  is strongly soft.

To show that (2) implies (3), let  $I \subseteq A$  be a closed ideal. Then  $\text{Cu}(I)$  is naturally an ideal in  $\text{Cu}(A)$  such that  $\text{Cu}(A/I)$  is canonically isomorphic to  $\text{Cu}(A)/\text{Cu}(I)$ ; see [14] and [3, Section 5.1]. The class  $[a + I] \in \text{Cu}(A/I)$  corresponds to the image of  $[a]$  under the quotient map  $\text{Cu}(A) \rightarrow \text{Cu}(A)/\text{Cu}(I)$ . By Lemma 4.16,  $[a + I]$  is strongly soft.

Thus, without loss of generality, we may assume that  $I = \{0\}$ . We may also assume that 0 is isolated in the spectrum  $\sigma(a)$ , which implies that there exists  $\varepsilon > 0$  with  $\sigma(a) \subseteq \{0\} \cup [\varepsilon, \infty)$ . This implies that  $\chi$  is continuous on  $\sigma(a)$ . Applying functional calculus, we obtain a projection  $p = \chi(a)$ , which satisfies  $[p] = [a]$ . We have  $[p] \ll [p]$ , and therefore  $[p] = 2[p]$  by Proposition 4.8. This implies that  $p \oplus p$  is Cuntz subequivalent, and thus also Murray-von Neumann subequivalent, to  $p$ . Hence,  $p$  is a properly infinite projection.

Finally, if  $A$  is residually stably finite, then no quotient of  $A$  contains a nonzero, properly infinite projection, which shows that (3) implies (1) by Proposition 3.6.  $\square$

**Question 4.18.** Let  $A$  be a stable  $C^*$ -algebra, and let  $x \in \text{Cu}(A)$  be strongly soft. Does there exist a soft element  $a \in A_+$  with  $x = [a]$ ?

## 5. Abundance of soft elements

In this section, we study  $C^*$ -algebras and Cu-semigroups with an *abundance of soft elements*, a notion introduced in Definition 5.2 below. We have shown in Proposition 4.17 that soft elements in  $C^*$ -algebras have a (strongly) soft Cuntz class, and this readily implies that if a stable  $C^*$ -algebra has an abundance of soft elements, then so does the associated Cuntz semigroup. The main result of this section shows that the converse also holds; see Theorem 5.14.

For nonstable  $C^*$ -algebras, we have to consider scaled Cu-semigroups. A *scale* in a Cu-semigroup  $S$  is a subset  $\Sigma \subseteq S$  that is closed under suprema of increasing sequences, that is downward-hereditary, and that generates  $S$  as an ideal; see [6, Definition 4.1]. In this case, the pair  $(S, \Sigma)$  is called a *scaled Cu-semigroup*.

Given a  $C^*$ -algebra  $A$ , the naturally associated scale  $\Sigma_A \subseteq \text{Cu}(A)$  is

$$\Sigma_A = \{x \in \text{Cu}(A) : x \leq [a] \text{ for some } a \in A_+\};$$

see [6, 4.2] and [47, Lemma 3.3(2)].

Given elements  $a$  and  $b$  in a  $C^*$ -algebra, recall that we write  $a \triangleleft b$  if  $a$  belongs to the closed ideal generated by  $b$ . We will use analogous notation in Cu-semigroups.

**Notation 5.1.** Given elements  $x$  and  $y$  in a Cu-semigroup  $S$ , we write  $x \triangleleft y$  if  $x$  belongs to the ideal of  $S$  generated by  $y$ , that is, if  $x \leq \infty y$ .

**Definition 5.2.** We say that a  $C^*$ -algebra  $A$  has an *abundance of soft elements* if for every  $a \in A_+$  and every  $\varepsilon > 0$  there exists a positive, soft element  $b \in \overline{aAa}$  such that  $(a - \varepsilon)_+ \triangleleft b$ .

We say that a scaled Cu-semigroup  $(S, \Sigma)$  has an *abundance of strongly soft elements* if for every  $x', x \in \Sigma$  satisfying  $x' \ll x$  there exists a strongly soft element  $y \in S$  such that  $x' \triangleleft y \leq x$ .

**Remark 5.3.** Let  $S$  be a Cu-semigroup, which we consider as a scaled Cu-semigroup with trivial scale  $\Sigma = S$ . In this case, Definition 5.2 says that  $S$  has an *abundance of strongly soft elements* if for every  $x', x \in S$  satisfying  $x' \ll x$  there exists a strongly soft element  $y \in S$  such that  $x' \triangleleft y \leq x$ .

We start with basic properties of the relation  $\triangleleft$  for Cu-semigroups.

**Lemma 5.4.** Let  $S$  be a Cu-semigroup, and let  $x', x, y \in S$  satisfy  $x' \ll x \triangleleft y$ . Then:

- (1) There exists  $y' \in S$  such that  $x' \triangleleft y' \ll y$ .
- (2) If  $S$  satisfies (O6) and (O7), then there exists  $z \in S$  such that  $z \leq y$  and  $x' \triangleleft z \triangleleft x$ .

**Proof.** Statement (1) is clear. To verify (2), assume that  $S$  satisfies (O6) and (O7). Pick  $x'' \in S$  such that  $x' \ll x'' \ll x$ . Then  $x'' \ll x \leq \infty y$ , and we obtain  $n \in \mathbb{N}$  such that  $x'' \leq ny$ .

It follows from (O6), applied for  $x' \ll x'' \leq ny = y + \dots + y$ , that there exist  $e_1, \dots, e_n \in S$  such that

$$x' \ll e_1 + \dots + e_n, \quad \text{and} \quad e_j \leq x'', y \text{ for } j = 1, \dots, n.$$

Choose  $e'_j \in S$  such that

$$x' \ll e'_1 + \dots + e'_n, \quad \text{and} \quad e'_j \ll e_j \text{ for } j = 1, \dots, n.$$

It follows from (O7), applied for  $e'_j \ll e_j \leq y$ , that there exists  $z \in S$  such that

$$e'_1, \dots, e'_n \leq z \leq y, e_1 + \dots + e_n.$$

Then

$$x' \ll e'_1 + \dots + e'_n \leq nz, \quad \text{and} \quad z \leq e_1 + \dots + e_n \leq nx'',$$

and therefore  $x' \triangleleft z \triangleleft x$ .  $\square$

Next, we show a useful consequence of having an abundance of soft elements in a Cu-semigroup; see Proposition 5.6. In preparation, we first prove the following lemma.

**Lemma 5.5.** *Let  $(S, \Sigma)$  be a scaled Cu-semigroup with an abundance of strongly soft elements. Then for every strongly soft element  $x \in \Sigma$  and every  $x' \in S$  with  $x' \ll x$ , there exists a strongly soft element  $y \in S$  with  $x' \ll y \ll x$ .*

**Proof.** We adapt the proof of [3, Proposition 5.3.18]. Choose  $z', z \in S$  with  $x' \ll z' \ll z \ll x$ . Since  $x$  is strongly soft, we know by Proposition 4.14 that there exists  $t \in S$  such that

$$z + t \leq x \leq \infty t.$$

Take  $t' \in S$  such that

$$t' \ll t, \quad \text{and} \quad z \ll \infty t'.$$

Since  $t \leq x \in \Sigma$ , we have  $t \in \Sigma$ . Using that  $(S, \Sigma)$  has an abundance of soft elements, we obtain a strongly soft element  $u \in S$  such that

$$t' \triangleleft u \ll t.$$

Set  $y = z' + u$ . Then

$$x' \ll z' \leq z' + u = y, \quad \text{and} \quad y = z' + u \ll z + t \leq x.$$

Further, using that  $z' \leq \infty t' \leq \infty u$ , it follows from Theorem 4.15 that  $y = z' + u$  is strongly soft.  $\square$

As defined in [44, Definition 4.1], recall that a submonoid  $T$  of a Cu-semigroup  $S$  is a *sub-Cu-semigroup* of  $S$  if it is closed under suprema of increasing sequences and is such that, for every  $x \in T$  and  $x' \in S$  with  $x' \ll x$ , there exists  $y \in T$  such that  $x' \ll y \ll x$ .

**Proposition 5.6.** *Let  $S$  be a Cu-semigroup with an abundance of strongly soft elements. Then the set  $S_{\text{soft}}$  of strongly soft elements in  $S$  forms a sub-Cu-semigroup of  $S$ .*

**Proof.** By Theorem 4.15,  $S_{\text{soft}}$  is a submonoid of  $S$  that is closed under suprema of increasing sequences. Further, by Lemma 5.5, for all  $x \in S_{\text{soft}}$  and  $x' \in S$  with  $x' \ll x$ , there exists  $y \in S_{\text{soft}}$  such that  $x' \ll y \ll x$ .  $\square$

In [28, Definition 5.1], Kirchberg and Rørdam define a  $C^*$ -algebra  $A$  to have the *2-splitting property* if there exist full elements  $a, b \in A_+$  with  $a \perp b$ . They only apply their definition to unital  $C^*$ -algebras. We propose below a notion that considers the 2-splitting property for all hereditary sub- $C^*$ -algebras, and that also allows for a small ‘error’ (which is only relevant for nonunital subalgebras).

**Definition 5.7.** We say that a  $C^*$ -algebra  $A$  has the *hereditary 2-splitting property* if for every  $a \in A_+$  and every  $\varepsilon > 0$  there exist positive elements  $b, c \in \overline{aAa}$  such that  $b \perp c$  and  $(a - \varepsilon)_+ \triangleleft b, c$ .

We say that a scaled Cu-semigroup  $(S, \Sigma)$  has the *hereditary 2-splitting property* if for every  $x', x \in \Sigma$  satisfying  $x' \ll x$  there exist  $y, z \in S$  such that  $y + z \leq x$ , and  $x' \triangleleft y, z$ .

We first show that in Cu-semigroups satisfying (O5)-(O7) (which always hold in Cuntz semigroups of  $C^*$ -algebras), one can ensure that in the definition of the hereditary 2-splitting property at least one of the ‘splitting’ elements generates an ideal that not only contains  $x'$  but even  $x$ .

**Lemma 5.8.** *Let  $(S, \Sigma)$  be a scaled Cu-semigroup satisfying (O5)-(O7) and with the hereditary 2-splitting property. Let  $x', x \in \Sigma$  be such that  $x' \ll x$ . Then there exist  $y, z \in \Sigma$  such that*

$$y + z \leq x, \quad x' \triangleleft y, \quad \text{and} \quad x \triangleleft z.$$

**Proof.** Choose  $x_1, x_2, x_3 \in S$  such that

$$x' \ll x_1 \ll x_2 \ll x_3 \ll x.$$

Using the hereditary 2-splitting property for  $x_3 \ll x$ , we obtain  $s, t \in S$  such that

$$s + t \leq x, \quad \text{and} \quad x_3 \triangleleft s, t.$$

Applying Lemma 5.4(2) for  $x_1 \ll x_2 \triangleleft s$ , we obtain  $s'$  such that

$$x_1 \triangleleft s' \triangleleft x_2, \quad \text{and} \quad s' \leq s.$$

Applying Lemma 5.4(1) for  $x' \ll x_1 \triangleleft s'$  and for  $x_2 \ll x_3 \triangleleft t$ , we obtain  $s''$  and  $t'$  such that

$$x' \triangleleft s'' \ll s', \quad \text{and} \quad x_2 \triangleleft t' \ll t.$$

Using (O5) for  $s' + t \leq x$ , and  $s'' \ll s'$ , and  $t' \ll t$ , we obtain  $c \in S$  such that

$$s'' + c \leq x \leq s' + c, \quad \text{and} \quad t' \leq c.$$

We have  $s' \triangleleft x_2 \triangleleft t' \leq c$ , and therefore  $x \triangleleft c$ . Further,  $x' \triangleleft s''$ . This shows that  $y = s''$  and  $z = c$  have the desired properties.  $\square$

**Theorem 5.9.** *Let  $(S, \Sigma)$  be a scaled Cu-semigroup satisfying (O5)-(O7). Then the following are equivalent:*

- (1) *For every  $x \in \Sigma$  there exists a strongly soft  $y \in S$  such that  $x \triangleleft y \leq x$ .*
- (2)  *$(S, \Sigma)$  has an abundance of strongly soft elements, that is, for every  $x', x \in \Sigma$  with  $x' \ll x$ , there exists a strongly soft  $y \in S$  with  $x' \triangleleft y \leq x$ .*
- (3)  *$(S, \Sigma)$  has the hereditary 2-splitting property, that is, for every  $x', x \in \Sigma$  with  $x' \ll x$ , there exists  $y, z \in S$  with  $y + z \leq x$  and  $x' \triangleleft y, z$ .*

**Proof.** It is clear that (1) implies (2).

Assume (2), and let  $x', x \in \Sigma$  be such that  $x' \ll x$ . Take  $x'' \in \Sigma$  such that  $x' \ll x'' \ll x$ . Then, there exists a strongly soft element  $y$  with  $x'' \triangleleft y \leq x$ . Find  $y' \in S$  such that  $y' \ll y$  and  $x' \triangleleft y'$ . Using that  $y$  is strongly soft, it follows from Proposition 4.14 that there exists  $z \in S$  such that  $y' + z \leq y \leq \infty z$ . Thus, we get  $y' + z \leq y \leq x$  and  $x' \triangleleft y', z$ , as required.

To prove that (3) implies (1), let  $x \in \Sigma$ . We need to find a strongly soft  $y \in S$  with  $y \leq x \triangleleft y$ .

Choose a  $\ll$ -increasing sequence  $(x_n)_n$  with supremum  $x$ . We will inductively find elements  $z_n \in S$  for  $n \geq 0$  and  $y_n \in S$  for  $n \geq 1$  such that

$$y_{n+1} + z_{n+1} \leq z_n, \quad x_{n+1} \triangleleft y_{n+1}, \quad \text{and} \quad x \triangleleft z_n$$

for all  $n \in \mathbb{N}$ .

To begin, set  $z_0 = x$ . Now, let  $n \geq 0$ , and assume that we have chosen  $z_n$ . Using Lemma 5.4(1) for  $x_{n+1} \ll x \triangleleft z_n$ , we obtain  $z'_n$  such that

$$x_{n+1} \triangleleft z'_n \ll z_n.$$

Applying Lemma 5.8 for  $z'_n \ll z_n$ , we obtain  $y_{n+1}, z_{n+1}$  such that

$$y_{n+1} + z_{n+1} \leq z_n, \quad z'_n \triangleleft y_{n+1}, \quad \text{and} \quad z_n \triangleleft z_{n+1}.$$

Then  $x_{n+1} \triangleleft z'_n \triangleleft y_{n+1}$  and  $x \triangleleft z_n \triangleleft z_{n+1}$ , as desired.

Next, we alter the elements  $y_n$  so that they generate an increasing sequence of ideals. Given  $n \geq 1$ , apply Lemma 5.4(2) for  $x_{n-1} \ll x_n \triangleleft y_n$  to obtain  $y'_n$  such that

$$x_{n-1} \triangleleft y'_n \triangleleft x_n, \quad \text{and} \quad y'_n \leq y_n.$$

Set  $y = \sum_{n=1}^{\infty} y'_n$ . We get  $y'_n \triangleleft x_n \triangleleft y'_{n+1}$  for every  $n \geq 1$ . Therefore,  $y$  is strongly soft by Lemma 4.13. For each  $n \geq 1$ , we have

$$\begin{aligned} y'_1 + \dots + y'_{n-1} + y'_n &\leq y_1 + \dots + y_{n-1} + y_n + z_n \\ &\leq y_1 + \dots + y_{n-1} + z_{n-1} \leq \dots \leq y_1 + z_1 \leq z_0 = x \end{aligned}$$

and therefore  $y = \sup_n \sum_{j=1}^n y'_j \leq x$ . Further, for every  $n \geq 1$ , we have

$$x_n \triangleleft y'_n \leq y,$$

and, consequently,  $x = \sup_n x_n \triangleleft y$ .  $\square$

**Remark 5.10.** In Theorem 5.9, the implications ‘(1)⇒(2)⇒(3)’ hold for arbitrary Cu-semigroups. The assumptions (O5)-(O7) are only needed for the implication ‘(1)⇐(3)’.

The following lemma is the  $C^*$ -analogue of Lemma 4.13.

**Lemma 5.11.** *Let  $A$  be a  $C^*$ -algebra, and let  $(a_n)_n$  be a sequence of positive, contractive, pairwise orthogonal elements in  $A$ . Assume that  $a_n \triangleleft a_{n+1}$  for every  $n \in \mathbb{N}$ . Then  $a = \sum_{n=0}^\infty \frac{1}{2^n} a_n$  is soft.*

**Proof.** We verify condition (3) of Proposition 3.6. Let  $\varepsilon > 0$ . Choose  $N \in \mathbb{N}$  such that  $\frac{1}{2^N} \leq \varepsilon$ . For  $n \geq N$ , we have  $\frac{1}{2^n} a_n - \varepsilon \leq 0$ , and therefore  $(\frac{1}{2^n} a_n - \varepsilon)_+ = 0$ . Since the  $a_n$ ’s are pairwise orthogonal, we have

$$(a - \varepsilon)_+ = \sum_{n=0}^\infty (\frac{1}{2^n} a_n - \varepsilon)_+ = \sum_{n=0}^N (\frac{1}{2^n} a_n - \varepsilon)_+.$$

Set  $b = \sum_{n=N+1}^\infty \frac{1}{2^n} a_n$ . Then  $b \perp (a - \varepsilon)_+$ . For  $n \geq N + 1$ , it is clear that  $a_n \in \overline{\text{span}}AbA$ . For  $n \leq N$ , it follows from the assumption that  $a_n \in \overline{\text{span}}AbA$ . This implies that  $a \in \overline{\text{span}}AbA$ , as desired.  $\square$

The following lemma is the  $C^*$ -analogue of Lemma 5.4.

**Lemma 5.12.** *Let  $A$  be a  $C^*$ -algebra, and let  $a, b \in A_+$  satisfy  $a \triangleleft b$ . Then:*

- (1) *For every  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $(a - \varepsilon)_+ \triangleleft (b - \delta)_+$ .*
- (2) *For every  $\varepsilon > 0$ , there exists  $c \in \overline{bAb}_+$  such that  $(a - \varepsilon)_+ \triangleleft c \triangleleft a$ .*

**Proof.** To prove (1), take  $\varepsilon > 0$ . Using [10, Corollary II.5.2.13], let  $N \in \mathbb{N}$  and  $r_n \in A$  for  $n = 1, \dots, N$  be such that  $\|a - \sum_{n=1}^N r_n^* b r_n\| < \varepsilon$ . Then, there exists  $\delta > 0$  such that

$$\left\| a - \sum_{n=1}^N r_n^* (b - \delta)_+ r_n \right\| < \varepsilon$$

Applying [26, Lemma 2.5(ii)], we obtain an element  $r \in A$  such that  $(a - \varepsilon)_+ = \sum_{n=1}^N (r_n r)^* (b - \delta)_+ (r_n r)$ , as required.

To prove (2), take  $\varepsilon > 0$ . Set  $I = \overline{\text{span}}AaA$ . Then  $\overline{bAb} \cap I$  generates  $I$  as a closed ideal. Thus, using an approximate unit in  $\overline{bAb} \cap I$ , we can find a positive element  $c \in \overline{bAb} \cap I$  such that  $a$  is at distance at most  $\varepsilon$  from the ideal generated by  $c$ . Using a similar argument as in (1), one can show that  $(a - \varepsilon)_+ \triangleleft c$ . Thus,  $c$  has the desired properties.  $\square$

**Lemma 5.13.** *Let  $A$  be a  $C^*$ -algebra, let  $a, c \in A_+$ , let  $\varepsilon > 0$ , and let  $(s_n)_n$  be a sequence of strongly soft elements in  $\text{Cu}(A)$  such that  $[a] \leq \infty s_n$  for all  $n$ , and*

$$\dots \ll s_2 \ll s_1 \ll s_0 \leq [c].$$

*Then there exists a soft element  $b \in \overline{cAc}$  such that  $(a - \varepsilon)_+ \triangleleft b$ .*

**Proof.** For each  $n \in \mathbb{N}$ , using that  $s_{n+1} \ll s_n$  and that  $s_n$  is strongly soft, we obtain  $t_{n+1} \in \text{Cu}(A)$  such that

$$s_{n+1} + t_{n+1} \ll s_n, \quad \text{and} \quad s_{n+1} \ll \infty t_{n+1}.$$

Choose  $s'_n, t'_{n+1} \in \text{Cu}(A)$  such that

$$s'_n \ll s_n, \quad s_{n+1} + t_{n+1} \ll s'_n, \quad t'_{n+1} \ll t_{n+1}, \quad \text{and} \quad s_{n+1} \ll \infty t'_{n+1}.$$

Next, we inductively choose  $c_n \in A_+$  for  $n \in \mathbb{N}$  and  $d_n \in A_+$  for  $n \geq 1$  such that  $c_{n+1}, d_{n+1}$  are orthogonal elements in  $\overline{c_n A c_n}$ , and such that

$$s'_n \leq [c_n] \leq s_n, \quad \text{and} \quad t'_n \leq [d_n] \leq t_n.$$

For  $n = 0$ , set  $c_0 = c$ . Now fix  $n \in \mathbb{N}$  and assume that we have already chosen the elements  $c_0, \dots, c_n \in A_+$  and  $d_1, \dots, d_n \in A_+$ . Then

$$s'_{n+1} \ll s_{n+1}, \quad t'_{n+1} \ll t_{n+1}, \quad \text{and} \quad s_{n+1} + t_{n+1} \leq s'_n \leq [c_n].$$

Applying [36, Lemma 2.3(ii)], we find orthogonal elements  $c_{n+1}$  and  $d_{n+1}$  in  $\overline{c_n A c_n}$  such that

$$s'_{n+1} \ll [c_{n+1}] \ll s_{n+1}, \quad \text{and} \quad t'_{n+1} \ll [d_{n+1}] \ll t_{n+1},$$

which finishes the inductive argument.

Now choose a strictly decreasing sequence  $(\gamma_n)_n$  of positive numbers such that

$$\dots < \gamma_2 < \gamma_1 < \gamma_0 = \varepsilon.$$

Given  $n \geq 1$ , we have

$$[a] \leq \infty s_n \leq \infty t'_n \leq \infty [d_n],$$

and therefore  $(a - \gamma_{n+1})_+ \triangleleft a \triangleleft d_n$ .

Using the equality  $(a - \gamma_n)_+ = ((a - \gamma_{n+1})_+ - (\gamma_n - \gamma_{n+1}))_+$ , and by applying Lemma 5.12 (2), we obtain a positive element  $b_n$  in  $\overline{d_n A d_n}$  such that

$$(a - \gamma_n)_+ \triangleleft b_n \triangleleft (a - \gamma_{n+1})_+.$$

By construction, the elements  $b_1, b_2, \dots$  are pairwise orthogonal, and  $b_n \triangleleft b_{n+1}$  for each  $n \geq 1$ . Thus, the element  $b = \sum_{n=1}^{\infty} \frac{1}{2^k \|b_k\|} b_k$  is soft by Lemma 5.11. Note that, since each  $b_n$  is in  $\overline{c A c}$ , we have  $b \in \overline{c A c}$ . Moreover, one gets

$$(a - \varepsilon)_+ = (a - \gamma_0)_+ \triangleleft b_1 \triangleleft b,$$

as desired.  $\square$

**Theorem 5.14.** *Let  $A$  be a  $C^*$ -algebra. Then the following are equivalent:*

- (1)  *$A$  has an abundance of soft elements.*
- (2)  *$A$  has the hereditary 2-splitting property.*
- (3)  *$(\text{Cu}(A), \Sigma_A)$  has an abundance of strongly soft elements.*
- (4)  *$(\text{Cu}(A), \Sigma_A)$  has the hereditary 2-splitting property.*

**Proof.** To show that (1) implies (2), assume that  $A$  has an abundance of soft elements, and let  $a \in A_+$  and  $\varepsilon > 0$ . We need to find positive elements  $b, c \in \overline{aAa}$  such that  $b \perp c$  and  $(a - \varepsilon)_+ \triangleleft b, c$ . By assumption, there exists a soft element  $d \in \overline{aAa}$  with  $(a - \frac{\varepsilon}{2})_+ \triangleleft d$ . Note that

$$(a - \varepsilon)_+ = ((a - \frac{\varepsilon}{2})_+ - \frac{\varepsilon}{2})_+.$$

We can therefore apply Lemma 5.12(1) to obtain  $\delta > 0$  such that

$$(a - \varepsilon)_+ \triangleleft (d - \delta)_+.$$

Using that  $d$  is soft, we get from Proposition 3.6(3) a positive element  $c \in \overline{dAd}$  with  $c \perp (d - \delta)_+$  and  $d \triangleleft c$ . Set  $b = (d - \delta)_+$ . Then  $b$  and  $c$  have the desired properties.

Let us show that (2) implies (4). Let  $x', x \in \Sigma_A$  with  $x' \ll x$ . We need to find  $y, z \in \text{Cu}(A)$  with  $y + z \leq x$  and  $x' \triangleleft y, z$ .

By [47, Lemma 3.3(1)], we obtain  $a \in A_+$  such that  $x' \ll [a] \ll x$ . Choose  $\varepsilon > 0$  such that  $x' \ll [(a - \varepsilon)_+]$ . By assumption, we obtain orthogonal positive elements  $b, c \in \overline{aAa}$  with  $(a - \varepsilon)_+ \triangleleft b, c$ . Then  $y = [b]$  and  $z = [c]$  have the desired properties.

Since Cuntz semigroups of  $C^*$ -algebras satisfy (O5)-(O7), it follows from Theorem 5.9 that (4) implies (3). Finally, to show that (3) implies (1), assume that  $(\text{Cu}(A), \Sigma_A)$  has an abundance of strongly soft elements and take  $a \in A_+$  and  $\varepsilon > 0$ . We need to find a soft element  $b \in \overline{aAa}$  with  $(a - \varepsilon)_+ \triangleleft b$ .

Using that  $(\text{Cu}(A), \Sigma_A)$  has an abundance of strongly soft elements for  $[a] \in \Sigma_A$  and  $[(a - \frac{\varepsilon}{3})_+] \ll [a]$ , we obtain a strongly soft  $s_0 \in \text{Cu}(A)$  such that

$$[(a - \frac{\varepsilon}{3})_+] \triangleleft s_0 \leq [a].$$

Since  $[(a - \frac{2\varepsilon}{3})_+] \ll [(a - \frac{\varepsilon}{3})_+]$ , we can choose  $s'_0 \in \text{Cu}(A)$  such that

$$[(a - \frac{2\varepsilon}{3})_+] \triangleleft s'_0 \ll s_0.$$

Applying Lemma 5.5 for  $s'_0 \ll s_0$ , we obtain a strongly soft  $s_1 \in \text{Cu}(A)$  with  $s'_0 \ll s_1 \ll s_0$ . Inductively, we obtain a sequence  $(s_n)_n$  of strongly soft elements in  $\text{Cu}(A)$  such that

$$\dots \ll s_2 \ll s_1 \ll s_0 \leq [a],$$

and such that  $[(a - \frac{2\varepsilon}{3})_+] \triangleleft s'_0 \triangleleft s_n$  for all  $n$ .

Applying Lemma 5.13 (with  $\frac{\varepsilon}{3}$ ), we obtain a soft element  $b \in \overline{aAa}$  such that

$$(a - \varepsilon)_+ = (((a - \frac{2\varepsilon}{3})_+ - \frac{\varepsilon}{3})_+) \triangleleft b,$$

as desired.  $\square$

**Corollary 5.15.** *A stable  $C^*$ -algebra has an abundance of soft elements if and only if its Cuntz semigroup has an abundance of strongly soft elements.*

Let  $A$  be a (nonstable)  $C^*$ -algebra. If  $A \otimes \mathbb{K}$  has an abundance of soft elements, then so does  $A$ . Using that  $\text{Cu}(A) \cong \text{Cu}(A \otimes \mathbb{K})$ , we deduce that  $A$  has an abundance of soft elements whenever its Cuntz semigroup does. The converse remains unclear:

**Question 5.16.** Let  $A$  be a  $C^*$ -algebra. If  $A$  has an abundance of soft elements, does its stabilization as well?

With a view towards Theorem 5.14, the above question is equivalent to the following: If  $A$  is a  $C^*$ -algebra such that  $(Cu(A), \Sigma_A)$  has an abundance of strongly soft elements, does  $Cu(A)$  have an abundance of strongly soft elements? One can also ask this question in the more abstract setting of scaled  $Cu$ -semigroups that satisfy additional properties, like (O5)-(O8).

### 6. Completely soft elements in $C^*$ -algebras

We introduce completely soft operators as those positive elements in a  $C^*$ -algebra whose every cut-down is soft; see Definition 6.1. We characterize these elements by spectral properties; see Proposition 6.2. The main result of this section is Theorem 6.10, where we show that a  $C^*$ -algebra has an abundance of such elements whenever it has an abundance of soft elements. To construct completely soft elements in a  $C^*$ -algebra, we follow a strategy introduced in [41] and view positive elements as suitable ‘paths’ of open projections.

**Definition 6.1.** We say that a positive element  $a$  in a  $C^*$ -algebra is *completely soft* if for every  $\varepsilon \geq 0$  the element  $(a - \varepsilon)_+$  is soft.

**Proposition 6.2.** *Let  $A$  be a  $C^*$ -algebra, and  $a \in A_+$ . Then the following are equivalent:*

- (1) *The element  $a$  is completely soft.*
- (2) *For every closed ideal  $I \subseteq A$ , the spectrum of  $a + I \in A/I$  is  $[0, \|a + I\|]$ .*
- (3) *For every closed ideal  $I \subseteq A$ , the spectrum of  $a + I \in A/I$  is connected and contains 0.*

**Proof.** Let  $\varepsilon \geq 0$ , let  $I \subseteq A$  be a closed ideal, and let  $\pi_I: A \rightarrow A/I$  denote the quotient map. Then  $(\pi_I(a) - \varepsilon)_+ = \pi_I((a - \varepsilon)_+)$ . Using the spectral mapping theorem (see, for example, [10, Proposition II.2.3.2]), we see that  $\varepsilon$  is contained in the closure of  $\text{sp}(\pi_I(a)) \cap (\varepsilon, \infty)$  if and only if 0 is not isolated in the spectrum of  $(\pi_I(a) - \varepsilon)_+$ .

By Proposition 3.6,  $a$  is completely soft if and only if for every  $\varepsilon \geq 0$  and every closed ideal  $I \subseteq A$ , the element  $\pi_I((a - \varepsilon)_+)$  is either zero or 0 is not isolated in its spectrum. Using the above considerations, this is in turn equivalent to (2), which is easily seen to also be equivalent to (3).  $\square$

An immediate consequence of Proposition 6.2 is that images of completely soft elements in quotients are again completely soft, which is analogous to Proposition 3.8. It remains unclear if the analog of Proposition 3.7 also holds:

**Question 6.3.** Let  $\varphi: A \rightarrow B$  be a completely positive, order-zero map between  $C^*$ -algebras. Given a completely soft element  $a \in A_+$ , is  $\varphi(a)$  completely soft?

**6.4.** Let  $A$  be a  $C^*$ -algebra. Recall that a projection  $p \in A^{**}$  is said to be *open* if it is the weak\*-limit of an increasing net in  $A_+$ , and denote by  $\mathcal{O}(A)$  the set of all open projections in  $A$ . The sub- $C^*$ -algebra  $pA^{**}p \cap A$  is hereditary for every open projection  $p$  and, conversely, every hereditary sub- $C^*$ -algebra of  $A$  is of this form for some unique open projection of  $A$ ; see [31, p.77f].

As in [41, Section 2], we define the relation  $\prec$  on  $\mathcal{O}(A)$  by setting  $p \prec q$  if there exists  $a \in A_+$  such that  $p \leq a \leq q$ . That is, if there exists  $a \in A_+$  such that  $p = pa$  and  $a = aq$ . We say that  $p \in \mathcal{O}(A)$  is *soft* if the associated hereditary sub- $C^*$ -algebra  $A_p = pA^{**}p \cap A$  is soft. We use  $\text{supp}(a)$  to denote the support projection of any positive element  $a \in A_+$ . Note that  $\text{supp}(a)$  belongs to  $\mathcal{O}(A)$ .

Following [41, Definition 2.4], we say that a map  $f: (-\infty, 0] \rightarrow \mathcal{O}(A)$  is a *path* if  $f(s) = \sup\{f(s') : s' < s\}$  for all  $s \in (-\infty, 0]$ , and if  $f(s) \prec f(t)$  whenever  $s < t$  in  $(-\infty, 0]$ .

Given a positive contraction  $a \in A$ , the map  $f_a: (-\infty, 0] \rightarrow \mathcal{O}(A)$  given by  $f_a(t) = \text{supp}((a+t)_+)$  is a path with  $f_a(-1) = 0$ . Conversely, by [41, Proposition 2.6], every path  $f$  with  $f(-1) = 0$  arises this way from a positive contraction.

**Lemma 6.5.** *Let  $A$  be a  $C^*$ -algebra that has an abundance of soft elements, let  $p, q \in \mathcal{O}(A)$  satisfy  $p \prec q$ , and assume that  $q$  is soft. Then there exists  $r \in \mathcal{O}(A)$  such that  $r$  is soft and  $p \prec r \prec q$ .*

**Proof.** Given any  $\varepsilon > 0$ , let  $f_\varepsilon: [0, \infty) \rightarrow [0, 1]$  be the map that takes the value 0 in  $[0, \varepsilon/2]$ , 1 in  $[\varepsilon, \infty)$ , and is linear from  $\varepsilon/2$  to  $\varepsilon$ .

Since  $p \prec q$ , there exists  $a \in A_+$  such that  $p \leq a \leq q$ . Thus, one has

$$p \leq f_1(a) \ll f_{1/2}(a) \ll f_{1/4}(a) \leq q.$$

Using that  $f_{1/4}(a) \in A_q$ , it follows that there exists an element in  $(A_q)_+$  that acts as a unit for  $f_{1/2}(a)$ . By Proposition 2.9, we obtain a positive element  $d \in A_q$  with  $f_{1/2}(a) \triangleleft d \perp f_{1/2}(a)$ . Applying Lemma 5.12(1), we obtain  $\delta > 0$  such that  $(f_{1/2}(a) - \frac{1}{2})_+ \triangleleft (d - \delta)_+$ . Note that one has  $f_1(a) \triangleleft (f_{1/2}(a) - \frac{1}{2})_+$ .

Applying that  $A$  has an abundance of soft elements for  $(d - \frac{\delta}{2})_+$  and  $\frac{\delta}{2}$ , we obtain a soft element  $e$  such that

$$\left( (d - \frac{\delta}{2})_+ - \frac{\delta}{2} \right)_+ \triangleleft e, \quad \text{and} \quad e \in \overline{(d - \frac{\delta}{2})_+ A (d - \frac{\delta}{2})_+},$$

and we may assume that  $e$  is contractive.

Then, one has

$$f_1(a) \triangleleft (f_{1/2}(a) - \frac{1}{2})_+ \triangleleft (d - \delta)_+ = \left( (d - \frac{\delta}{2})_+ - \frac{\delta}{2} \right)_+ \triangleleft e.$$

Further, using that  $f_1(a) \perp e$  and  $f_1(a) \triangleleft e$ , it follows that  $f_1(a) + e$  is soft and, consequently, that the open projection  $r = \text{supp}(f_1(a) + e)$  is also soft. Since  $f_1(a) + e$  is contractive, we have  $p \leq f_1(a) \leq f_1(a) + e \leq \text{supp}(f_1(a) + e) = r$  and thus  $p \prec r$ .

We also have  $e \in \overline{(d - \frac{\delta}{2})_+ A (d - \frac{\delta}{2})_+}$  and  $(d - \frac{\delta}{2})_+ \ll f_{\delta/2}(d)$ , and therefore  $e \ll f_{\delta/2}(d)$ . We further have  $f_1(a) \ll f_{1/2}(a)$ , and  $f_{1/2}(a) \perp f_{\delta/2}(d)$ . This implies that

$$f_1(a) + e \ll f_{1/2}(a) + f_{\delta/2}(d).$$

Therefore,

$$r = \text{supp}(f_1(a) + e) \leq f_{1/2}(a) + f_{\delta/2}(d) \leq q,$$

and thus  $r \prec q$ .  $\square$

In preparation for the proof of Lemma 6.7, we recall a result of Bice and Koszmider from [9]. Following [9, Definition 2.1], we write  $a \ll_\varepsilon b$  for positive elements  $a$  and  $b$  in a  $C^*$ -algebra and  $\varepsilon > 0$  if  $\|a - ab\| < \varepsilon$ .

**Theorem 6.6** ([9, Corollary 4.2]). *For all  $\varepsilon > 0$  there exists  $\delta > 0$  such that the following holds: If  $a, b, c, d$  are positive contractions in a  $C^*$ -algebra  $A$  such that*

$$a \ll b \ll_\delta c \ll d$$

*then there exists a unitary  $u \in \tilde{A}$  such that*

$$uau^* \ll d, \quad \text{and} \quad \|u - 1\| < \varepsilon.$$

Given open projections  $p, q, r \in \mathcal{O}(A)$  such that  $p, q \prec r$ , note that generally there is no open projection  $r'$  satisfying  $p, q \prec r' \prec r$ . However, using the above result from [9], we prove in Lemma 6.7 below that an approximate version of this statement does hold.

We write  $p \prec_\varepsilon q$  if there exists  $a \in A_+$  such that  $\|p - pa\| < \varepsilon$  and  $a \leq q$ .

**Lemma 6.7.** *Let  $p, q, r$  be open projections in a  $C^*$ -algebra  $A$  such that  $p, q \prec r$ . Take  $\varepsilon > 0$ . Then there exists  $r' \in \mathcal{O}(A)$  such that  $p \prec r' \prec r$  and  $q \prec_\varepsilon r'$ .*

**Proof.** Using functional calculus, we find positive, contractive elements  $a', a, b', b$  in  $A$  such that

$$p \leq a' \ll a \leq r, \quad \text{and} \quad q \leq b' \ll b \leq r.$$

Fix  $\delta > 0$  such that the statement in Theorem 6.6 is satisfied for  $\varepsilon/3$  and  $\delta$ , where note that we may assume  $\delta \leq \varepsilon/6$ . Using functional calculus and an approximate unit in  $A_r$ , we find positive, contractive elements  $c', c \in A$  such that

$$a \ll_\delta c' \ll c, \quad \text{and} \quad b \ll_\delta c' \ll c.$$

Now, by Theorem 6.6, there exists a unitary  $u \in \widetilde{A}_r$  such that

$$ua'u^* \ll c, \quad \text{and} \quad \|1 - u\| < \frac{\varepsilon}{3}.$$

Then

$$a' \ll u^*cu, \quad \text{and} \quad b \ll_\delta c' \ll_{2\varepsilon/3} u^*cu,$$

and therefore

$$\begin{aligned} \|b - b(u^*cu)\| &\leq \|b - bc'\| + \|bc' - bc'(u^*cu)\| + \|bc'(u^*cu) - b(u^*cu)\| \\ &< \delta + \frac{2\varepsilon}{3}\|b\| + \delta\|u^*cu\| \leq \varepsilon, \end{aligned}$$

that is,  $b \ll_\varepsilon u^*cu$ . Now  $r' = \text{supp}(u^*cu)$  has the desired properties.  $\square$

**Lemma 6.8.** *Let  $A$  be a  $C^*$ -algebra and let  $(p_j)_j$  be a  $\prec$ -directed family of soft open projections in  $A$ . Then the open projection  $\sup_j p_j$  is soft.*

**Proof.** Let  $p$  be the supremum of  $(p_j)_j$ . Then,

$$A \cap pA^{**}p = A \cap \overline{\left( \bigcup_j p_j A^{**} p_j \right)}.$$

It follows that each  $A \cap p_j A^{**} p_j$  is a sub- $C^*$ -algebra of  $A \cap pA^{**}p$ , and that the family  $(A \cap p_j A^{**} p_j)_j$  approximates  $A \cap pA^{**}p$ . Since each  $A \cap p_j A^{**} p_j$  is soft, we know from Proposition 2.17 that  $A \cap pA^{**}p$  is soft. By definition,  $p$  is soft.  $\square$

**Theorem 6.9.** *Let  $A$  be a  $C^*$ -algebra that has an abundance of soft elements, and let  $a \in A_+$  be soft. Then there exists a contractive, completely soft element  $b$  such that  $\overline{aAa} = \overline{bAb}$ .*

**Proof.** Set  $p(0) = \text{supp}(a)$ . Let us first prove that there exists a sequence of soft open projections  $p(-\frac{1}{2^n}) \in \mathcal{O}(A)$  such that

$$\text{supp}((a - \frac{1}{2^n})_+) \prec_{1/2^n} p(-\frac{1}{2^n}) \prec p(-\frac{1}{2^{n+1}}) \prec p(0)$$

for each  $n \geq 1$ .

For  $n = 1$ , it follows from Lemma 6.5 applied for  $\text{supp}((a - \frac{1}{2})_+) \prec p(0)$  that one can find a soft open projection  $p(-\frac{1}{2})$  with the required properties.

Now fix  $n \in \mathbb{N}$  and assume that we have found the projections  $p(-\frac{1}{2^k})$  for each  $k \leq n$ . Applying Lemma 6.7 for  $p(-\frac{1}{2^n}), \text{supp}((a - \frac{1}{2^{n+1}})_+) \prec p(0)$  and  $\varepsilon = \frac{1}{2^{n+1}}$  we obtain  $r \in \mathcal{O}(A)$  such that

$$p(-\frac{1}{2^n}) \prec r \prec p(0), \quad \text{and} \quad \text{supp}((a - \frac{1}{2^{n+1}})_+) \prec_{1/2^{n+1}} r.$$

Since  $r \prec p(0)$ , it follows from Lemma 6.5 that there exists a soft open projection  $p(-\frac{1}{2^{n+1}})$  such that  $r \prec p(-\frac{1}{2^{n+1}}) \prec p(0)$ , as desired. In particular, note that  $p(0) = \sup_n p(-\frac{1}{2^n})$ .

Repeated application of the previous construction to each  $p(-\frac{1}{2^n})$  allows us to find soft open projections  $p(t)$  for every dyadic number  $t \in [-1, 0]$  such that  $p(-1) = 0$ , and such that  $p(s) \prec p(t)$  whenever  $s < t$ .

We set  $r(-1) = 0$  and

$$r(t) = \sup \{p(s) : s \in [-1, t] \text{ dyadic}\},$$

for each  $t \in (-1, 0]$ .

Note that

$$r(0) \leq p(0), \quad \text{and} \quad r(0) \geq \sup_n p(-\frac{1}{2^n}) = p(0),$$

which implies  $r(0) = p(0)$ . Further, each  $r(t)$  is soft by Lemma 6.8.

It follows that  $r: [-1, 0] \rightarrow \mathcal{O}(A)$  is a path in the sense of [41, Definition 2.4]. By [41, Proposition 2.6], there is a unique positive element  $b \in A_+$  such that  $\text{supp}((b+t)_+) = r(t)$  for each  $t \in [-1, 0]$ . Since  $r(-1) = 0$ , the element  $b$  is contractive. Further, since  $\text{supp}(b) = r(0) = p(0) = \text{supp}(a)$ , we have  $\overline{aAa} = \overline{bAb}$ .

Finally, each cut-down of  $b$  is soft by construction, as desired.  $\square$

Recall from Definition 5.2 that we say that a  $C^*$ -algebra  $A$  has an *abundance of soft elements* if for every  $a \in A_+$  and every  $\varepsilon > 0$  there exists a positive, soft element  $b \in \overline{aAa}$  such that  $(a - \varepsilon)_+ \triangleleft b$ . One could therefore phrase statement (2) in the next result by saying that ‘ $A$  has an abundance of completely soft elements’.

**Theorem 6.10.** *Let  $A$  be a  $C^*$ -algebra. Then the following are equivalent:*

- (1) *The  $C^*$ -algebra  $A$  has an abundance of soft elements.*
- (2) *For every  $a \in A_+$  and  $\varepsilon > 0$  there exists a positive, completely soft element  $b \in \overline{aAa}$  such that  $(a - \varepsilon)_+ \triangleleft b$ .*

**Proof.** It is clear that (2) implies (1). Conversely, assume that  $A$  has an abundance of soft elements. To verify (2), take  $a \in A_+$  and  $\varepsilon > 0$ . Since  $A$  has an abundance of soft elements, one can find a soft element  $c \in \overline{aAa}$  such that  $(a - \varepsilon)_+ \in \overline{\text{span}AcA}$ . By Theorem 6.9, there exists a completely soft element  $b \in A_+$  such that  $\overline{cAc} = \overline{bAb}$ . This implies that  $b \in \overline{aAa}$  and  $(a - \varepsilon)_+ \in \overline{\text{span}AbA}$ , which shows (2).  $\square$

### 7. Softness and the Global Glimm Property

Following [27, Definition 4.12], we say that a  $C^*$ -algebra  $A$  has the *Global Glimm Property* if for every  $a \in A_+$ , every  $\varepsilon > 0$  and every  $k \geq 2$  there exists a  $*$ -homomorphism  $\varphi: C_0(0, 1] \otimes M_k \rightarrow \overline{aAa}$  such that  $(a - \varepsilon)_+$  is in the ideal generated by the image of  $\varphi$ .

Every  $C^*$ -algebra with the Global Glimm Property has no nonzero elementary ideal-quotients (that is, it is *nowhere scattered* in the sense of [45]). It is an open problem, known as the *Global Glimm Problem*, to determine if every nowhere scattered  $C^*$ -algebra has the Global Glimm Property; see [47].

We shed new light on this problem by proving that the Global Glimm Property implies having an abundance of soft elements, which in turn implies nowhere scatteredness; see Propositions 7.4 and 7.7. This decomposes the Global Glimm Problem into two subproblems; see Question 7.8.

To prove these results, we use that nowhere scatteredness and the Global Glimm Property of a  $C^*$ -algebra  $A$  are reflected in divisibility properties (recalled in Paragraph 7.1) of its Cuntz semigroup  $\text{Cu}(A)$ . More precisely,  $A$  is nowhere scattered if and only if  $\text{Cu}(A)$  is weakly  $(2, \omega)$ -divisible; see [45, Theorem 8.9]. Further,  $A$  has the Global Glimm Property if and only if  $\text{Cu}(A)$  is  $(2, \omega)$ -divisible; see [47, Theorem 3.6].

This translates the Global Glimm Problem into a question about Cuntz semigroups: Does weak  $(2, \omega)$ -divisibility imply  $(2, \omega)$ -divisibility?

In [47] we identified two conditions on a  $\text{Cu}$ -semigroup that precisely capture when the desired implication of divisibility properties holds: ideal-filteredness and property (V). More precisely, a  $\text{Cu}$ -semigroup satisfying (O5)-(O8) is  $(2, \omega)$ -divisible if and only if it is weakly  $(2, \omega)$ -divisible, ideal-filtered and has property (V); see [47, Theorem 6.3]. Below, we give straightforward generalizations of ideal-filteredness and property (V) to scaled  $\text{Cu}$ -semigroups (Definitions 7.9 and 7.10), and we note that the main result of [47] can be generalized to the scaled setting.

We show that an abundance of strongly soft elements implies property (V) (Lemma 7.11), which implies that a  $\text{Cu}$ -semigroup satisfying (O5)-(O8) is  $(2, \omega)$ -divisible if and only if it is ideal-filtered and has an abundance of strongly soft elements; see Theorem 7.12. It follows that a  $C^*$ -algebra has the Global Glimm Property if and only if it has an abundance of soft elements and its scaled Cuntz semigroup is ideal-filtered; see Theorem 7.14.

**7.1.** As defined in [36, Definition 5.1], a  $\text{Cu}$ -semigroup  $S$  is said to be *weakly  $(2, \omega)$ -divisible* if, for every pair  $x' \ll x$  in  $S$ , there exist finitely many elements  $y_1, \dots, y_n \in S$  such that  $x' \leq y_1 + \dots + y_n$  and  $2y_j \leq x$  for each  $j \leq n$ .

Similarly, one says that  $S$  is  *$(2, \omega)$ -divisible* if, whenever  $x' \ll x$  in  $S$ , then there exists  $y \in S$  and  $n \in \mathbb{N}$  such that  $2y \leq x$  and  $x' \leq ny$ .

In [47, Lemma 3.5], we showed that a  $\text{Cu}$ -semigroup satisfying (O5)-(O7) is  $(2, \omega)$ -divisible whenever all elements in a scale are. The next result shows that the analog statement holds for weak  $(2, \omega)$ -divisibility.

**Lemma 7.2.** *Let  $(S, \Sigma)$  be a scaled  $\text{Cu}$ -semigroup satisfying (O6) and assume that every element in  $\Sigma$  is weakly  $(2, \omega)$ -divisible. Then  $S$  is weakly  $(2, \omega)$ -divisible.*

**Proof.** To show that every element in  $S$  is weakly  $(2, \omega)$ -divisible, let  $x', x \in S$  satisfy  $x' \ll x$ . Pick  $x'' \in S$  with  $x' \ll x'' \ll x$ . Since  $\Sigma$  is a scale, there exist  $n$  and  $y_1, \dots, y_n \in \Sigma$  such that  $x'' \leq y_1 + \dots + y_n$ . Applying (O6), we obtain  $z_1, \dots, z_n$  such that

$$x' \ll z_1 + \dots + z_n, \quad \text{and} \quad z_j \leq x \text{ for } j = 1, \dots, n.$$

Pick  $z'_j$  such that

$$x' \ll z'_1 + \dots + z'_n, \quad \text{and} \quad z'_j \ll z_j \text{ for } j = 1, \dots, n.$$

Each  $z_j$  belongs to  $\Sigma$  and is therefore weakly  $(2, \omega)$ -divisible. We thus obtain elements  $v_{j,1}, \dots, v_{j,N_j}$  such that

$$z'_j \leq v_{j,1} + \dots + v_{j,N_j}, \quad \text{and} \quad 2v_{j,k} \leq z_j \text{ for } k = 1, \dots, N_j.$$

Then  $2v_{j,k} \leq z_j \leq x$  for each  $j$  and  $k$ , and also

$$x' \leq \sum_j z'_j \leq \sum_{j,k} v_{j,k}.$$

This shows that the elements  $v_{j,k}$  have the desired properties.  $\square$

Next, we prove that a  $C^*$ -algebra is nowhere scattered whenever it has an abundance of strongly soft elements; see Proposition 7.4 below. In preparation for the proof, we need the following lemma.

**Lemma 7.3.** *Let  $S$  be a Cu-semigroup satisfying (O6), and let  $x \in S$  be weakly soft. Then  $x$  is weakly  $(2, \omega)$ -divisible.*

**Proof.** To verify that  $x$  is weakly  $(2, \omega)$ -divisible, let  $x' \in S$  satisfy  $x' \ll x$ . Choose  $x'' \in S$  such that  $x' \ll x'' \ll x$ . Since  $x$  is weakly soft, we obtain  $n \geq 1$  and  $t_1, \dots, t_n \in S$  such that

$$x'' + t_1 \ll x, \quad \dots, \quad x'' + t_n \ll x, \quad \text{and} \quad x'' \ll t_1 + \dots + t_n.$$

Applying (O6) to  $x' \ll x'' \ll t_1 + \dots + t_n$ , we obtain  $z_1, \dots, z_n \in S$  such that

$$x' \ll z_1 + \dots + z_n, \quad z_1 \ll x'', t_1, \quad \dots, \quad z_n \ll x'', t_n.$$

Then

$$2z_j \ll x'' + t_j \ll x$$

for each  $j = 1, \dots, n$ . Thus,  $z_1, \dots, z_n$  have the desired properties.  $\square$

**Proposition 7.4.** *Let  $(S, \Sigma)$  be a scaled Cu-semigroup satisfying (O6) that has an abundance of strongly soft elements. Then  $S$  is weakly  $(2, \omega)$ -divisible.*

*Consequently, if a  $C^*$ -algebra has an abundance of soft elements, then it is nowhere scattered.*

**Proof.** By Lemma 7.2, it suffices to verify that every element in  $\Sigma$  is weakly  $(2, \omega)$ -divisible. Let  $x', x \in \Sigma$  satisfy  $x' \ll x$ . Pick  $x'', x''' \in S$  with  $x' \ll x'' \ll x''' \ll x$ . Then there exists a strongly soft element  $y \in S$  satisfying  $x''' \triangleleft y \ll x$ . Take  $y' \in S$  with  $y' \ll y$  and  $x'' \triangleleft y'$ . By Lemma 7.3,  $y$  is weakly  $(2, \omega)$ -divisible, which means that we can find  $n$  and  $z_1, \dots, z_n \in S$  such that

$$y' \leq z_1 + \dots + z_n, \quad \text{and} \quad 2z_j \leq y \text{ for } j = 1, \dots, n.$$

Using that  $x' \ll x'' \triangleleft y'$ , we obtain  $N \in \mathbb{N}$  such that  $x' \leq Ny'$ . Then the elements with the desired properties are given by considering  $N$  copies of each of  $z_1, \dots, z_n$ .

For the second part of the statement, let  $A$  be a  $C^*$ -algebra with an abundance of soft elements. By Theorem 5.14, the scaled Cu-semigroup  $(Cu(A), \Sigma_A)$  has an abundance of strongly soft elements. Since Cuntz semigroups of  $C^*$ -algebras always satisfy (O6), we can apply the above argument to deduce that  $Cu(A)$  is weakly  $(2, \omega)$ -divisible. Hence,  $A$  is nowhere scattered by [45, Theorem 8.9].  $\square$

Proposition 7.7 below showcases the exact difference between having an abundance of strongly soft elements and being  $(2, \omega)$ -divisible: In the first case, for every element  $x$  one can find a strongly soft element  $y$  such that  $y \leq x \leq \infty y$  (Theorem 5.9). In the second case,  $y$  can be chosen such that  $2y \leq x \leq \infty y$ .

In the proof of Lemma 7.6 below, we will apply the following useful consequence of (O5):

**Lemma 7.5** ([47, Lemma 2.2]). *Let  $S$  be a Cu-semigroup satisfying (O5), and let  $k \in \mathbb{N}$  and  $x', x, z \in S$  satisfy  $x' \ll x$  and  $(k + 1)x \leq z$ . Then there exists  $y \in S$  such that  $kx' + y \leq z \leq (k + 1)y$  and  $x' \ll y$ .*

**Lemma 7.6.** *Let  $k \in \mathbb{N}$ , and let  $S$  be a  $(2, \omega)$ -divisible Cu-semigroup satisfying (O5). Then, whenever  $(x_n)_n$  is a  $\ll$ -increasing sequence in  $S$ , there exists a sequence  $(y_n)_n$  such that*

$$\sum_{n=0}^{\infty} ky_n \leq \sup_n x_n, \quad \text{and} \quad y_n, x_{n+1} \ll \infty y_{n+1}$$

for every  $n \geq 0$ .

**Proof.** We will use below that  $S$  is  $(k + 1, \omega)$ -divisible by [47, Lemma 3.4].

Let  $(x_n)_n$  be a  $\ll$ -increasing sequence in  $S$ , and let  $x$  be its supremum. Using induction on  $n$ , we will find elements  $y_j, c_j \in S$  for  $j \in \mathbb{N}$  such that

$$ky_{j+1} + c_{j+1} \leq c_j \leq \infty c_{j+1}, \quad y_j \ll c_j, \quad \text{and} \quad y_j, x_{j+1} \ll \infty y_{j+1},$$

for all  $j \geq 0$ , and with  $ky_0 + c_0 \leq x \leq (k + 1)c_0$ .

First, applying that  $S$  is  $(k + 1, \omega)$ -divisible for  $x_0 \ll x$ , we find  $z \in S$  such that

$$(k + 1)z \ll x, \quad \text{and} \quad x_0 \ll \infty z.$$

Choose  $y_0 \in S$  such that

$$y_0 \ll z, \quad \text{and} \quad x_0 \ll \infty y_0.$$

Then, it follows from Lemma 7.5 that there exists  $c_0 \in S$  with

$$ky_0 + c_0 \leq x \leq (k + 1)c_0, \quad \text{and} \quad y_0 \ll c_0,$$

and it is readily checked that  $c_0, y_0$  satisfy the required conditions.

Now fix  $n \in \mathbb{N}$  and assume that we have found the elements  $y_j, c_j$  for every  $j \leq n$ . In particular, we have  $x \leq \infty c_0 \leq \infty c_n$ . Choose  $c'_n \in S$  such that

$$c'_n \ll c_n, \quad y_n \ll c'_n, \quad \text{and} \quad x_{n+1} \ll \infty c'_n.$$

Applying that  $S$  is  $(k + 1, \omega)$ -divisible for  $c'_n \ll c_n$ , one finds  $z_{n+1} \in S$  such that

$$(k + 1)z_{n+1} \ll c_n, \quad \text{and} \quad c'_n \ll \infty z_{n+1}.$$

Choose  $y_{n+1} \in S$  such that

$$y_{n+1} \ll z_{n+1}, \quad \text{and} \quad c'_n \ll \infty y_{n+1}.$$

Applying Lemma 7.5, we find  $c_{n+1} \in S$  with

$$ky_{n+1} + c_{n+1} \leq c_n \leq (k+1)c_{n+1}, \quad \text{and} \quad y_{n+1} \ll c_{n+1},$$

which finishes the inductive argument.

Finally, note that for every  $n \in \mathbb{N}$ , one has

$$\begin{aligned} ky_0 + \dots + ky_n &\leq ky_0 + \dots + ky_n + c_n \leq ky_0 + \dots + ky_{n-1} + c_{n-1} \\ &\leq \dots \leq ky_0 + c_0 \leq x. \end{aligned}$$

Taking supremum on  $n$ , we obtain the required result.  $\square$

**Proposition 7.7.** *Let  $S$  be a  $(2, \omega)$ -divisible Cu-semigroup satisfying (O5). Then for every  $x \in S$  and  $k \in \mathbb{N}$  there exists a strongly soft element  $y \in S$  such that  $ky \leq x \leq \infty y$ . In particular,  $S$  has an abundance of strongly soft elements.*

*Consequently, if a  $C^*$ -algebra has the Global Glimm Property, then it has an abundance of soft elements.*

**Proof.** Let  $x \in S$  and  $k \in \mathbb{N}$ , and let  $(x_n)_n$  be a  $\ll$ -increasing sequence in  $S$  with supremum  $x$ . Using Lemma 7.6, we find a sequence  $(y_n)_n$  such that

$$\sum_{n=0}^{\infty} ky_n \leq x, \quad \text{and} \quad y_n, x_{n+1} \ll \infty y_{n+1}$$

for every  $n \geq 0$ .

Set  $y = \sum_{n=0}^{\infty} y_n$ , which satisfies  $ky \leq x$  by construction. Then, since we have

$$x_n \leq \infty y_n \leq \infty y,$$

for each  $n$ , we deduce  $x \leq \infty y$ . Further,  $y$  is strongly soft by Lemma 4.13.

For the second part of the statement, let  $A$  be a  $C^*$ -algebra with the Global Glimm Property. Then  $\text{Cu}(A)$  is  $(2, \omega)$ -divisible by [47, Theorem 3.6]. Since Cuntz semigroups of  $C^*$ -algebras always satisfy (O5), we can apply the above argument to deduce that  $\text{Cu}(A)$  has an abundance of strongly soft elements. Hence,  $A$  (and in fact, even  $A \otimes \mathbb{K}$ ) has an abundance of soft elements by Theorem 5.14.  $\square$

If a  $C^*$ -algebra has the Global Glimm Property, then it has an abundance of soft elements by Proposition 7.7; and if a  $C^*$ -algebra has an abundance of soft elements, then it is nowhere scattered by Proposition 7.4. Therefore, the Global Glimm Problem, which asks if every nowhere scattered  $C^*$ -algebra has the Global Glimm Property, decomposes into the following subquestions:

**Question 7.8.** If a  $C^*$ -algebra is nowhere scattered, does it have an abundance of soft elements? If a  $C^*$ -algebra has an abundance of soft elements, does it have the Global Glimm Property?

In [47, Definitions 4.1, 5.1], we defined ideal-filteredness and property (V) for Cu-semigroups. We will use the following straightforward generalizations to scaled Cu-semigroups.

**Definition 7.9.** We say that a scaled Cu-semigroup  $(S, \Sigma)$  is *ideal-filtered* if for all  $v', v \in S$  and  $x, y \in \Sigma$  satisfying

$$v' \ll v \ll \infty x, \infty y,$$

there exists  $z \in S$  such that

$$v' \ll \infty z, \quad \text{and} \quad z \ll x, y.$$

**Definition 7.10.** We say that a scaled Cu-semigroup  $(S, \Sigma)$  has *property (V)* if, given  $d'_1, d_1, d'_2, d_2, c, x \in \Sigma$  such that

$$d'_1 \ll d_1, \quad d'_2 \ll d_2, \quad d_1, d_2 \ll c, \quad \text{and} \quad c + d_1, c + d_2 \ll x,$$

there exist  $y, z \in S$  satisfying

$$y + z \leq x, \quad \text{and} \quad d'_1 + d'_2 \leq \infty y, \infty z.$$

**Lemma 7.11.** *Let  $(S, \Sigma)$  be a scaled Cu-semigroup with an abundance of strongly soft elements. Then  $(S, \Sigma)$  has property (V).*

**Proof.** Let  $d'_1, d_1, d'_2, d_2, c, x \in \Sigma$  be such that

$$d'_1 \ll d_1, \quad d'_2 \ll d_2, \quad d_1, d_2 \ll c, \quad \text{and} \quad c + d_1, c + d_2 \ll x.$$

Choose  $x'', x' \in S$  such that

$$x'' \ll x' \ll x, \quad \text{and} \quad c + d_1, c + d_2 \ll x''.$$

Using that  $(S, \Sigma)$  has an abundance of strongly soft elements for  $x' \ll x$ , we find a strongly soft element  $s \in S$  with  $x' \triangleleft s \ll x$ . Choose  $y \in S$  such that  $x'' \triangleleft y \ll s \ll x$ .

Since  $s$  is strongly soft, we know by Proposition 4.14 that there exists  $z \in S$  satisfying  $y + z \leq s \leq \infty z$ . It is readily checked that the elements  $y, z$  have the desired properties.  $\square$

**Theorem 7.12.** *Let  $S$  be a Cu-semigroup satisfying (O5)-(O8). Then the following are equivalent:*

- (1)  $S$  is  $(2, \omega)$ -divisible.
- (2) Every element in  $\Sigma$  is  $(2, \omega)$ -divisible.
- (3) Every element in  $\Sigma$  is weakly  $(2, \omega)$ -divisible and  $(S, \Sigma)$  is ideal-filtered and has property (V).
- (4)  $(S, \Sigma)$  is ideal-filtered and for every  $x$  in  $S$ , there exists a strongly soft element  $y \in S$  such that  $2y \leq x \leq \infty y$ ;
- (5)  $(S, \Sigma)$  is ideal-filtered and has an abundance of strongly soft elements.

**Proof.** By [47, Lemma 3.5], (1) and (2) are equivalent.

The results and proofs of Section 6 in [47] have straightforward generalizations to the scaled version of (weak)  $(2, \omega)$ -divisibility, ideal-filteredness, and property (V). In particular, the scaled version of [47, Theorem 6.3] shows that (2) and (3) are equivalent.

Using Proposition 7.7, we see that (1) implies (4), and it is clear that (4) implies (5). Finally, by Lemma 7.11 and Proposition 7.4, (5) implies (3).  $\square$

**Corollary 7.13.** *Let  $S$  be a Cu-semigroup satisfying (O5)-(O8). Assume that  $S = S_{\text{soft}}$ . Then  $S$  is  $(2, \omega)$ -divisible if and only if  $S$  is ideal-filtered.*

**Theorem 7.14.** *Let  $A$  be a  $C^*$ -algebra. Then the following are equivalent:*

- (1)  *$A$  has the Global Glimm Property.*
- (2)  *$A$  has an abundance of soft elements and  $(\text{Cu}(A), \Sigma_A)$  is ideal-filtered.*
- (3)  *$A$  has an abundance of soft elements and  $\text{Cu}(A)$  is ideal-filtered.*

**Proof.** Let  $A$  be a  $C^*$ -algebra. It follows from [47, Theorem 3.6] that  $A$  has the Global Glimm Property if and only if  $\text{Cu}(A)$  is  $(2, \omega)$ -divisible. Thus, [47, Theorem 6.3] together with Proposition 7.7 show that (1) implies (3), and it is clear that (3) implies (2).

Using Theorem 7.12, we know that  $\text{Cu}(A)$  is  $(2, \omega)$ -divisible if and only if  $\text{Cu}(A)$  has an abundance of strongly soft elements and  $(\text{Cu}(A), \Sigma_A)$  is ideal-filtered. By Theorem 5.14,  $A$  has an abundance of soft elements if and only if strongly soft elements are abundant in  $(\text{Cu}(A), \Sigma_A)$ . This shows that (2) implies (1), as desired.  $\square$

## References

- [1] R. Antoine, F. Perera, L. Robert, H. Thiel, Edwards' condition for quasitraces on  $C^*$ -algebras, Proc. R. Soc. Edinb., Sect. A 151 (2021) 525–547.
- [2] R. Antoine, F. Perera, L. Robert, H. Thiel,  $C^*$ -algebras of stable rank one and their Cuntz semigroups, Duke Math. J. 171 (2022) 33–99.
- [3] R. Antoine, F. Perera, H. Thiel, Tensor products and regularity properties of Cuntz semigroups, Mem. Am. Math. Soc. 251 (2018), viii+191.
- [4] R. Antoine, F. Perera, H. Thiel, Abstract bivariant Cuntz semigroups, Int. Math. Res. Not. (2020) 5342–5386.
- [5] R. Antoine, F. Perera, H. Thiel, Abstract bivariant Cuntz semigroups II, Forum Math. 32 (2020) 45–62.
- [6] R. Antoine, F. Perera, H. Thiel, Cuntz semigroups of ultraproduct  $C^*$ -algebras, J. Lond. Math. Soc. (2) 102 (2020) 994–1029.
- [7] P. Ara, F. Perera, A.S. Toms,  $K$ -theory for operator algebras. Classification of  $C^*$ -algebras, in: Aspects of Operator Algebras and Applications, in: Contemp. Math., vol. 534, Amer. Math. Soc., Providence, RI, 2011, pp. 1–71.
- [8] M.A. Asadi-Vasfi, H. Thiel, E. Vilalta, Ranks of soft operators in nowhere scattered  $C^*$ -algebras, preprint, arXiv:2310.00663 [math.OA], 2023.
- [9] T. Bice, P. Koszmider,  $C^*$ -algebras with and without  $\llcorner$ -increasing approximate units, J. Math. Anal. Appl. 477 (2019) 1396–1418.
- [10] B. Blackadar, Operator algebras, in: Theory of  $C^*$ -Algebras and von Neumann Algebras, Operator Algebras and Non-commutative Geometry, III, in: Encyclopaedia of Mathematical Sciences, vol. 122, Springer-Verlag, Berlin, 2006.
- [11] J. Bosa, Stable elements and property (S), J. Math. Anal. Appl. 512 (2022) 126171.
- [12] J. Bosa, J. Gabe, A. Sims, S. White, The nuclear dimension of  $\mathcal{O}_\infty$ -stable  $C^*$ -algebras, Adv. Math. 401 (2022) 108250.
- [13] N.P. Brown, A. Ciuperca, Isomorphism of Hilbert modules over stably finite  $C^*$ -algebras, J. Funct. Anal. 257 (2009) 332–339.
- [14] A. Ciuperca, L. Robert, L. Santiago, The Cuntz semigroup of ideals and quotients and a generalized Kasparov stabilization theorem, J. Oper. Theory 64 (2010) 155–169.
- [15] K.T. Coward, G.A. Elliott, C. Ivanescu, The Cuntz semigroup as an invariant for  $C^*$ -algebras, J. Reine Angew. Math. 623 (2008) 161–193.
- [16] J. Cuntz, On the continuity of semi-norms on operator algebras, Math. Ann. 220 (1976) 171–183.
- [17] G.A. Elliott, L. Robert, L. Santiago, The cone of lower semicontinuous traces on a  $C^*$ -algebra, Am. J. Math. 133 (2011) 969–1005.
- [18] G.A. Elliott, M. Rørdam, Perturbation of Hausdorff moment sequences, and an application to the theory of  $C^*$ -algebras of real rank zero, in: Operator Algebras: The Abel Symposium 2004, in: Abel Symp., vol. 1, Springer, Berlin, 2006, pp. 97–115.
- [19] C. Farsi, Soft  $C^*$ -algebras, Proc. Edinb. Math. Soc. (2) 45 (2002) 59–65.
- [20] E. Gardella, F. Perera, The modern theory of Cuntz semigroups of  $C^*$ -algebras, preprint, arXiv:2212.02290 [math.OA], 2023.
- [21] E. Gardella, H. Thiel, Weighted homomorphisms between  $C^*$ -algebras, preprint, arXiv:2203.16702 [math.OA], 2022.
- [22] K.R. Goodearl, D.E. Handelman, Stenosis in dimension groups and AF  $C^*$ -algebras, J. Reine Angew. Math. 332 (1982) 1–98.
- [23] I. Hirshberg, M. Rørdam, W. Winter,  $C_0(X)$ -algebras, stability and strongly self-absorbing  $C^*$ -algebras, Math. Ann. 339 (2007) 695–732.
- [24] J.v.B. Hjelmborg, M. Rørdam, On stability of  $C^*$ -algebras, J. Funct. Anal. 155 (1998) 153–170.
- [25] H.E.n. Jensen, Scattered  $C^*$ -algebras, Math. Scand. 41 (1977) 308–314.

- [26] E. Kirchberg, M. Rørdam, Non-simple purely infinite  $C^*$ -algebras, *Am. J. Math.* 122 (2000) 637–666.
- [27] E. Kirchberg, M. Rørdam, Infinite non-simple  $C^*$ -algebras: absorbing the Cuntz algebra  $\mathcal{O}_\infty$ , *Adv. Math.* 167 (2002) 195–264.
- [28] E. Kirchberg, M. Rørdam, When central sequence  $C^*$ -algebras have characters, *Int. J. Math.* 26 (2015) 1550049.
- [29] E. Ortega, F. Perera, M. Rørdam, The corona factorization property, stability, and the Cuntz semigroup of a  $C^*$ -algebra, *Int. Math. Res. Not.* (2012) 34–66.
- [30] E. Ortega, M. Rørdam, H. Thiel, The Cuntz semigroup and comparison of open projections, *J. Funct. Anal.* 260 (2011) 3474–3493.
- [31] G.K. Pedersen,  *$C^*$ -Algebras and Their Automorphism Groups*, Lond. Math. Soc. Monogr., vol. 14, Academic Press, London, 1979.
- [32] F. Perera, A.S. Toms, Recasting the Elliott conjecture, *Math. Ann.* 338 (2007) 669–702.
- [33] L. Robert, Classification of inductive limits of 1-dimensional NCCW complexes, *Adv. Math.* 231 (2012) 2802–2836.
- [34] L. Robert, The cone of functionals on the Cuntz semigroup, *Math. Scand.* 113 (2013) 161–186.
- [35] L. Robert, Remarks on  $\mathcal{Z}$ -stable projectionless  $C^*$ -algebras, *Glasg. Math. J.* 58 (2016) 273–277.
- [36] L. Robert, M. Rørdam, Divisibility properties for  $C^*$ -algebras, *Proc. Lond. Math. Soc.* (3) 106 (2013) 1330–1370.
- [37] L. Robert, A. Tikuisis, Nuclear dimension and  $\mathcal{Z}$ -stability of non-simple  $C^*$ -algebras, *Trans. Am. Math. Soc.* 369 (2017) 4631–4670.
- [38] M. Rørdam, On the structure of simple  $C^*$ -algebras tensored with a UHF-algebra. II, *J. Funct. Anal.* 107 (1992) 255–269.
- [39] M. Rørdam, A simple  $C^*$ -algebra with a finite and an infinite projection, *Acta Math.* 191 (2003) 109–142.
- [40] M. Rørdam, W. Winter, The Jiang-Su algebra revisited, *J. Reine Angew. Math.* 642 (2010) 129–155.
- [41] H. Thiel, Diffuse traces and Haar unitaries, *Amer. J. Math.* (2020) (to appear), arXiv:2009.06940 [math.OA].
- [42] H. Thiel, Ranks of operators in simple  $C^*$ -algebras with stable rank one, *Commun. Math. Phys.* 377 (2020) 37–76.
- [43] H. Thiel, The generator rank of  $C^*$ -algebras, *J. Funct. Anal.* 280 (2021) 108874.
- [44] H. Thiel, E. Vilalta, Covering dimension of Cuntz semigroups II, *Int. J. Math.* 32 (2021), 27 p., Paper No. 2150100.
- [45] H. Thiel, E. Vilalta, Nowhere scattered  $C^*$ -algebras, *J. Noncommut. Geom.* (2023), <https://doi.org/10.4171/JNCG/526>, arXiv:2112.09877 [math.OA].
- [46] H. Thiel, E. Vilalta, Covering dimension of Cuntz semigroups, *Adv. Math.* 394 (2022) 108016.
- [47] H. Thiel, E. Vilalta, The Global Glimm Property, *Trans. Am. Math. Soc.* 376 (2023) 4713–4744.
- [48] A.S. Toms, On the classification problem for nuclear  $C^*$ -algebras, *Ann. Math.* (2) 167 (2008) 1029–1044.
- [49] W. Winter, Nuclear dimension and  $\mathcal{Z}$ -stability of pure  $C^*$ -algebras, *Invent. Math.* 187 (2012) 259–342.
- [50] W. Winter, J. Zacharias, Completely positive maps of order zero, *Münster J. Math.* 2 (2009) 311–324.